



Mathematisches
Forschungsinstitut
Oberwolfach



Oberwolfach Preprints

OWP 2015 - 12

MICHAEL BATE, SEBASTIAN HERPEL, BENJAMIN MARTIN
AND GERHARD RÖHRLE

Cocharacter-Closure and Spherical Buildings

Mathematisches Forschungsinstitut Oberwolfach gGmbH
Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

Starting in 2007, the MFO publishes a preprint series which mainly contains research results related to a longer stay in Oberwolfach. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 30 free hard copies (DIN A4, black and white copy) by surface mail.

Of course, the full copy right is left to the authors. The MFO only needs the right to publish it on its website *www.mfo.de* as a documentation of the research work done at the MFO, which you are accepting by sending us your file.

In case of interest, please send a **pdf file** of your preprint by email to *rip@mfo.de* or *owlf@mfo.de*, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany

Tel +49 7834 979 50
Fax +49 7834 979 55
Email admin@mfo.de
URL www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.
Copyright of the content is held by the authors.

COCHARACTER-CLOSURE AND SPHERICAL BUILDINGS

MICHAEL BATE, SEBASTIAN HERPEL, BENJAMIN MARTIN, AND GERHARD RÖHRLE

In memory of Robert Steinberg

ABSTRACT. Let k be a field, let G be a reductive k -group and V an affine k -variety on which G acts. In this note we continue our study of the notion of cocharacter-closed $G(k)$ -orbits in V . In earlier work we used a rationality condition on the point stabilizer of a G -orbit to prove Galois ascent/descent and Levi ascent/descent results concerning cocharacter-closure for the corresponding $G(k)$ -orbit in V . In the present paper we employ building-theoretic techniques to derive analogous results.

1. INTRODUCTION

Let k be a field and let G be a reductive linear algebraic group acting on an affine variety V , with G , V and the action all defined over k . In this paper we continue the study, initiated in [6], [4] and [1], of the notion of *cocharacter-closed orbits* in V for the group $G(k)$ of k -rational points of G , and of interactions with the geometry of the spherical building $\Delta(\mathbb{K})$ associated to G (for precise definitions, see below). The philosophy of this paper is as follows (cf. [1]): given a point v in V , we are interested in *Galois ascent/descent* questions—given a separable algebraic extension k'/k of fields, how is the $G(k')$ -orbit of v related to the $G(k)$ -orbit of v ?—and *Levi ascent/descent* questions—given a k -defined torus S of the stabilizer G_v , how is the $C_G(S)(k)$ -orbit of v related to the $G(k)$ -orbit of v ? These questions are related, and have natural interpretations in $\Delta(\mathbb{K})$.

Our results complement those of [1]: they give similar conclusions but under different assumptions. It was shown in [1] (see Proposition 2.6 below) that Galois descent—passing from $G(k')$ -orbits to $G(k)$ -orbit—is always well-behaved. Certain results on Galois ascent were also proved [1, Thm. 5.7] under hypotheses on the stabilizer G_v . The mantra in this paper is that when the Centre Conjecture (see Theorem 1.2 below) is known to hold, one can use it to prove Galois ascent results (and hence deduce Levi ascent/descent results). The idea is that when the extension k'/k is separable and normal, these rationality questions can be interpreted in terms of the action of the Galois group of k'/k on the building; moreover, if one has such rationality questions under control, then it is easier to handle Levi ascent/descent because one may assume that the torus S is split [1, Thm. 5.4].

When k is algebraically closed (or more generally when k is perfect), our setup is also intimately related to the *optimality formalism* of Kempf-Rousseau-Hesselink, [11], [17], [9]. Indeed, one may interpret this formalism in the language of the Centre Conjecture (see [4, Sec. 1]). The idea is to study the G -orbits in V via limits along cocharacters of G : limits are formally defined below, but given v in V , if we take the set of cocharacters λ of G for which

2010 *Mathematics Subject Classification.* 20G15 (14L24).

Key words and phrases. Affine G -variety, cocharacter-closed orbit, rationality, spherical building, the Centre Conjecture.

the limit $\lim_{a \rightarrow 0} \lambda(a) \cdot v$ exists, and interpret this set in terms of the set of rational points $\Delta(\mathbb{Q})$ of the building of G , then we obtain a convex subset Σ_v of $\Delta(\mathbb{Q})$. In case $G \cdot v$ is not Zariski-closed, one can find a fixed point in the set Σ_v and an associated *optimal parabolic subgroup* P of G with many nice properties: in particular, the stabilizer G_v normalizes P . It is not currently known in general how to produce analogues of these optimality results over arbitrary fields (or even whether such results exist); see [6, Sec. 1] for further discussion. Our first main theorem gives a rational analogue of the Kempf-Rousseau-Hesselink ideas when Σ_{v, k_s} (the points of Σ_v coming from k_s -defined cocharacters of G) happens to be a subcomplex of $\Delta(\mathbb{Q})$, and also answers in this case the ascent/descent questions posed earlier.

Theorem 1.1. *Let $v \in V$. Suppose Σ_{v, k_s} is a subcomplex of $\Delta_{k_s}(\mathbb{Q})$. Then the following hold:*

- (i) *Suppose $v \in V(k)$ and $G(k_s) \cdot v$ is not cocharacter-closed over k_s . Let S be any k -defined torus of G_v and set $L = C_G(S)$. Then there exists $\sigma \in Y_k(L)$ such that $\lim_{a \rightarrow 0} \sigma(a) \cdot v$ exists and lies outside $G(k_s) \cdot v$.*
- (ii) *Suppose $v \in V(k)$. For any separable algebraic extension k'/k , $G(k') \cdot v$ is cocharacter-closed over k' if and only if $G(k) \cdot v$ is cocharacter-closed over k .*
- (iii) *Let S be any k -defined torus of G_v and set $L = C_G(S)$. Then $G(k) \cdot v$ is cocharacter-closed over k if and only if $L(k) \cdot v$ is cocharacter-closed over k .*

The following result—Tits’ Centre Conjecture—plays a crucial part in the proof of Theorem 1.1:

Theorem 1.2. *Let Δ be a thick building and let Σ be a convex subcomplex of Δ such that Σ is not completely reducible. Then there is a simplex of Δ that is fixed by every building automorphism of Δ that stabilizes Σ . (We call such a simplex a centre of Σ .)*

For definitions and further details, see [16]. The conjecture was proved by Mühlherr-Tits [14], Leeb-Ramos-Cuevas [12] and Ramos-Cuevas [16], and a uniform proof for chamber subcomplexes has also now been given by Mühlherr-Weiss [15]. The hypothesis that Σ_{v, k_s} is a subcomplex allows us to apply the Centre Conjecture. This condition arises naturally in the theory of complete reducibility for subgroups of G and Lie subalgebras of $\text{Lie}(G)$, and our results yield applications to complete reducibility (see Theorem 1.4 below).

It was shown in [1, Thm. 5.7] that the conclusions of Theorem 1.1(ii) and (iii) hold if G_v has a maximal torus that is k -defined. Our second main result gives alternative hypotheses on G_v , this time of a group-theoretic nature, for the conclusions of Theorem 1.1 to hold. The proof of this result also relies in an essential way on known cases of a strengthened version of the Centre Conjecture (this time from [4]).

Theorem 1.3. *Let $v \in V(k)$. Suppose that (a) G_v^0 is nilpotent, or (b) every simple component of G^0 has rank 1. Then the following hold:*

- (i) *Suppose $G(k_s) \cdot v$ is not cocharacter-closed over k_s . Let S be any k -defined torus of G_v and set $L = C_G(S)$. Then there exists $\sigma \in Y_k(L)$ such that $G_v(k_s)$ normalizes $P_\sigma(G^0)$ and $\lim_{a \rightarrow 0} \sigma(a) \cdot v$ exists and lies outside $G(k_s) \cdot v$.*
- (ii) *For any separable algebraic extension k'/k , $G(k') \cdot v$ is cocharacter-closed over k' if and only if $G(k) \cdot v$ is cocharacter-closed over k .*
- (iii) *Let S be any k -defined torus of G_v and set $L = C_G(S)$. Then $G(k) \cdot v$ is cocharacter-closed over k if and only if $L(k) \cdot v$ is cocharacter-closed over k .*

The hypothesis in Theorem 1.1(i) that v is a k -point ensures that the subset Σ_v is Galois-stable, and it is also needed in our proof of Theorem 1.3 (but see Remark 4.6). Sometimes, however, one can get away with a weaker hypothesis. This happens for G -complete reducibility in the final section of the paper, where we prove the following ascent/descent result:

Theorem 1.4. *Suppose that G is connected. Let H be a subgroup of G . Let S be a k -defined torus of $C_G(H)$ and set $L = C_G(S)$. Then H is G -completely reducible over k if and only if H is L -completely reducible over k .*

Remark 1.5. (i) Theorem 1.4 gives an alternative proof and also slightly generalizes Serre's Levi ascent/descent result [18, Prop. 3.2]; cf. [2, Cor. 3.21, Cor. 3.22]. For in the statement of Theorem 1.4, we do not require H to be a subgroup of $G(k)$.
(ii) The counterpart of Theorem 1.1(ii) (Galois descent/ascent for G -complete reducibility) was proved in [3].

We spend much of the paper recalling relevant results from geometric invariant theory and the theory of buildings. Although the basic ideas are familiar, we need to extend many of them: for instance, the material on quasi-states in Section 3.4 was covered in [4] for algebraically closed fields, but we need it for arbitrary fields. We work with the geometric realisations of buildings rather than with buildings as abstract simplicial complexes; some care is needed when the reductive group G has positive-dimensional centre.

The paper is laid out as follows. In Section 2, we set up notation and collect terminology and results relating to cocharacter-closedness. In Section 3, we translate our setup into the language of spherical buildings; we use notation and results from [4] on buildings, some of which we extend slightly. In Section 4, we combine the technology from both of the preceding sections to give proofs of our main results. In the final section we give our applications to the theory of complete reducibility.

2. NOTATION AND PRELIMINARIES

Let k denote a field with separable closure k_s and algebraic closure \bar{k} . Let $\Gamma := \text{Gal}(k_s/k) = \text{Gal}(\bar{k}/k)$ denote the Galois group of k_s/k . Throughout, G denotes a (possibly non-connected) reductive linear algebraic group defined over k , and V denotes a k -defined affine variety upon which G acts k -morphically. Let $G(k)$, $G(k_s)$, $V(k)$, $V(k_s)$ denote the k - and k_s -points of G and V ; we usually identify G with $G(\bar{k})$ and V with $V(\bar{k})$. If X is a variety then we denote its Zariski closure by \overline{X} .

More generally, we need to consider k -points and k_s -points in subgroups that are not necessarily k -defined or k_s -defined; note that if k is not perfect then even when v is a k -point, the stabilizer G_v need not be k -defined. If k'/k is an algebraic field extension and H is a closed subgroup of G then we set $H(k') = H(\bar{k}) \cap G(k')$, and we say that a torus S of H is k' -defined if it is k' -defined as a torus of the k -defined group G . Note that a k_s -defined torus of H is a torus of $\overline{H(k_s)}$.

2.1. Cocharacters and G -actions. Given a k -defined algebraic group H , we let $Y(H)$ denote the set of cocharacters of H , with $Y_k(H)$ and $Y_{k_s}(H)$ denoting the sets of k -defined and k_s -defined cocharacters, respectively. The group H acts on $Y(H)$ via the conjugation action of H on itself. This gives actions of the group of k -points $H(k)$ on $Y_k(H)$ and the group of k_s -points $H(k_s)$ on $Y_{k_s}(H)$. There is also an action of the Galois group Γ on $Y(H)$

which stabilizes $Y_{k_s}(H)$, and the Γ -fixed elements of $Y_{k_s}(H)$ are precisely the elements of $Y_k(H)$. We write $Y = Y(G)$, $Y_k = Y_k(G)$ and $Y_{k_s} = Y_{k_s}(G)$.

Definition 2.1. A function $\|\cdot\| : Y \rightarrow \mathbb{R}_{\geq 0}$ is called a Γ -invariant, G -invariant norm if:

- (i) $\|g \cdot \lambda\| = \|\lambda\| = \|\gamma \cdot \lambda\|$ for all $\lambda \in Y$, $g \in G$ and $\gamma \in \Gamma$;
- (ii) for any maximal torus T of G , there is a positive definite integer-valued form (\cdot, \cdot) on $Y(T)$ such that $\|\lambda, \lambda\| = \|\lambda\|^2$ for any $\lambda \in Y(T)$.

Such a norm always exists. For take a k -defined maximal torus T and any positive definite integer-valued form on $Y(T)$. Since T splits over a finite extension of k , we can average the form over the Weyl group W and over the finite Galois group of the extension to obtain a W -invariant Γ -invariant form on $Y(T)$, which defines a norm satisfying (ii). One can extend this norm to all of Y because any cocharacter is G -conjugate to one in $Y(T)$; this procedure is well-defined since the norm on $Y(T)$ is W -invariant. See [11] for more details. If G is simple then $\|\cdot\|$ is unique up to nonzero scalar multiples. We fix such a norm once and for all.

For each cocharacter $\lambda \in Y$ and each $v \in V$, we define a morphism of varieties $\phi_{v,\lambda} : \bar{k}^* \rightarrow V$ via the formula $\phi_{v,\lambda}(a) = \lambda(a) \cdot v$. If this morphism extends to a morphism $\widehat{\phi}_{v,\lambda} : \bar{k} \rightarrow V$, then we say that $\lim_{a \rightarrow 0} \lambda(a) \cdot v$ exists, and set this limit equal to $\widehat{\phi}_{v,\lambda}(0)$; note that such an extension, if it exists, is necessarily unique.

Definition 2.2. For $\lambda \in Y$ (resp. $\lambda \in Y_k$) and $v \in V$, we say that λ *destabilizes* v provided $\lim_{a \rightarrow 0} \lambda(a) \cdot v$ exists, and if $\lim_{a \rightarrow 0} \lambda(a) \cdot v$ exists and does not belong to $G \cdot v$ (resp. does not belong to $G(k) \cdot v$), then we say λ *properly destabilizes* v (resp. *properly destabilizes* v over k).

2.2. R-parabolic subgroups. When $V = G$ and G is acting by conjugation, for each $\lambda \in Y$ we get a set $P_\lambda := \{g \in G \mid \lim_{a \rightarrow 0} \lambda(a)g\lambda(a)^{-1} \text{ exists}\}$; this is a parabolic subgroup of G . We distinguish these parabolic subgroups by calling them *Richardson-parabolic* or *R-parabolic* subgroups. For basic properties of these subgroups, see [2, Sec. 6]. We recall here that $L_\lambda = C_G(\text{Im}(\lambda))$ is called an *R-Levi subgroup* of P_λ , $R_u(P_\lambda)$ is the set of elements sent to $1 \in G$ in the limit, and $P_\lambda = R_u(P_\lambda) \rtimes L_\lambda$. Further, $R_u(P_\lambda)$ acts simply transitively on the set of all L_μ such that $P_\mu = P_\lambda$ (that is, on the set of all R-Levi subgroups of P_λ): note that this is a transitive action of $R_u(P_\lambda)$ on the set of *subgroups* of the form L_μ , not on the set of cocharacters for which $P_\mu = P_\lambda$. Most of these things work equally well over the field k : for example, if λ is k -defined then P_λ , L_λ and $R_u(P_\lambda)$ are; moreover, given any k -defined R-parabolic subgroup P , $R_u(P)(k)$ acts simply transitively on the set of k -defined R-Levi subgroups of P [6, Lem. 2.5]. Note that if P is k -defined and G is connected then $P = P_\lambda$ for some k -defined λ , but this can fail if G is not connected [6, Sec. 2].

When H is a reductive subgroup of G the inclusion $Y(H) \subseteq Y(G)$ means that we get an R-parabolic subgroup of H and of G attached to any $\lambda \in Y(H)$. When we use the notation P_λ , L_λ , etc., we are always thinking of λ as a cocharacter of G . If we need to restrict attention to the subgroup H for some reason, we write $P_\lambda(H)$, $L_\lambda(H)$, etc.

2.3. The sets $Y(\mathbb{Q})$ and $Y(\mathbb{R})$. Form the set $Y(\mathbb{Q})$ by taking the quotient of $Y \times \mathbb{N}_0$ by the relation $\lambda \sim \mu$ if and only if $n\lambda = m\mu$ for some $m, n \in \mathbb{N}$, and extend the norm function to $Y(\mathbb{Q})$ in the obvious way. For any torus T in G , $Y(T, \mathbb{Q}) := Y(T) \otimes_{\mathbb{Z}} \mathbb{Q}$ is a vector space over \mathbb{Q} . Now one can form real spaces $Y(T, \mathbb{R}) := Y(T, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{R}$ for each maximal torus

T of G and a set $Y(\mathbb{R})$ by glueing the $Y(T, \mathbb{R})$ together according to the way the spaces $Y(T, \mathbb{Q})$ fit together [4, Sec. 2.2]. The norm extends to these sets. One can define sets $Y_k(\mathbb{Q})$, $Y_k(\mathbb{R})$, $Y_k(T, \mathbb{Q})$, $Y_k(T, \mathbb{R})$, $Y_{k_s}(\mathbb{Q})$, $Y_{k_s}(\mathbb{R})$, $Y_{k_s}(T, \mathbb{Q})$ and $Y_{k_s}(T, \mathbb{R})$ analogously by restricting attention to k -defined cocharacters and maximal tori, or k_s -defined cocharacters and maximal tori, as appropriate. For the rest of the paper, \mathbb{K} denotes either of \mathbb{Q} or \mathbb{R} when the distinction is not important. The sets $Y(\mathbb{K})$, $Y_k(\mathbb{K})$ and $Y_{k_s}(\mathbb{K})$ inherit G -, $G(k)$ -, $G(k_s)$ - and Γ -actions from those on Y , Y_k and Y_{k_s} , as appropriate, and each element $\lambda \in Y(\mathbb{K})$ still corresponds to an R-parabolic subgroup P_λ and an R-Levi subgroup L_λ of G (see [4, Sec. 2.2] for the case $\mathbb{K} = \mathbb{R}$). If H is a reductive subgroup of G then we write $Y(H, \mathbb{Q})$, etc., to denote the above constructions for H instead of G .

2.4. G -varieties and cocharacter-closure. We recall the following fundamental definition from [1, Def. 1.2], which extends the one given in [6, Def. 3.8].

Definition 2.3. A subset S of V is said to be *cocharacter-closed over k (for G)* if for every $v \in S$ and $\lambda \in Y_k$ such that $v' := \lim_{a \rightarrow 0} \lambda(a) \cdot v$ exists, we have $v' \in S$.

This notion is explored in detail in [1]. In this section, we content ourselves with collecting some results from that paper, together with the earlier paper [6]. These results, most of which are also needed in the sequel, give a flavour of what is known about the notion of cocharacter-closure in the case that the subset involved is a single $G(k)$ -orbit.

Remark 2.4. The geometric orbit $G \cdot v$ is Zariski-closed if and only if it is cocharacter-closed over \bar{k} , by the Hilbert-Mumford Theorem [11, Thm. 1.4].

The next result is [1, Cor. 5.1].

Theorem 2.5. *Suppose $v \in V$ is such that $G(k) \cdot v$ is cocharacter-closed over k . Then whenever $v' = \lim_{a \rightarrow 0} \lambda(a) \cdot v$ exists for some $\lambda \in Y_k$, there exists $u \in R_u(P_\lambda)(k)$ such that $v' = u \cdot v$.*

The next result is [1, Prop. 5.5].

Proposition 2.6. *Let $v \in V$ such that $G_v(k_s)$ is Γ -stable and let k'/k be a separable algebraic extension. If $G(k') \cdot v$ is cocharacter-closed over k' , then $G(k) \cdot v$ is cocharacter-closed over k .*

The next result is [1, Thm. 5.4].

Theorem 2.7. *Suppose S is a k -defined torus of G_v and set $L = C_G(S)$.*

- (i) *If $G(k) \cdot v$ is cocharacter-closed over k , then $L(k) \cdot v$ is cocharacter-closed over k .*
- (ii) *If S is k -split, then $G(k) \cdot v$ is cocharacter-closed over k if and only if $L(k) \cdot v$ is cocharacter-closed over k .*

We note that, as described in the introduction, one of the main points of this paper is to show that the converse of Proposition 2.6 holds under certain extra hypotheses, and that the hypothesis of splitness can be removed in Theorem 2.7(ii) under the same hypotheses; see also [1, Thm. 1.5].

Our final result is a strengthening of [1, Lem. 5.6].

Lemma 2.8. *Let V be an affine G -variety over k and let $v \in V(k)$. Suppose there exists $\lambda \in Y_{k_s}$ such that λ properly destabilizes v over \bar{k} . Then there exists $\mu \in Y_k$ such that $v' = \lim_{a \rightarrow 0} \mu(a) \cdot v$ exists, v' is not $G(k_s)$ -conjugate to v and $G_v(k_s)$ normalizes P_μ . In particular, $G(k) \cdot v$ is not cocharacter-closed over k .*

Proof. This follows from the arguments of [1, Lem. 5.6]. \square

Remark 2.9. The hypotheses of Lemma 2.8 are satisfied if $\lambda \in Y_{k_s}(Z(G^0))$ destabilizes v but does not fix v . For if $v' := \lim_{a \rightarrow 0} \lambda(a) \cdot v$ is G -conjugate to v then v' is $R_u(P_\lambda)$ -conjugate to v [6, Thm. 3.3]; but $R_u(P_\lambda) = 1$, so this cannot happen.

3. SPHERICAL BUILDINGS AND TITS' CENTRE CONJECTURE

Our aim in this section is to construct objects $\Delta_k(\mathbb{K})$ for $\mathbb{K} = \mathbb{R}$ or \mathbb{Q} which correspond to the geometric realization of the spherical building of G^0 over k (or the rational points thereof) when G^0 is a semisimple group. These are slightly more general objects (possibly with a contribution from $Z(G^0)$) when G^0 is reductive. Recall that Γ denotes the Galois group of k_s/k . Most of the notation and terminology below is developed in full detail in the paper [4]—we point the reader in particular to the constructions in [4, Sec. 2, Sec. 6.3, Sec. 6.4]. For the purposes of this paper, we need to extend some of the results in *loc. cit.* (for example by incorporating the effect of the Galois group Γ), but rather than reiterating all the details, we just gather enough material to make our exposition here coherent.

3.1. Definition of $\Delta_k(\mathbb{K})$. We first form the *vector building* $V_k(\mathbb{K})$ by identifying λ in $Y_k(\mathbb{K})$ with $u \cdot \lambda$ for every $u \in R_u(P_\lambda)(k)$. The restriction of the norm function on $Y_k(\mathbb{K})$ to $V_k(\mathbb{K})$, because it is G -invariant, gives a well-defined function, also called a norm, on $V_k(\mathbb{K})$, and hence turns $V_k(\mathbb{K})$ into a metric space.

Definition 3.1. (i) Define $\Delta_k(\mathbb{R})$ to be the unit sphere in $V_k(\mathbb{R})$ and $\Delta_k(\mathbb{Q})$ to be the projection of $V_k(\mathbb{Q}) \setminus \{0\}$ onto $\Delta_k(\mathbb{R})$.

(ii) Two points of $\Delta_k(\mathbb{K})$ are called *opposite* if they are antipodal on the sphere $\Delta_k(\mathbb{R})$.

(iii) It is clear that the conjugation action of $G(k)$ on Y_k gives rise to an action of $G(k)$ on $\Delta_k(\mathbb{K})$ by isometries, and there is a natural $G(k)$ -equivariant, surjective map $\zeta : Y_k(\mathbb{K}) \setminus \{0\} \rightarrow \Delta_k(\mathbb{K})$.

(iv) The *apartments* of $\Delta_k(\mathbb{K})$ are the sets $\Delta_k(T, \mathbb{K}) := \zeta(Y_k(T, \mathbb{K}))$ where T runs over the maximal k -split tori of G .

(v) The metric space $\Delta_k(\mathbb{K})$ and its apartments have a simplicial structure, because any point of $\Delta_k(\mathbb{K})$ gives rise to a k -defined parabolic subgroup of G^0 ; the simplicial complex consists of the proper k -defined parabolic subgroups of G^0 , ordered by reverse inclusion. We write $\tilde{\Delta}_k$ for the spherical building of G over k regarded purely as a simplicial complex. The simplicial spherical buildings of G^0 and of $[G^0, G^0]$ are the same. Our notion of opposite is compatible with the usual one for parabolic subgroups: if $\lambda \in Y(G)$ then $P_{-\lambda}$ is an opposite parabolic to P_λ .

To avoid tying ourselves in knots, when the distinction is not important to the discussion at hand, we loosely refer to either of the objects $\Delta_k(\mathbb{Q})$ and $\Delta_k(\mathbb{R})$ as *the building of G over k* .

One can make analogous definitions of objects $\Delta_{k_s}(\mathbb{K})$ and $\Delta(\mathbb{K}) = \Delta_{\bar{k}}(\mathbb{K})$ over k_s and \bar{k} , respectively, with corresponding systems of apartments and maps ζ .

Because we are interested in rationality results, we need to know the relationship between $\Delta_k(\mathbb{K})$ and $\Delta_{k_s}(\mathbb{K})$. Given a k -defined reductive subgroup H of G , we also want to relate $\Delta_k(H, \mathbb{K})$ to $\Delta_k(\mathbb{K})$, where $\Delta_k(H, \mathbb{K})$ denotes the building of H over k . It is easy to see that the Γ -action on cocharacters descends (via ζ) to Γ -actions by isometries on $\Delta_{k_s}(\mathbb{K})$ and $\Delta(\mathbb{K})$.

Lemma 3.2. (i) *There are naturally occurring copies of $\Delta_k(\mathbb{K})$ inside $\Delta_{k_s}(\mathbb{K})$ and $\Delta(\mathbb{K})$. We can in fact identify $\Delta_k(\mathbb{K})$ with the set of Γ -fixed points of $\Delta_{k_s}(\mathbb{K})$.*
(ii) *Let H be a k -defined reductive subgroup of G . Then there is a naturally occurring copy of $\Delta_k(H, \mathbb{K})$ inside $\Delta_k(\mathbb{K})$.*

Proof. It is clear that $Y_k(\mathbb{K}) \subseteq Y_{k_s}(\mathbb{K}) \subseteq Y(\mathbb{K})$, and $Y_k(\mathbb{K})$ is precisely the set of Γ -fixed points in $Y_{k_s}(\mathbb{K})$. Since $R_u(P_\lambda)(k)$ acts simply transitively on the set of k -defined R-Levi subgroups of P_λ , two k -defined cocharacters λ and μ are $R_u(P_\lambda)$ -conjugate if and only if they are $R_u(P_\lambda)(k_s)$ -conjugate if and only if they are $R_u(P_\lambda)(k)$ -conjugate. Following this observation through the definition of $\Delta_k(\mathbb{K})$, $\Delta_{k_s}(\mathbb{K})$ and $\Delta_{\bar{k}}(\mathbb{K})$ is enough to prove the first assertion of (i). It is clear that $\Delta_k(\mathbb{K})$ is fixed by Γ . Conversely, let $x \in \Delta_{k_s}(\mathbb{K})$ be fixed by Γ . Let P be the parabolic subgroup associated to x . Then P is k_s -defined and Γ -stable, so P is k -defined. Pick a k -defined maximal torus T of P . There exists $\lambda \in Y_{k_s}(T, \mathbb{K})$ such that $P = P_\lambda$. Each $\gamma \in \Gamma$ maps λ to a $R_u(P)(k_s)$ -conjugate of λ . But T is Γ -stable, so Γ fixes λ . Hence $x \in \Delta_k(\mathbb{K})$, as required.

(ii). In analogy with the first assertion of (i) (although it is slightly more subtle), the key observation is that if $\lambda, \mu \in Y_k(H)$ are $R_u(P_\lambda(G))(k)$ -conjugate, then they are in fact $R_u(P_\lambda(H))(k)$ -conjugate (see [5, Lem. 3.3(i)]). Observe also that the restriction of a Γ - and G -invariant norm on Y to $Y(H)$ gives a Γ - and H -invariant norm on $Y(H)$. \square

Henceforth, we write $\Delta_k(\mathbb{K}) \subseteq \Delta_{k_s}(\mathbb{K}) \subseteq \Delta(\mathbb{K})$ and $\Delta_k(H, \mathbb{K}) \subseteq \Delta_k(\mathbb{K})$ without any further comment. One note of caution: the inclusion $\Delta_k(H, \mathbb{K}) \subseteq \Delta_k(\mathbb{K})$ does not in general respect the simplicial structures on these objects.

3.2. Convex subsets. Because any two parabolic subgroups of G contain a common maximal torus, any two points $x, y \in \Delta(\mathbb{K})$ are contained in a common apartment and, as long as these points are not opposite each other, there is a unique geodesic $[x, y]$ joining them. This geodesic does not depend on the apartment we find containing x and y ; in particular, this can be done inside $\Delta_k(\mathbb{K})$ if $x, y \in \Delta_k(\mathbb{K})$ and inside $\Delta_k(H)$ if $x, y \in \Delta_k(H)$ for some reductive subgroup H of G . This leads to the following key definitions:

Definition 3.3. (i) A subset $\Sigma \subseteq \Delta(\mathbb{K})$ is called *convex* if whenever $x, y \in \Sigma$ are not opposite then $[x, y] \subseteq \Sigma$. It follows from the discussion above that $\Delta_k(\mathbb{K})$ is a convex subset of $\Delta(\mathbb{K})$.
(ii) Given a convex subset Σ of $\Delta(\mathbb{K})$, its preimage $C := \zeta^{-1}(\Sigma) \cup \{0\}$ in $Y(\mathbb{K})$ is a union of cones $C_T := C \cap Y(T, \mathbb{K})$, where T runs over the maximal tori of G . The subset Σ is called *polyhedral* if each C_T is a polyhedral cone and Σ is said to have *finite type* if the set of cones $\{g \cdot C_{g^{-1}Tg} \mid g \in G\}$ is finite for all T .

More generally, there is an addition operation on the set $V(\mathbb{K})$, given as follows. Let $\varphi: Y(\mathbb{K}) \rightarrow V(\mathbb{K})$ be the canonical projection. Choose a maximal torus T of G and $\lambda, \mu \in Y(T, \mathbb{K})$ such that $\varphi(\lambda) = x$ and $\varphi(\mu) = y$; we define $x + y \in V(\mathbb{K})$ by $x + y = \varphi(\lambda + \mu)$.

It can be shown that this does not depend on the choice of T ; moreover, for any $g \in G$, $g \cdot (x + y) = g \cdot x + g \cdot y$.

3.3. The destabilizing locus and complete reducibility. For this paper, a particularly important class of convex subsets arises from G -actions on affine varieties. Given an affine G -variety V and a point $v \in V$, set

$$\Sigma_v := \{\zeta(\lambda) \mid \lambda \in Y \text{ and } \lim_{a \rightarrow 0} \lambda(a) \cdot v \text{ exists}\} \subseteq \Delta(\mathbb{Q}).$$

We call this subset the *destabilizing locus* for v ; it is a convex subset of $\Delta(\mathbb{Q})$ by [4, Lem. 5.5] (note that Σ_v coincides with $E_{V, \{v\}}(\mathbb{Q})$ in the language of [4]). Similarly we write $\Sigma_{v,k}$ (resp. Σ_{v,k_s}) for the image in $\Delta_k(\mathbb{Q})$ (resp. $\Delta_{k_s}(\mathbb{Q})$) of the k -defined (resp. k_s -defined) characters destabilizing v . If H is a reductive subgroup of G , then we write $\Sigma_{v,k}(H)$ for the destabilizing locus for v with respect to H .

Definition 3.4. A subset Σ of $\Delta(\mathbb{K})$ is called *completely reducible* if every point of Σ has an opposite in Σ .

Lemma 3.5. *Let $v \in V$. Then:*

- (i) *Given $\lambda \in Y_k$ such that $\zeta(\lambda) \in \Sigma_{v,k}$, λ has an opposite in $\Sigma_{v,k}$ if and only if there exists $u \in R_u(P_\lambda)(k)$ such that $u \cdot \lambda$ fixes v , if and only if there exists $u \in R_u(P_\lambda)(k)$ such that $\lim_{a \rightarrow 0} \lambda(a) \cdot v = u^{-1} \cdot v$.*
- (ii) *The subset $\Sigma_{v,k}$ is completely reducible if and only if $G(k) \cdot v$ is cocharacter-closed over k .*
- (iii) *The subset Σ_v is completely reducible if and only if the orbit $G \cdot v$ is closed in V .*

Proof. We have that Σ_v (resp. $\Sigma_{v,k}$) is completely reducible if and only if for every $\lambda \in Y$ (resp. $\lambda \in Y_k$) such that $\lim_{a \rightarrow 0} \lambda(a) \cdot v$ exists, there is some $u \in R_u(P_\lambda)$ (resp. $u \in R_u(P_\lambda)(k)$) such that both $u \cdot \lambda$ and $-(u \cdot \lambda)$ destabilize v . But this is true if and only if $u \cdot \lambda$ fixes v , which is equivalent to the fact that $\lim_{a \rightarrow 0} \lambda(a) \cdot v = u^{-1} \cdot v$, by [6, Lem. 2.12]. This gives part (i). Part (ii) now follows from Theorem 2.5, and part (iii) from Remark 2.4. \square

3.4. The strong Centre Conjecture and quasi-states. The aim of the paper [4] is to study a strengthened version of Tits' Centre Conjecture for $\Delta_{k_s}(\mathbb{K})$. Let \mathcal{G} denote the group of transformations of $\Delta(\mathbb{K})$ generated by the isometries arising from the action of G and the action of Γ . Note that elements of \mathcal{G} map k_s -defined parabolic subgroups of G to k_s -defined parabolic subgroups of G , so they give rise to automorphisms of the simplicial building $\tilde{\Delta}_{k_s}$. Given a convex subset Σ of $\Delta(\mathbb{K})$, we call a point of Σ a \mathcal{G} -*centre* if it is fixed by all the elements of \mathcal{G} that stabilize Σ .

We can now formulate the original Centre Conjecture in our setting. Given a convex subset Σ of $\Delta_{k_s}(\mathbb{K})$, we call a point of Σ a \mathcal{G} -*centre* if it is fixed by all the elements of \mathcal{G} that stabilize Σ setwise.

Theorem 3.6. *Suppose $\Sigma \subseteq \Delta(\mathbb{K})$ is a convex non-completely reducible subcomplex. Then Σ has a \mathcal{G} -centre.*

Proof. Theorem 1.2 asserts the existence of a stable simplex in the subcomplex (note that the simplicial structure on $\Delta(\mathbb{K})$ does not “see” the difference between G and G^0 , or between G^0 and its semisimple part, so the proof of the centre conjecture for subcomplexes of spherical buildings still works for the more general class of objects we have described).

Now any element of \mathcal{G} that fixes a simplex also fixes its barycentre (because the action is via isometries), and we are done. \square

In the strong Centre Conjecture [4, Conj. 2.10], one replaces convex non-completely reducible subcomplexes with convex non-completely reducible subsets. Most of [4] deals with the special case when $k = \bar{k}$ and considers only the isometries of $\Delta_k(\mathbb{K})$ coming from the action of G . We need to take the action of Γ into account, so we briefly indicate some of the key changes that must be made to the constructions in [4] in order to make the results go through; see also the comments in [4, Sec. 6.3].

Definition 3.7. We recall the notion of a \mathbb{K} -quasi-state Ξ from [4, Def. 3.1]: this is an assignment of a finite set of characters $\Xi(T)$ to each maximal torus T of G satisfying certain conditions (see *loc. cit.* for a precise statement).

The groups G and Γ act on quasi-states: given a \mathbb{K} -quasi-state Ξ and $g \in G$ and $\gamma \in \Gamma$ we define new quasi-states $g_*\Xi$ and $\gamma_*\Xi$ by

$$g_*\Xi(T) := g! \Xi(g^{-1}Tg), \quad \gamma_*\Xi(T) := \gamma!(\gamma^{-1} \cdot T),$$

where for a character χ of a torus T , $g! \chi$ is a character of the torus gTg^{-1} given by $(g! \chi)(gtg^{-1}) := \chi(t)$ for all $t \in T$, and similarly $\gamma! \chi$ is a character of $\gamma \cdot T$ given by $(\gamma! \chi)(\gamma \cdot t) := \chi(t)$ for all $t \in T$.

We say a quasi-state is *defined over a field k'* if it assigns k' -defined characters to k' -defined maximal tori. Recall also the notions of *boundedness*, *admissibility* and *quasi-admissibility* for \mathbb{K} -quasi-states, [4, Def. 3.1, Def. 3.2].

With these definitions in hand, we can extend [4, Lem. 3.8] as follows:

Lemma 3.8. *Let Υ be a \mathbb{K} -quasi-state which is defined over k_s and define $\Xi := \bigcup_{\gamma \in \Gamma} \gamma_* \Upsilon$ by $\Xi(T) := \bigcup_{\gamma \in \Gamma} (\gamma_* \Upsilon)(T)$ for each maximal torus T of G . Then Ξ is a \mathbb{K} -quasi-state which is defined over k_s , and it is bounded (resp. quasi-admissible, admissible at λ) if Υ is. Moreover, by construction, Ξ is Γ -stable.*

Proof. There are two points which need to be made in order for the arguments already in the proof of [4, Lem. 3.8] to go through. First note that given a k_s -defined maximal torus T of G the set of Galois conjugates of T is finite (because T is defined over some finite extension of k). This means that, because Υ is k_s -defined, $\Xi(T)$ is still finite, so Ξ is a \mathbb{K} -quasi-state. Now, for boundedness we need to check that if Υ is bounded then the set $\bigcup_{\gamma \in \Gamma} \left(\bigcup_{g \in G} g_*(\gamma_* \Upsilon)(T) \right)$ is finite for some (and hence all) k_s -defined maximal tori T of G . Since we can choose any k_s -defined maximal torus T , we choose one that is actually k -defined, and then

$$\begin{aligned} g_*(\gamma_* \Upsilon)(T) &= g!(\gamma_* \Upsilon)(g^{-1}Tg) \\ &= g!(\gamma! \Upsilon(\gamma^{-1} \cdot (g^{-1}Tg))) \\ &= g!(\gamma! \Upsilon((\gamma^{-1} \cdot g)^{-1}T(\gamma^{-1} \cdot g))) \\ &= \gamma!(\gamma^{-1} \cdot g)! \Upsilon((\gamma^{-1} \cdot g)^{-1}T(\gamma^{-1} \cdot g)) \\ &= \gamma!((\gamma^{-1} \cdot g)_* \Upsilon)(T). \end{aligned}$$

Therefore, we can write

$$\bigcup_{\gamma \in \Gamma} \left(\bigcup_{g \in G} (g_*(\gamma_* \Upsilon))(T) \right) = \bigcup_{\gamma \in \Gamma} \gamma! \left(\bigcup_{g \in G} (\gamma^{-1} \cdot g)_* \Upsilon(T) \right).$$

Now, since Υ is bounded, the second union on the RHS is finite for every γ , and because Υ is k_s -defined and the set of Galois conjugates of a k_s -defined character is finite, the whole RHS is finite. This proves the boundedness assertion. The other assertions follow as in *loc. cit.* \square

Using Lemma 3.8 we can ensure that, when appropriate, the states and quasi-states constructed during the course of the paper [4] are Galois-stable. In particular, we get the following variant of [4, Thm. 5.5]:

Theorem 3.9. *Suppose $\Sigma \subseteq \Delta(\mathbb{Q})$ is a convex polyhedral non-completely reducible subset of finite type contained in a single apartment of $\Delta(\mathbb{Q})$. Then Σ has a \mathcal{G} -centre. In particular, if Σ is stabilized by all of Γ , then there exists a Γ -fixed point in Σ .*

Proof. Using Lemma 3.8, one can ensure that the quasi-state constructed in [4, Lem. 5.2] which is used in the proof of [4, Thm. 5.5] is also stable under the relevant elements of \mathcal{G} . The proofs in *loc. cit.* now go through. \square

Remark 3.10. In our application of Theorem 3.9 to the proof of Theorem 1.3 below, Σ is a Γ -stable subset of $\Delta_{k_s}(\mathbb{Q})$ and we want to show that Σ has a Γ -fixed point. A striking feature of Theorem 3.9 is that we do **not** require the apartment containing Σ to be k_s -defined: it can be any apartment of the building $\Delta(\mathbb{K}) = \Delta_{\bar{k}}(\mathbb{K})$.

We note here that, unfortunately, we do not know *a priori* that any cocharacter corresponding to the fixed point given by Theorem 3.9 properly destabilizes v . Moreover, we may need to consider cocharacters of $Z(G^0)$, which do not correspond to simplices of the spherical building at all. These technical issues are at the heart of many of the complications in the proofs in Section 4 below.

4. PROOFS OF THE MAIN RESULTS

Having put in place the building-theoretic technology needed for our proofs in the previous section, we start this section with a few more technical results to be used for the main theorems. As always, V denotes a k -defined affine G -variety, and $v \in V$. One obstacle to proving Theorems 1.1 and 1.3 is that we need to deal with cocharacters that live in $Z(G^0)$, which are not detected by the simplicial building (cf. the proof of Theorem 3.6). An extra problem for Theorem 1.3 is that we need to factor out some simple components of G^0 . The following results let us deal with these difficulties.

Let N be a product of certain simple factors of G^0 , and let S be a torus of $Z(G^0)$. Let M be the product of the remaining simple factors of G^0 together with $Z(G^0)$. Suppose that N and S are normal in G (this implies that M is normal in G as well), and that N and S both fix v . Set $G_1 = G/NS$ and let $\pi: G \rightarrow G_1$ be the canonical projection. Since $Z(G^0)$ is k_s -defined and k_s -split, S is k_s -defined, and it is clear that N is k_s -defined. So G_1 and π are k_s -defined. We have a k_s -defined action of G_1 on the fixed point set V^{NS} (note that V^{NS} is G -stable).

- Lemma 4.1.** (i) For any $\mu_1 \in Y_{k_s}(G_1)$, there exist $n \in \mathbb{N}$ and $\mu \in Y_{k_s}(M)$ such that $\pi \circ \mu = n\mu_1$.
- (ii) Let $\lambda \in Y_{k_s}$. Then λ destabilizes v over k_s in V if and only if $\pi \circ \lambda$ destabilizes v over k_s in V^{NS} . Moreover, if this is the case then $\lim_{a \rightarrow 0} \lambda(a) \cdot v = \lim_{a \rightarrow 0} (\pi \circ \lambda)(a) \cdot v$ belongs to $R_u(P_\lambda(G))(k_s) \cdot v$ if and only if it belongs to $R_u(P_{\pi \circ \lambda}(G_1))(k_s) \cdot v$.
- (iii) $G_1(k_s) \cdot v$ is cocharacter-closed over k_s if and only if $G(k_s) \cdot v$ is cocharacter-closed over k_s .

Proof. (i) Let $\mu_1 \in Y_{k_s}(G_1)$. Since μ_1 is k_s -defined, we can choose a k_s -defined maximal torus $T_1 \subseteq G_1$ with $\mu_1 \in Y_{k_s}(T_1)$. Since π is separable and k_s -defined, $\pi^{-1}(T_1) \subseteq G$ is k_s -defined [20, Cor. 11.2.14]. Hence $\pi^{-1}(T_1)$ has a k_s -defined maximal torus T . Let $T' = T \cap M$, a \bar{k} -defined torus of M . Now $\pi(T') = \pi(T)$ is a maximal torus of G_1 by [7, Prop. 11.14(1)]; but $\pi(T')$ is contained in T_1 , so we must have $\pi(T') = T_1$. The surjection $T' \rightarrow T_1$ induces a surjection $\mathbb{Q} \otimes_{\mathbb{Z}} Y_{\bar{k}}(T') \rightarrow \mathbb{Q} \otimes_{\mathbb{Z}} Y_{\bar{k}}(T_1)$ (the map before tensoring maps onto a finite-index subgroup: e.g., by transposing the injective map on character groups [21, Thm. 7.3]), hence there exist $n \in \mathbb{N}$ and $\mu \in Y_{\bar{k}}(T')$ such that $\pi \circ \mu = n\mu_1$. As $\mu \in Y_{\bar{k}}(T) = Y_{k_s}(T)$, μ is k_s -defined as required.

(ii). The first assertion is immediate, as is the assertion that the limits coincide. Since π is an epimorphism, we have $\pi(R_u(P_\lambda(G))) = R_u(P_{\pi \circ \lambda}(G_1))$ (see [8, Cor. 2.1.9]). Moreover, since λ normalizes NS , the restriction of π to $R_u(P_\lambda(G))$ is separable (see [8, Prop. 2.1.8(3) and Rem. 2.1.11]) and k_s -defined, and hence is surjective on k_s -points (cf. [21, Cor. 18.5]). This implies that if the common limit v' is in $R_u(P_{\pi \circ \lambda}(G_1))(k_s) \cdot v$, it is contained in $R_u(P_\lambda(G))(k_s) \cdot v$. The reverse implication is clear, since π is k_s -defined.

(iii). Suppose $G(k_s) \cdot v$ is not cocharacter-closed over k_s . Then there exists $\lambda \in Y_{k_s}$ such that $\lim_{a \rightarrow 0} \lambda(a) \cdot v$ exists but does not belong to $R_u(P_\lambda(G))(k_s) \cdot v$. Then $\lim_{a \rightarrow 0} (\pi \circ \lambda)(a) \cdot v$ exists but does not belong to $R_u(P_\lambda(G_1))(k_s) \cdot v$, by part (ii). Hence $G_1(k_s) \cdot v$ is not cocharacter-closed over k_s , by Theorem 2.5.

Now suppose $G_1(k_s) \cdot v$ is not cocharacter-closed over k_s . Then there exists $\mu_1 \in Y_{k_s}(G_1)$ such that $v' := \lim_{a \rightarrow 0} \mu_1(a) \cdot v$ exists and does not belong to $R_u(P_{\mu_1}(G_1))(k_s) \cdot v$. Replacing μ_1 with a positive multiple $n\mu_1$ of μ_1 if necessary, it follows from part (i) that there exists $\mu \in Y_{k_s}$ such that $\pi \circ \mu = \mu_1$. Then $\lim_{a \rightarrow 0} \mu(a) \cdot v$ is equal to v' and v' does not belong to $R_u(P_\lambda(G))(k_s) \cdot v$, by part (ii). Hence $G(k_s) \cdot v$ is not cocharacter-closed over k_s , by Theorem 2.5. \square

Remark 4.2. We insist in Lemma 4.1(i) that λ be a cocharacter of M because we need this in the proof of Theorem 1.3.

Lemma 4.3. Let G be connected, let S be a k_s -torus of G_v and set $L = C_G(S)$. Suppose that for every $\lambda, \mu \in Y_{k_s}$ such that $P_\lambda = P_\mu$, either both of λ and μ destabilize v or neither does. Then for every $\lambda, \mu \in Y_{k_s}(L)$ such that $P_\lambda(L) = P_\mu(L)$, either both of λ and μ destabilize v or neither does.

Proof. Let $\lambda, \mu \in Y_k(L)$ such that $P_\lambda(L) = P_\mu(L)$. We can choose $\sigma \in Y_k(S)$ such that $L_\sigma = L$ [1, Lem. 2.5]. Then $P_\lambda(L_\sigma) = P_\mu(L_\sigma)$ and there exists $n \in \mathbb{N}$ such that $P_{n\sigma + \lambda} = P_\lambda(L_\sigma)R_u(P_\sigma)$ and $P_{n\sigma + \mu} = P_\mu(L_\sigma)R_u(P_\sigma)$ [2, Lem. 6.2(i)]. By hypothesis, either both of $n\sigma + \lambda$ and $n\sigma + \mu$ destabilize v , or neither one does. In the first case, since σ fixes v , both λ and μ must destabilize v . Conversely, in the second case neither λ nor μ can destabilize v . \square

Lemma 4.4. *Let T be a maximal torus of G , let $\mu_1, \dots, \mu_r \in Y(T) \setminus \{0\}$, let $\mu = \sum_{i=1}^r \mu_i$ and assume $\mu \neq 0$. Suppose $g \in G$ and $g \cdot \zeta(\mu_i) = \zeta(\mu_i)$ for all $1 \leq i \leq r$. Then $g \cdot \zeta(\mu) = \zeta(\mu)$.*

Proof. Clearly, there is a permutation $\tau \in S_r$ such that none of the sums $\sum_{i=1}^t \mu_{\tau(i)}$ is 0 for $1 \leq t \leq r$. Consider the special case $r = 2$ (the general case follows easily by induction on r). Recall the addition operation $+$ on $V(\mathbb{K})$ and the canonical projection $\varphi: Y(\mathbb{K}) \rightarrow V(\mathbb{K})$ from Section 3.2. Let $\xi: V(\mathbb{K}) \setminus \{0\} \rightarrow \Delta(\mathbb{K})$ be the canonical projection. Note that φ and ξ are G -equivariant. Moreover, as g fixes $\zeta(\mu_1)$ and g acts as an isometry, g fixes $\varphi(\mu_1)$, and likewise g fixes $\varphi(\mu_2)$. We have

$$\begin{aligned} g \cdot \zeta(\mu) &= g \cdot \zeta(\mu_1 + \mu_2) = g \cdot \xi(\varphi(\mu_1 + \mu_2)) = \xi(g \cdot \varphi(\mu_1 + \mu_2)) = \xi(g \cdot (\varphi(\mu_1) + \varphi(\mu_2))) \\ &= \xi(g \cdot \varphi(\mu_1) + g \cdot \varphi(\mu_2)) = \xi(\varphi(\mu_1) + \varphi(\mu_2)) = \xi(\varphi(\mu_1 + \mu_2)) = \zeta(\mu_1 + \mu_2) = \zeta(\mu), \end{aligned}$$

as required. \square

We now have everything in place to prove Theorem 1.1.

Proof of Theorem 1.1. For part (i), suppose $v \in V(k)$ and $G(k_s) \cdot v$ is not cocharacter-closed over k_s . Clearly there is no harm in assuming S is a maximal k -defined torus of G_v , so we shall do this. Since the closed subgroup $\overline{G_v(k_s)}$ generated by $G_v(k_s)$ is k_s -defined and Γ -stable, it is k -defined. Hence S is a maximal torus of $\overline{G_v(k_s)}$; in particular, S is a maximal k_s -defined torus of G_v . Set $H = C_G(S)$. If $\sigma \in Y_k(H)$ and σ destabilizes v but does not fix v then σ properly destabilizes v over k_s for G , by [1, Lem. 2.8]. Hence it is enough to prove that such a σ exists.

By Theorem 2.7(ii), $H(k_s) \cdot v$ is not cocharacter-closed over k_s . So we can choose $\mu \in Y_{k_s}(H)$ such that μ properly destabilizes v over k_s for H . If $\mu \in Y_{k_s}(Z(H^0))$ then we are done by Lemma 2.8 and Remark 2.9. So assume otherwise. Then $P_\mu(H^0)$ is a proper subgroup of H^0 . By Lemma 4.3, $\Sigma_{v,k_s}(H)$ is a subcomplex of $\Delta_{k_s}(H, \mathbb{K})$. It follows from Lemma 3.5 that $\tilde{\Sigma}_{v,k_s}(H)$ (the image of $\Sigma_{v,k_s}(H)$ in the simplicial building $\tilde{\Delta}_{k_s}(H)$) is not completely reducible, since if Q is an opposite parabolic to $P_\mu(H^0)$ in H^0 then there exists $\mu' \in Y_{k_s}(H)$ such that $P_{\mu'} = Q$ and μ' is opposite to μ , which is impossible. Clearly, $\tilde{\Sigma}_{v,k_s}(H)$ is Γ - and $H_v(k_s)$ -stable, so by Theorem 3.6 there is a Γ - and $H_v(k_s)$ -fixed simplex $s \in \tilde{\Sigma}_{v,k_s}(H)$, corresponding to some proper parabolic subgroup P of H^0 . There exists $\sigma \in Y_k(H)$ such that $P = P_\sigma(H^0)$ [6, Lem. 2.5(ii)], and σ destabilizes v by construction. Now $\sigma \notin Y_{k_s}(Z(H^0))$ since P is proper. But every k_s -defined torus of $H_v(k_s)$ is contained in $Z(H^0)$ (since S is contained in $Z(H^0)$), so σ does not fix v . As σ commutes with S , it follows from [1, Lem. 2.8] that $v' := \lim_{a \rightarrow 0} \sigma(a) \cdot v$ does not lie in $H(k_s) \cdot v$. This completes the proof of (i).

For part (ii), Proposition 2.6 shows that if $G(k') \cdot v$ is cocharacter-closed over k' then $G(k) \cdot v$ is cocharacter-closed over k . For the other direction, suppose that $G(k') \cdot v$ is not cocharacter-closed over k' . Again by Proposition 2.6, we may assume $k' = k_s$. Applying part (i) with $S = 1$, we find $\sigma \in Y_k$ such that σ properly destabilizes v over k_s . In particular, $G(k) \cdot v$ is not cocharacter-closed over k . This finishes part (ii).

Part (iii) of Theorem 1.1 follows using similar arguments to those in the proof of [1, Thm. 5.7(ii)]. Let S be a k -defined torus of G_v and let $L = C_G(S)$. First, by the argument of [1, Lem. 6.2], we can assume without loss that $v \in V(k_s)$ without changing Σ_{v,k_s} . Second, by [1, Lem. 6.3] and the argument of the proof of [1, Thm. 6.1], we can pass to a suitable G -variety W and find $w \in W(k)$ such that $\Sigma_{w,k_s} = \bigcap_{\gamma \in \Gamma} \gamma \cdot \Sigma_{v,k_s}$; in particular, Σ_{w,k_s} is a subcomplex

of $\Delta_k(\mathbb{Q})$ and $\Sigma_{w,k} = \Sigma_{v,k}$. The arguments of [1, Sec. 6] also show that S fixes w . Hence we can assume without loss that $v \in V(k)$. As before, Lemma 4.3 implies that $\Sigma_{v,k_s}(L)$ is a subcomplex of $\Delta_{k_s}(L, \mathbb{K})$. We may thus apply part (ii) and assume $k = k_s$. But then S is k -split, so the result follows from Theorem 2.7. \square

Remark 4.5. We do not know how to prove that $P_\sigma(G^0)$ from Theorem 1.1(i) is normalized by $G_v(k_s)$, but the proof does show that $P_\sigma(G^0)$ is normalized by $H_v(k_s)$.

We finish the section by proving Theorem 1.3.

Proof of Theorem 1.3. For part (i), suppose $v \in V(k)$ and $G(k_s) \cdot v$ is not cocharacter-closed over k_s . Recall that a connected algebraic group is nilpotent if and only if it contains just one maximal torus (see [10, 21.4 Prop. B]). Let G_i be a simple component of G^0 . If $\text{rank}(G_i) = 1$ and $\dim(G_i)_v \geq 2$ then $(G_i)_v^0$ must contain a Borel subgroup B_i of G_i : but then the orbit map $G_i \rightarrow G_i \cdot v$ factors through the connected projective variety G_i/B_i and hence is constant, so $(G_i)_v = G_i$.

Let N be the product of the simple components of G^0 that fix v , and let K be the product of the remaining simple components of G^0 together with $Z(G^0)^0$. Then N and K are Γ -stable, so they are k -defined. The next step is to factor out N and reduce to the case when the stabilizer has nilpotent identity component. As in the proof of Theorem 1.1, $\overline{G_v(k_s)}$ is k -defined and we may assume S is a maximal k -defined torus of G_v and a k_s -defined maximal torus of $\overline{G_v(k_s)}$. We can choose k -defined tori S_0 of K and S_2 of N such that $S = S_0 S_2$. Note that K_v^0 is nilpotent—this holds by assumption in case (a), and by the above argument in case (b)—so S_0 is the unique maximal k -defined torus of $\overline{K_v(k_s)}$.

Let $H_0 = N_G(S_0)$, let $H = N_{H_0}(N)$ and let $M = C_K(S_0)$. Then H_0 is k -defined [8, Lem. A.2.9], so H is k -defined as it is Γ -stable and has finite index in H^0 . Note that $H^0 = NM = C_G(S_0)^0$, so N is a product of simple components of H^0 and M is the product of $Z(H^0)^0 = S^0 Z(G^0)^0$ with the remaining simple components of H^0 . The subgroup M of H is normal, so it is Γ -stable and hence k -defined. Now M_v^0 is nilpotent since K_v^0 is, so M_v^0 has a unique maximal torus S' —in particular, $S_0 \subseteq S'$ and S_0 is the unique maximal torus of $\overline{M_v(k_s)}$. Since $G_v(k_s)$ normalizes N and K , $G_v(k_s)$ normalizes N and S_0 , so $G_v(k_s) \subseteq H$; it follows that $H_v(k_s) = G_v(k_s)$.

Let $H_1 = H/N S_0$ and let $\pi: H \rightarrow H_1$ be the canonical projection. We wish to find $\lambda_1 \in Y_{k_s}(H_1)$ such that λ_1 properly destabilizes v over k_s and $P_{\lambda_1}(H_1)$ is k -defined. Note that no nontrivial k_s -defined cocharacter of H_1 fixes v ; for if $0 \neq \lambda_1 \in Y_{k_s}(H_1)$ fixes v then by Lemma 4.1, there exist $n \in \mathbb{N}$ and $\lambda \in Y_{k_s}(M)$ such that $\pi \circ \lambda = n\lambda_1$, and $\langle \text{Im}(\lambda) \cup S \rangle$ is a k_s -defined torus of $\overline{G_v(k_s)}$ that properly contains S , contradicting the maximality of S . Clearly, $(H_1)_v^0$ is nilpotent with unique maximal torus $S'_1 := \pi(S')$. Since $H^0 = C_G(S_0)^0$, $H(k_s) \cdot v$ is not cocharacter-closed over k_s , by Theorem 2.7(ii) and [1, Cor. 5.3]. Hence $H_1(k_s) \cdot v$ is not cocharacter-closed over k_s (Lemma 4.1). Let $\lambda_1 \in Y_{k_s}(H_1)$ such that λ_1 destabilizes v . By Lemmas 2.8 and 4.1, we can assume λ_1 does not properly destabilize v over \bar{k} . Therefore, there exists $u \in R_u(P_{\lambda_1}(H_1))$ such that $u \cdot \lambda_1$ fixes v ; then $u \cdot \lambda_1$ must be a cocharacter of S'_1 . It follows that $\Sigma_{v,k_s}(H_1) \subseteq \Delta_{k_s}(T_1, \mathbb{Q})$, where T_1 is a fixed maximal torus of H_1 that contains S'_1 . Note that T_1 and S'_1 need not be k -defined, or even k_s -defined. As $H_1(k_s) \cdot v$ is not cocharacter-closed over k_s , $\Sigma_{v,k_s}(H_1)$ is not completely reducible (Lemma 3.5(ii)). Now $\Sigma_{v,k_s}(H_1)$ is stabilized by Γ and by $(H_1)_v(k_s)$, so it follows from Theorem 3.9 that $\Sigma_{v,k_s}(H_1)$ contains a Γ -fixed and $(H_1)_v(k_s)$ -fixed point x_1 . We can write $x_1 = \zeta(\mu_1)$ for some

$\mu_1 \in Y_{k_s}(H_1)$. Then μ_1 destabilizes v but does not fix v ; moreover, $P_{\mu_1}(H_1^0)$ is Γ -stable and is normalized by $(H_1)_v(k_s)$. In particular, $P_{\mu_1}(H_1^0)$ is k -defined.

By Lemma 4.1, there exist $n \in \mathbb{N}$ and $\mu \in Y_{k_s}(M)$ such that $\pi \circ \mu = n\mu_1$ and μ destabilizes v ; note that μ does not fix v , because μ_1 does not. The map π gives a bijection between the parabolic subgroups of M^0 and the parabolic subgroups of H_1^0 . So $P_\mu(M^0)$ is Γ -stable—because $P_{\mu_1}(H_1^0)$ is—and hence is defined over k . As $\pi(H_v(k_s))$ is contained in $(H_1)_v(k_s)$ and $(H_1)_v(k_s)$ normalizes $P_{\mu_1}(H_1^0)$, $H_v(k_s)$ normalizes $P_\mu(M^0)$.

After replacing μ if necessary with an $R_u(P_\mu(H^0))(k_s)$ -conjugate of μ , we can assume that μ is a cocharacter of a k -defined maximal torus T of $P_\mu(H^0)$. Let $\mu^{(1)}, \dots, \mu^{(r)}$ be the Γ -conjugates of μ . These are cocharacters of T , so they all commute with each other. Set $\sigma = \sum_{i=1}^r \mu^{(i)}$, a k -defined cocharacter of T . Note that σ centralizes $S = S_0S_2$. Now $\pi \circ \sigma$ destabilizes v but does not fix v (since $\pi \circ \sigma \neq 0$), so σ does not fix v . This implies by [1, Lem. 2.8] that σ properly destabilizes v over k_s for G . Since $H_v(k_s)$ is Γ -stable and fixes $\zeta(\mu)$, $H_v(k_s)$ fixes $\zeta(\mu^{(i)})$ for all $1 \leq i \leq r$. It follows from Lemma 4.4 that $H_v(k_s)$ fixes $\zeta(\sigma)$: that is, for all $h \in H_v(k_s)$, there exists $u \in R_u(P_\sigma(H^0))(k_s)$ such that $h \cdot \sigma = u \cdot \sigma$. Hence $P_\sigma(G^0)$ is normalized by $H_v(k_s) = G_v(k_s)$. This completes the proof of (i).

Parts (ii) and (iii) now follow as in the proof of Theorem 1.1 (there is no need to reduce to the case when v is a k -point in (iii) because we already assume this). \square

Remark 4.6. It can be shown that Theorem 1.3(iii) actually holds without the assumption that v is a k -point. Here is a sketch of the proof. It is enough to prove that Levi ascent holds. Without loss, assume S is a maximal k -defined torus of G_v . As in the proof of Theorem 1.1(iii), we replace v with a k -point w of a k -defined G -variety W , with the property that $\Sigma_{w,k_s} \subseteq \Sigma_{v,k_s}$ and $\Sigma_{w,k_s} = \Sigma_{v,k_s}$. By the arguments of [1, Sec. 6], we can assume that S and N fix w . Suppose $G(k) \cdot v$ is not cocharacter-closed over k . Then $G(k) \cdot w$ is not cocharacter-closed over k , so $L(k_s) \cdot w$ is not cocharacter-closed over k_s , by Galois descent and split Levi ascent. It follows that $H_1(k_s) \cdot w$ is not cocharacter-closed over k_s . We do not know whether hypotheses (a) and (b) of Theorem 1.3 hold for w . The key point, however, that makes the proof of Theorem 1.3(i) work is that $\Sigma_{v,k_s}(H_1)$ is contained in a single apartment of $\Delta_{k_s}(H_1, \mathbb{Q})$. The analogous property holds for $\Sigma_{w,k_s}(H_1)$ since $\Sigma_{w,k_s}(H_1) \subseteq \Sigma_{v,k_s}(H_1)$. Hence Galois ascent holds and $H_1(k) \cdot w$ is not cocharacter-closed over k . Then $H_1(k) \cdot v$ is not cocharacter-closed over k , and the result follows.

5. APPLICATIONS TO G -COMPLETE REDUCIBILITY

Many of the constructions in this paper, and in the key references [6], [4], [1], were inspired originally by the study of Serre's notion of G -complete reducibility for subgroups of G . We refer the reader to [19] and [2] for a thorough introduction to the theory. We simply record the basic definition here:

Definition 5.1. A subgroup H of G is said to be G -completely reducible over k if whenever H is contained in a k -defined R-parabolic subgroup P of G , there exists a k -defined R-Levi subgroup L of P containing H . (We do not assume that H is k -defined.)

The following is [1, Thm. 9.3].

Theorem 5.2. *Let H be a subgroup of G and let $\mathbf{h} \in H^n$ be a generic tuple of H (see [6, Def. 5.4]). Then H is G -completely reducible over k if and only if $G(k) \cdot \mathbf{h}$ is cocharacter-closed over k , where G acts on G^n by simultaneous conjugation.*

Theorem 5.2 allows us to prove results about G -complete reducibility over k using our results on geometric invariant theory. If G is connected, and $\mathbf{h} \in G^n$ is a generic tuple for a subgroup H of G , then $\Sigma_{\mathbf{h}}$ is a subcomplex of $\Delta_G(\mathbb{Q})$, since for any $\lambda \in Y$, λ destabilizes \mathbf{h} if and only if $H \subseteq P_\lambda$; this means that we are in the territory of Theorem 1.1.

Proof of Theorem 1.4. Let \mathbf{h} be a generic tuple of H . Then $\Sigma_{\mathbf{h},k_s}$ is a subcomplex of Δ_{G,k_s} , and $C_G(H) = G_{\mathbf{h}}$. The result now follows from Theorems 5.2 and 1.1(iii). \square

This theory has a counterpart for Lie algebras of $\mathfrak{g} := \text{Lie}(G)$. The basic definitions and results were covered for algebraically closed fields in [13] and [5, Sec. 3.3], but the extension to arbitrary fields is straightforward (cf. [5, Rem. 4.16]).

Definition 5.3. A Lie subalgebra \mathfrak{h} of G is G -completely reducible over k if whenever P is a k -defined parabolic subgroup of G such that $\mathfrak{h} \subseteq \text{Lie}(P)$, there exists a k -defined Levi subgroup L of P such that $\mathfrak{h} \subseteq \text{Lie}(L)$. (We do not assume that \mathfrak{h} is k -defined.)

The group G acts on \mathfrak{g}^n via the simultaneous adjoint action for any $n \in \mathbb{N}$. The next result follows from [5, Lem. 3.8] and the arguments in the proofs of [5, Thm. 4.12(iii)] (cf. [5, Thm. 3.10(iii)]).

Theorem 5.4. *Let \mathfrak{h} be a Lie subalgebra of \mathfrak{g} and let $\mathbf{h} \in \mathfrak{h}^n$ such that the components of \mathbf{h} generate \mathfrak{h} as a Lie algebra. Then \mathfrak{h} is G -completely reducible over k if and only if $G(k) \cdot \mathbf{h}$ is cocharacter-closed over k .*

We now give the applications of our earlier results to G -complete reducibility over k for Lie algebras.

Theorem 5.5. *Let \mathfrak{h} be a Lie subalgebra of \mathfrak{g} .*

- (i) *Suppose \mathfrak{h} is k -defined, and let k'/k be a separable algebraic extension. Then \mathfrak{h} is G -completely reducible over k' if and only if \mathfrak{h} is G -completely reducible over k .*
- (ii) *Let S be a k -defined torus of $C_G(\mathfrak{h})$ and set $L = C_G(S)$. Then \mathfrak{h} is G -completely reducible over k if and only if \mathfrak{h} is L -completely reducible over k .*

Proof. Pick $\mathbf{h} \in \mathfrak{h}^n$ for some $n \in \mathbb{N}$ such that the components of \mathbf{h} generate \mathfrak{h} as a Lie algebra. If \mathfrak{h} is k -defined then we can assume that $\mathbf{h} \in \mathfrak{h}(k)^n$. Part (i) now follows from Theorems 5.4 and 1.1(ii), and part (ii) from Theorems 5.4 and 1.1(iii). \square

Acknowledgments: The authors acknowledge the financial support of EPSRC Grant EP/L005328/1 and of Marsden Grants UOC1009 and UOA1021. Part of the research for this paper was carried out while the authors were staying at the Mathematical Research Institute Oberwolfach supported by the ‘‘Research in Pairs’’ programme. Also, parts of this paper were written during mutual visits to Auckland, Bochum and York.

REFERENCES

- [1] M. Bate, S. Herpel, B. Martin, G. Röhrle, *Cocharacter closure and the rational Hilbert-Mumford Theorem*, <http://arxiv.org/abs/1411.7849>
- [2] M. Bate, B. Martin, G. Röhrle, *A geometric approach to complete reducibility*, *Invent. Math.* **161**, no. 1 (2005), 177–218.
- [3] ———, *On Tits’ Centre Conjecture for fixed point subcomplexes*, *C. R. Acad. Sci. Paris Ser. I Math.* **347**, (2009) 353–356.

- [4] ———, *The strong Centre Conjecture: an invariant theory approach*, J. Algebra **372** (2012), 505–530.
- [5] M. Bate, B. Martin, G. Röhrle, R. Tange, *Complete reducibility and conjugacy classes of tuples in algebraic groups and Lie algebras*, Math. Z. **269** (2011), 809–832.
- [6] ———, *Closed orbits and uniform S -instability in geometric invariant theory*, Trans. Amer. Math. Soc., **365** (2013), no. 7, 3643–3673.
- [7] A. Borel, *Linear algebraic groups*, Graduate Texts in Mathematics, **126**, Springer-Verlag 1991.
- [8] B. Conrad, O. Gabber, and G. Prasad, *Pseudo-reductive groups*, New Mathematical Monographs, vol. 17, Cambridge University Press, Cambridge, 2010.
- [9] W.H. Hesselink, *Uniform instability in reductive groups*, J. Reine Angew. Math. **303/304** (1978), 74–96.
- [10] J.E. Humphreys, *Linear Algebraic Groups*, Springer-Verlag, New York, 1975.
- [11] G.R. Kempf, *Instability in invariant theory*, Ann. Math. **108** (1978), 299–316.
- [12] B. Leeb, C. Ramos-Cuevas, *The center conjecture for spherical buildings of types F_4 and E_6* , Geom. Funct. Anal. **21** (2011), no. 3, 525–559.
- [13] G. McNinch, *Completely reducible Lie subalgebras*, Transform. Groups **12** (2007), no. 1, 127–135.
- [14] B. Mühlherr, J. Tits, *The center conjecture for non-exceptional buildings*, J. Algebra **300** (2006), no. 2, 687–706.
- [15] B. Mühlherr, R.M. Weiss, *Receding polar regions of a spherical building and the center conjecture*, Ann. Inst. Fourier (Grenoble) **63** (2013), no. 2, 479–513.
- [16] C. Ramos-Cuevas, *The center conjecture for thick spherical buildings*, Geom. Dedicata **166** (2013), 349–407.
- [17] G. Rousseau, *Immeubles sphériques et théorie des invariants*, C.R.A.S. **286** (1978), 247–250.
- [18] J.-P. Serre, *La notion de complète réductibilité dans les immeubles sphériques et les groupes réductifs*, Séminaire au Collège de France, résumé (1997).
- [19] ———, *Complète réductibilité*, Séminaire Bourbaki, 56ème année, 2003–2004, n° 932.
- [20] T.A. Springer, *Linear algebraic groups*, Second edition. Progress in Mathematics, 9. Birkhäuser Boston, Inc., Boston, MA, 1998.
- [21] W.C. Waterhouse, *Introduction to affine group schemes*, Graduate Texts in Mathematics, vol. 66, Springer-Verlag, New York, 1979.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF YORK, YORK YO10 5DD, UNITED KINGDOM
E-mail address: michael.bate@york.ac.uk

FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, D-44780 BOCHUM, GERMANY
E-mail address: sebastian.herpel@rub.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ABERDEEN, KING’S COLLEGE, FRASER NOBLE BUILDING, ABERDEEN AB24 3UE, UNITED KINGDOM
E-mail address: b.martin@abdn.ac.uk

FAKULTÄT FÜR MATHEMATIK, RUHR-UNIVERSITÄT BOCHUM, D-44780 BOCHUM, GERMANY
E-mail address: gerhard.roehrle@rub.de