



Mathematisches
Forschungsinstitut
Oberwolfach

Member of the



Oberwolfach Preprints

OWP 2022 - 09

DIMITAR GRANTCHAROV, IVAN PENKOV AND
VERA SERGANOVA

Bounded Weight Modules for Basic Classical
Lie Superalgebras at Infinity

Mathematisches Forschungsinstitut Oberwolfach gGmbH
Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

The MFO publishes a preprint series **Oberwolfach Preprints (OWP)**, ISSN 1864-7596, which mainly contains research results related to a longer stay in Oberwolfach, as a documentation of the research work done at the MFO. In particular, this concerns the Oberwolfach Research Fellows program (and the former Research in Pairs program) and the Oberwolfach Leibniz Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it in electronic and printed form. Every OWRP group or Oberwolfach Leibniz Fellow may receive on request 20 free hard copies (DIN A4, black and white copy) by surface mail.

The full copyright is left to the authors. With the submission of a manuscript, the authors warrant that they are the creators of the work, including all graphics. The authors grant the MFO a perpetual, irrevocable, non-exclusive right to publish it on the MFO's institutional repository. Since the right is non-exclusive, the MFO enables parallel or later publications, e.g. on the researcher's personal website, in arXiv or in a journal. Whether the other journals also accept preprints or postprints can be checked, for example, via the Sherpa Romeo service.

In case of interest, please send a **pdf file** of your preprint by email to owrf@mfo.de. The file should be sent to the MFO within 12 months after your stay at the MFO.

The preprint (and a published paper) should contain an acknowledgement like: *This research was supported through the program "Oberwolfach Research Fellows" (resp. "Oberwolfach Leibniz Fellows") by the Mathematisches Forschungsinstitut Oberwolfach in [year].*

There are no requirements for the format of the preprint, except that the paper size (or format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX). Additionally, each preprint will get a Digital Object Identifier (DOI).

We cordially invite the researchers within the OWRP and OWLF program to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany

Tel +49 7834 979 50
Fax +49 7834 979 55
Email admin@mfo.de
URL www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.
Copyright of the content is held by the authors.

DOI 10.14760/OWP-2022-09

BOUNDED WEIGHT MODULES FOR BASIC CLASSICAL LIE SUPERALGEBRAS AT INFINITY

DIMITAR GRANTCHAROV, IVAN PENKOV AND VERA SERGANOVA

ABSTRACT. We classify simple bounded weight modules over the complex simple Lie superalgebras $\mathfrak{sl}(\infty|\infty)$ and $\mathfrak{osp}(m|2n)$, when at least one of m and n equals ∞ . For $\mathfrak{osp}(m|2n)$ such modules are of spinor-oscillator type, i.e., they combine into one the known classes of spinor $\mathfrak{o}(m)$ -modules and oscillator-type $\mathfrak{sp}(2n)$ -modules. In addition, we characterize the category of bounded weight modules over $\mathfrak{osp}(m|2n)$ (under the assumption $\dim \mathfrak{osp}(m|2n) = \infty$) by reducing its study to already known categories of representations of $\mathfrak{sp}(2n)$, where n possibly equals ∞ . When classifying simple bounded weight $\mathfrak{sl}(\infty|\infty)$ -modules, we prove that every such module is integrable over one of the two infinite-dimensional ideals of the Lie algebra $\mathfrak{sl}(\infty|\infty)_{\bar{0}}$. We finish the paper by establishing some first facts about the category of bounded weight $\mathfrak{sl}(\infty|\infty)$ -modules.

2020 MSC: Primary 17B65, 17B10

Keywords and phrases: direct limit Lie superalgebra, Clifford superalgebra, Weyl superalgebra, weight module, annihilator.

INTRODUCTION

The representation theory of the three simple infinite-dimensional finitary complex Lie algebras $\mathfrak{sl}(\infty)$, $\mathfrak{o}(\infty)$, and $\mathfrak{sp}(\infty)$ has made notable progress in the last three decades, see for instance [DPS], [DP], [PSer1], [PSer2], [PStyr], [SS]. For a summary of highlights of this theory see [PH]. The theory of representations of the super-counterparts of the Lie algebras $\mathfrak{sl}(\infty)$, $\mathfrak{o}(\infty)$, and $\mathfrak{sp}(\infty)$ is still much less developed. For a finite-dimensional Lie superalgebra \mathfrak{k} , the category of all representations of \mathfrak{k} is almost never equivalent to the category of all representations of the Lie algebra $\mathfrak{k}_{\bar{0}}$, the even part of \mathfrak{k} . However, in that case there is a general result claiming that a category of representations of \mathfrak{k} with fixed strongly typical central character is equivalent to a corresponding category of representations of $\mathfrak{k}_{\bar{0}}$.

This result does not provide a clear guideline for the case of Lie superalgebras of infinite rank since the center of the enveloping algebra of Lie superalgebras like $\mathfrak{sl}(\infty|\infty)$ or $\mathfrak{osp}(\infty|\infty)$ is trivial. Nevertheless, in the study of reasonably small categories of representations over the Lie superalgebras $\mathfrak{sl}(\infty|\infty)$ and $\mathfrak{osp}(\infty|\infty)$, one may rely on different intuition and obtain results not necessarily following the above pattern. For instance, in [S] it is shown that the category of tensor modules over the

Lie superalgebra $\mathfrak{osp}(\infty|\infty)$ (respectively, over $\mathfrak{sl}(\infty|\infty)$) is equivalent to the categories of tensor modules over each of the Lie algebras $\mathfrak{o}(\infty)$ and $\mathfrak{sp}(\infty)$ (respectively, over $\mathfrak{sl}(\infty)$). A somewhat similar phenomenon can be seen in the paper [CP], where it is proved that the categories of integrable bounded weight modules over various Lie superalgebras like $\mathfrak{sl}(\infty|\infty)$ or $\mathfrak{osp}(\infty|\infty)$ are semisimple.

In the present paper we study the categories of arbitrary (i.e., not necessarily integrable) bounded weight modules over the complex Lie superalgebras $\mathfrak{osp}(m|2n)$, where at least one of m or n equals ∞ , and over the Lie superalgebra $\mathfrak{sl}(\infty|\infty)$. Before describing our results we should recall that for the infinite-dimensional Lie algebras $\mathfrak{sl}(\infty)$, $\mathfrak{o}(\infty)$, $\mathfrak{sp}(\infty)$ simple bounded weight modules have been classified in [GP] and their structure has been further studied in [C].

Our first main result claims that any simple bounded weight module over an infinite-dimensional Lie superalgebra $\mathfrak{osp}(m|2n)$ has just length two (or one for a trivial module) over the Lie algebra $\mathfrak{osp}(m|2n)_{\bar{0}} = \mathfrak{o}(m) \oplus \mathfrak{sp}(n)$. Moreover, such a module (unless it is a natural or trivial module) is determined by a pair (S, N) , where S is a spinor $\mathfrak{o}(m)$ -module and N is a $\mathfrak{sp}(2n)$ -module of oscillator type, i.e., a close relative of the oscillator representations of $\mathfrak{sp}(2n)$. (The notions of spinor $\mathfrak{o}(m)$ -modules and oscillator-type $\mathfrak{sp}(2n)$ -modules make sense also for $m = \infty$ and $n = \infty$ due to the results of [GP].) This spectacular fact allows us to identify simple bounded weight $\mathfrak{osp}(m|2n)$ -modules, other than trivial and natural modules, as modules of “spinor-oscillator type”. The latter class of modules of $\mathfrak{osp}(m|2n)$ glues spinor and oscillator-type modules together, and is the ultimate super-symmetric version of both spinor $\mathfrak{o}(m)$ -modules and oscillator-type $\mathfrak{sp}(2n)$ -modules.

The classification of simple bounded weight $\mathfrak{sl}(\infty|\infty)$ -modules is also very interesting and constitutes our second main result. In particular, we show that every such module is integrable and semisimple with respect to a simple ideal of $\mathfrak{sl}(\infty|\infty)_{\bar{0}} \simeq (\mathfrak{sl}(\infty) \oplus \mathfrak{sl}(\infty)) \in \mathbb{C}$, and this nicely resembles the answer for the case of $\mathfrak{osp}(\infty|\infty)$ where a bounded weight $\mathfrak{osp}(\infty|\infty)$ -module is necessarily integrable and semisimple as an $\mathfrak{o}(\infty)$ -module.

Our main method of classification is reduction to weight modules of Weyl and Clifford superalgebras of infinitely many variables. We denote these superalgebras respectively by $D(\infty|\infty)$ and $Cl(\infty|\infty)$. There are natural homomorphisms $U(\mathfrak{osp}(\infty|\infty)) \rightarrow Cl(\infty|\infty)$ and $U(\mathfrak{sl}(\infty|\infty)) \rightarrow D(\infty|\infty)$, see Section 2.2. One of our central ideas is that, with the exception of Schur powers of the natural and conatural representations (for $\mathfrak{osp}(\infty|\infty)$ this exception applies only to the trivial and natural representations), all simple bounded weight $\mathfrak{osp}(\infty|\infty)$ - or $\mathfrak{sl}(\infty|\infty)$ -modules are annihilated by the kernel of the respective homomorphism. This facilitates a reduction of the study of simple bounded weight $\mathfrak{osp}(\infty|\infty)$ - and $\mathfrak{sl}(\infty|\infty)$ -modules, as well as of the respective categories of bounded weight modules, to the study of weight modules of the associative superalgebras $Cl(\infty|\infty)$ and $D(\infty|\infty)$ and their relevant subalgebras. The above method applies also to the case of $\mathfrak{osp}(m|2n)$ where m or n is finite, and to $\mathfrak{sl}(\infty|n)$ for $n \in \mathbb{Z}_{>0}$ as well.

Here is a brief description of the content of the paper. Section 1 is devoted to preliminaries. In Section 2 we undertake a study of the categories of weight modules over Clifford and Weyl superalgebras. In particular, we establish that any such simple module is multiplicity free. In Sections 3 and 4 we apply the above results to the case of $\mathfrak{osp}(m|2n)$ where at least one of m and n equals infinity. We show that any simple non-integrable bounded weight $\mathfrak{osp}(m|2n)$ -module is a spinor-oscillator module. Moreover, we prove that the category of spinor-oscillator representations is equivalent to the category of multiplicity free non-integrable weight modules over the Lie algebra $\mathfrak{osp}(m|2n)_{\bar{0}} = \mathfrak{o}(m) \oplus \mathfrak{sp}(2n)$.

The case of $\mathfrak{sl}(\infty|\infty)$ is discussed in Section 5. Here we give a classification of the simple bounded weight $\mathfrak{sl}(\infty|\infty)$ -representations and make a first step towards understanding the category of such representations. A deeper study of this category should be a separate project.

Acknowledgment. We thank Lucas Calixto for reading a preliminary draft of the paper. IP and VS acknowledge the outstanding hospitality of Mathematisches Forschungsinstitut Oberwolfach where this paper was almost completed. DG was supported in part by Simons Collaboration Grant 855678. IP was supported in part by DFG grant PE 980/8-1. VS was supported in part by NSF grant 2001191.

1. PRELIMINARIES

The base field is \mathbb{C} . By S_n we denote the symmetric group on n letters. A *superspace* is a \mathbb{Z}_2 -graded vector space where $\mathbb{Z}_2 := \mathbb{Z}/2\mathbb{Z}$, and a *superalgebra* is a \mathbb{Z}_2 -graded algebra. We use the indices $\bar{0}$ and $\bar{1}$ to indicate \mathbb{Z}_2 -gradings. A *purely even* (respectively, *purely odd*) superspace is a superspace V such that $V = V_{\bar{0}}$ (resp., $V = V_{\bar{1}}$). By Π we denote the parity change functor on superspaces: $(\Pi V)_{\bar{0}} = V_{\bar{1}}$, $(\Pi V)_{\bar{1}} = V_{\bar{0}}$. If $V = V_{\bar{0}} \oplus V_{\bar{1}}$ is a superspace, then the *dual* superspace equals $V_{\bar{0}}^* \oplus V_{\bar{1}}^*$, where $V_{\bar{0}}^* = \text{Hom}(V_{\bar{0}}, \mathbb{C})$, $V_{\bar{1}}^* = \Pi \text{Hom}(\Pi V_{\bar{1}}, \mathbb{C})$ and Hom stands here for homomorphisms of purely even spaces.

We write $S^k V$ and $\Lambda^k V$ for the k th symmetric and exterior powers for a superspace V . If W is a superspace of parity $p \in \mathbb{Z}_2$ (i.e., $W = W_{\bar{0}}$ for $p = \bar{0}$ and $W = W_{\bar{1}}$ for $p = \bar{1}$), then $S^k W$ (respectively, $\Lambda^k W$) is a superspace of parity $kp \in \mathbb{Z}_2$ (respectively, $kp + \bar{1} \in \mathbb{Z}_2$). For a general superspace $V = V_{\bar{0}} \oplus V_{\bar{1}}$ we have

$$S^k V = \bigoplus_{i+j=k} S^i V_{\bar{0}} \otimes \Lambda^j V_{\bar{1}}, \quad \Lambda^k V = \bigoplus_{i+j=k} \Lambda^i V_{\bar{0}} \otimes S^j V_{\bar{1}}.$$

An *even symmetric* (respectively, *even antisymmetric*) bilinear form on a superspace V is a parity-preserving linear operator $S^2 V \rightarrow \mathbb{C}$ (respectively, $\Lambda^2 V \rightarrow \mathbb{C}$).

In this paper we work with the Lie superalgebras $\mathfrak{gl}(a|b)$, $\mathfrak{sl}(a|b)$, $\mathfrak{osp}(2a|2b)$, $\mathfrak{osp}(2a+1|2b)$, where $a, b \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$. Their *defining representation* is the simple module of respective dimension $(a|b)$, $(a|b)$, $(2a|2b)$, $(2a+1|2b)$. In what follows we

use the term defining representation more loosely to include also the defining representation with changed parity. The Lie superalgebras $\mathfrak{gl}(a|b)$, $\mathfrak{sl}(a|b)$, $\mathfrak{osp}(2a|2b)$, $\mathfrak{osp}(2a+1|2b)$ can be equipped with a fixed even symmetric invariant form (\cdot, \cdot) . All homomorphisms of superalgebras are assumed to preserve the \mathbb{Z}_2 -grading. All modules over purely even associative algebras or Lie algebras are assumed to be purely even unless otherwise stated.

We assume that Cartan subalgebras of the Lie superalgebras considered are fixed, and use standard notation for the roots. Note that these Cartan subalgebras are purely even and all root spaces are either purely even or purely odd. Therefore the roots are designated as even or odd. Concretely, the even roots of $\mathfrak{gl}(a|b)$ and $\mathfrak{sl}(a|b)$ are $\varepsilon_i - \varepsilon_k, \delta_j - \delta_l$, while the odd roots are $\pm(\varepsilon_i - \delta_j)$, where $1 \leq i \neq k \leq a, 1 \leq j \neq l \leq b$. The even roots of $\mathfrak{osp}(2a|2b)$ are $\pm(\varepsilon_i \pm \varepsilon_k), \pm(\delta_j \pm \delta_l), \pm 2\delta_j$, and the odd roots are $\pm(\varepsilon_i - \delta_j)$. For $\mathfrak{osp}(2a+1|2b)$ we have in addition the even roots $\pm\varepsilon_i$ and the odd roots $\pm\delta_j$.

We should point out that for $a = \infty$ the Lie superalgebras $\mathfrak{osp}(2a+1|2b)$ and $\mathfrak{osp}(2a|2b)$ are isomorphic, and the difference in root systems is the result of different choices of Cartan subalgebras. A less brief discussion of the Lie superalgebras we consider and their root systems can be found in [CP].

Let \mathfrak{s} be a Lie algebra or Lie superalgebra with a fixed Cartan subalgebra $\mathfrak{h} = \mathfrak{h}_0$. A *weight* module M is an \mathfrak{s} -module that is semisimple as \mathfrak{h} -module. The \mathfrak{h} -isotypic components of M are the *weight spaces* of M : we denote them by M^λ for $\lambda \in \mathfrak{h}^*$. The weight spaces of M are superspaces. Every weight module M has a well-defined support:

$$\text{supp}M = \{\mu \in \mathfrak{h}^* \mid M^\mu \neq 0\}.$$

A *weight* module is *bounded* if the dimension $(d_0|d_1)$ of any weight space of M is less or equal to $(a|b)$ for some fixed $a, b \in \mathbb{Z}_{\geq 0}$, i.e., $d_0 \leq a, d_1 \leq b$. The *degree* $d(M)$ of a bounded weight module M equals the maximum value of the sum $d_0 + d_1$ over all weight spaces of M .

Each of our Lie superalgebras has (up to isomorphism) two natural modules which we denote by V and IV . These modules are weight modules, and for $\mathfrak{gl}(a|b)$ and $\mathfrak{sl}(a|b)$ we assume that the weight spaces of weight ε_i in V are purely odd and the weight spaces of weight δ_j in V are purely even. For $\mathfrak{osp}(2a+1|2b)$ and $\mathfrak{osp}(2a|2b)$ we make the opposite choice. We have

$$\text{supp}V = \begin{cases} \{\varepsilon_i, \delta_j \mid i, j > 0\} & \text{if } \mathfrak{g} = \mathfrak{sl}(a|b) \text{ or } \mathfrak{g} = \mathfrak{gl}(a|b), \\ \{0, \pm\varepsilon_i, \pm\delta_j \mid i, j > 0\} & \text{if } \mathfrak{g} = \mathfrak{osp}(2a+1|2b), \\ \{\pm\varepsilon_i, \pm\delta_j \mid i, j > 0\} & \text{if } \mathfrak{g} = \mathfrak{osp}(2a|2b). \end{cases}$$

For $\mathfrak{gl}(a|b)$ and $\mathfrak{sl}(a|b)$ modules V_* and IV_* are also well defined. They are characterized by equalities $\text{supp}V_* = -\text{supp}V$, $\text{supp}\text{IV}_* = -\text{supp}\text{IV}$, and by the fact that the weight spaces of weight $-\varepsilon_i$ in V_* are purely odd and the weight spaces of weight $-\varepsilon_i$ in IV_* are purely even.

We now recall some facts about *multiplicity free* weight \mathfrak{s} -modules for a finite-dimensional Lie algebra \mathfrak{s} , i.e., bounded weight \mathfrak{s} -modules M with $d(M) = 1$. Their classification has been part of a major effort to classify simple weight modules with finite-dimensional weight spaces. Some of the main contributors have been D. Britten, F. Lemire, S. Fernando, V. Futorny, G. Benkart, O. Mathieu, and Mathieu's paper [Mat] can be considered as the crown of this effort. It follows from a result of Fernando [Fer] that for $\mathfrak{s} = \mathfrak{o}(n)$, $n \geq 5$ every multiplicity free simple weight $\mathfrak{o}(n)$ is finite dimensional, hence is a trivial module, natural module, or a spinor module. For $\mathfrak{s} = \mathfrak{sp}(2n)$ the only multiplicity free simple finite-dimensional \mathfrak{s} -modules are the trivial and the natural modules, and there is a "coherent family" of infinite-dimensional multiplicity free simple weight \mathfrak{s} -modules [BBL], [Mat]. For every Borel subalgebra $\mathfrak{b} \supset \mathfrak{h}$, there are precisely two nonisomorphic multiplicity free simple \mathfrak{b} -highest weight modules in this family. These highest weight modules are known as oscillator or Shale-Weil modules, and every other infinite-dimensional multiplicity free simple weight module is obtained from one of them via twisted localization, see [Mat]. For $\mathfrak{s} = \mathfrak{sl}(n)$ the simple multiplicity free weight modules have been classified in [BBL] and have been further studied by O. Mathieu in [Mat]. In this paper we will not refer to the description of all simple multiplicity free weight modules for $\mathfrak{sl}(n)$ and $\mathfrak{sp}(2n)$, but for understanding our results it is essential to know that simple multiplicity free weight modules, and more generally simple bounded weight modules, are well studied.

For $\mathfrak{s} = \mathfrak{sl}(\infty)$, $\mathfrak{sp}(\infty)$, $\mathfrak{o}(\infty)$, simple bounded weight modules are described explicitly in [GP]. In the case of $\mathfrak{o}(\infty)$, any bounded weight module is *integrable*, i.e., it is a direct limit of finite-dimensional $\mathfrak{o}(n)$ -modules for $n \rightarrow \infty$. More precisely, if M is a simple bounded weight $\mathfrak{o}(\infty)$ -module, then M is a trivial module, a natural module, or a direct limit of spinor modules. We refer to the latter direct limits simply as *spinor $\mathfrak{o}(\infty)$ -modules*. For $\mathfrak{s} = \mathfrak{sp}(\infty)$ the result is similar. Namely, a simple bounded weight $\mathfrak{sp}(\infty)$ -module is a trivial module, a natural module, or a direct limit of simple multiplicity free infinite-dimensional $\mathfrak{sp}(2n)$ -modules for $n \rightarrow \infty$. A difference with the case of $\mathfrak{o}(\infty)$ is that a direct limit of simple multiplicity free infinite-dimensional modules is not integrable. We call such a direct limit a *simple weight $\mathfrak{sp}(\infty)$ -module of oscillator type*.

In the sequel we will need the following general lemma about associative superalgebras.

Lemma 1.1. *Let A be an associative superalgebra and X be a simple A -module. Then $X_{\bar{0}}$ and $X_{\bar{1}}$ are simple $A_{\bar{0}}$ -modules.*

Proof. If $Y \subset X_{\bar{0}}$ (respectively, $Y \subset X_{\bar{1}}$) is a proper nonzero $A_{\bar{0}}$ -submodule then AY is an A -submodule of Y and $(AY)_{\bar{0}} = Y$ (respectively, $(AY)_{\bar{1}} = Y$). \square

We conclude Section 1 with some facts concerning finite-dimensional Lie (super)algebras \mathfrak{s} . For a partition (equivalently, a Young diagram) μ , let \mathbb{S}_{μ} denote the corresponding Schur functor.

Proposition 1.2. *Let $\mathfrak{s} = \mathfrak{sl}(n)$ and V be the defining \mathfrak{s} -module. If $n \geq |\mu|$ then $d(\mathbb{S}_\mu V)$ equals the dimension of the simple $S_{|\mu|}$ -module Z_μ associated to μ .*

Proof. It suffices to consider the case $n = |\mu|$. Let $\{e_1, \dots, e_n\}$ be the standard \mathfrak{h} -eigenbasis of V . Let $\omega = \varepsilon_1 + \dots + \varepsilon_n$. Then the weight space $(V^{\otimes n})^\omega$ has a structure of $W \times S_n$ -module, where $W \simeq S_n$ is the Weyl group of $\mathfrak{sl}(n)$. Moreover, as an S_n -module $(V^{\otimes n})^\omega$ is isomorphic to the regular representation of S_n . Therefore, the isomorphism

$$(V^{\otimes n})^\omega \simeq \bigoplus_{\mu} (\mathbb{S}_\mu V)^\omega \otimes Z_\mu$$

forces $\dim(\mathbb{S}_\mu V)^\omega = \dim Z_\mu$. \square

Lemma 1.3. *Let \mathfrak{s} be a simple finite-dimensional Lie algebra, and $L(\mu), L(\nu)$ be simple finite-dimensional modules with respective highest weights μ, ν . Then $d(L(\mu + \nu)) \geq d(L(\mu))$.*

Proof. Let $\pi : L(\mu) \otimes L(\nu) \rightarrow L(\mu + \nu)$ be the unique surjective homomorphism. Then the restriction of π to $L(\mu)^\lambda \otimes L(\nu)^\nu$ is injective, where λ is a weight of $L(\mu)$ of maximal multiplicity. This implies the statement. \square

Lemma 1.4. *Let $\mathfrak{s} = \mathfrak{o}(2n + 1), \mathfrak{o}(2n)$, or $\mathfrak{sp}(2n)$. Then $L(\mu)$ is either multiplicity free or $d(L(\mu)) \geq n - 1$.*

Proof. Let ω_i be the i th fundamental weight of \mathfrak{s} . Set $\mathfrak{s} = \mathfrak{o}(2n + 1)$. Then $d(L(\omega_1)) = d(L(\omega_n)) = 1$. For $k = 2, \dots, n - 1$ we have $L(\omega_k) \simeq \Lambda^k V$, thus $d(L(\omega_k)) = \binom{n}{\lfloor k/2 \rfloor} \geq n - 1$. Next we note that

$$\begin{aligned} d(L(2\omega_1)) &= d(S^2 V) = n, \\ d(L(2\omega_n)) &= d(\Lambda^n V) = \binom{n}{\lfloor n/2 \rfloor} \geq n - 1, \end{aligned}$$

and

$$d(L(\omega_1 + \omega_n)) \geq d(L(\omega_1) \otimes L(\omega_n)) - d(L(\omega_n)) = n.$$

Consequently, for $\mu = \omega_1, \dots, \omega_n, 2\omega_1, 2\omega_n, \omega_1 + \omega_n$ we see that $d(L(\mu)) \geq n - 1$. For any other μ the inequality follows from Lemma 1.3.

The case of $\mathfrak{o}(2n)$ is similar.

Now let $\mathfrak{s} = \mathfrak{sp}(2n)$. Then $d(L(\omega_1)) = d(V) = 1$. For $k > 1$ we have $L(\omega_k) = \Lambda^k V / \Lambda^{k-2} V$. Hence $d(L(\omega_k)) = \binom{n}{\lfloor k/2 \rfloor} - \binom{n}{\lfloor k/2 \rfloor - 1} \geq n - 1$. Next, $L(2\omega_1)$ is the adjoint representation and hence $d(L(2\omega_1)) = n$. For $\mu \neq \omega_1, \dots, \omega_n, 2\omega_1$, the statement follows again from Lemma 1.3. \square

Lemma 1.5. *Let $\mathfrak{s} = \mathfrak{osp}(1|2n)$ and let $L(\mu)$ be a finite-dimensional simple \mathfrak{s} -module with highest weight μ relative to the Borel subsuperalgebra with simple roots $\delta_1 - \delta_2, \dots, \delta_{n-1} - \delta_n, \delta_n$. Assume $d(L(\mu)) < n$. Then $\mu = \delta_1, \mu = 0$, or $\mu = -\frac{1}{2}(\delta_1 + \dots + \delta_n)$.*

Proof. The restriction of $L(\lambda)$ to $\mathfrak{s}_0 = \mathfrak{sp}(2n)$ can have only simple constituents with highest weights 0 , δ_1 , or $-\frac{1}{2}(\delta_1 + \cdots + \delta_n)$, $-\frac{1}{2}(\delta_1 + \cdots + \delta_{n-1}) - \frac{3}{2}\delta_n$. The statement follows. \square

In this paper a *bounded* primitive ideal of $U(\mathfrak{s})$ is defined as a primitive ideal which annihilates a simple bounded weight \mathfrak{s} -module. It is a result of [PSer3] that if M and N are simple weight modules annihilated by the same bounded primitive ideal I , then M and N are bounded and $d(M) = d(N)$. This allows to define the *degree* of a bounded primitive ideal $I \subset U(\mathfrak{s})$ by setting $d(I) := d(M)$ for any simple bounded weight \mathfrak{s} -module M annihilated by I .

Lemma 1.6. *Let $\mathfrak{s} = \mathfrak{sp}(2n)$, $\mathfrak{sl}(n)$ and I be a bounded primitive ideal of $U(\mathfrak{s})$ of degree d . Assume that $U(\mathfrak{s})/I$ is infinite dimensional. Then either $d \geq \text{rk } \mathfrak{s} - 1$ or $d = 1$. If $d = 1$ and $\mathfrak{s} = \mathfrak{sl}(n)$, then $I = \text{Ann}_{U(\mathfrak{s})} L(a\omega_1)$ or $I = \text{Ann}_{U(\mathfrak{s})} L(a\omega_n)$ for some $a \notin \mathbb{Z}_{\geq 0}$. If $d = 1$ and $\mathfrak{s} = \mathfrak{sp}(2n)$, then I is the Joseph ideal (annihilator of an oscillator module).*

Proof. Assume first $d > 1$. The inequality $d \geq \text{rk } \mathfrak{s} - 1$ for $\mathfrak{s} = \mathfrak{sl}(n)$ follows from Lemma 2.25 in [GP].

We proceed to show that $d \geq \text{rk } \mathfrak{s} = n$ for $\mathfrak{s} = \mathfrak{sp}(2n)$. Theorem 12.2 in [Mat] implies $d = \frac{1}{2^{n-1}} \dim L_{\mathfrak{o}}(\lambda)$ for some simple finite-dimensional $\mathfrak{o}(2n)$ -module $L_{\mathfrak{o}}(\lambda)$ of highest weight $\lambda = \sum_{i=1}^n \lambda_i \varepsilon_i$ with $\lambda_i \in 1/2 + \mathbb{Z}$. Since $d > 1$, we have $\lambda \neq \omega_{n-1}, \omega_n$. Moreover, if $|\lambda_k| \neq |\lambda_{k+1}|$ for some $k \geq 1$, the stabilizer of λ in the Weyl group has at most $k!(n-k)!$ elements. Therefore the orbit of λ has at least $\binom{n}{k} 2^{n-1}$ elements, implying $d \geq n$. Consider now the case when all absolute values $|\lambda_i|$ are equal. Under this assumption, there are two possibilities: (i) all λ_i are equal, or (ii) $\lambda_1 = \cdots = \lambda_{n-1} = -\lambda_n$. We set $\mu = \lambda - (\varepsilon_{n-1} + \varepsilon_n)$ in case (i) and $\mu = \lambda - (\varepsilon_{n-1} - \varepsilon_n)$ in case (ii). Then μ is a weight of $L_{\mathfrak{o}}(\lambda)$ and the Weyl group orbit of μ has at least $n2^{n-1}$ elements. This implies again $d \geq n$. \square

2. CLIFFORD AND WEYL SUPERALGEBRAS AND WEIGHT MODULES OVER THEM

2.1. Definitions and main properties. Let $a, b \in \mathbb{Z}_{\geq 0} \cup \{\infty\}$. The *Weyl superalgebra* $D(a|b)$ is the associative superalgebra with generators $\{x_i, \partial_i \mid i = 1, \dots, a; -1, \dots, -b\}$ of parity

$$\bar{x}_i = \bar{\partial}_i = \begin{cases} 0 & \text{if } i > 0 \\ 1 & \text{if } i < 0 \end{cases},$$

satisfying the relations

$$[x_i, x_j] = [\partial_i, \partial_j] = 0, \quad [\partial_i, x_j] = \delta_{ij},$$

where $[u, v] := uv - (-1)^{\bar{u}\bar{v}}vu$ and δ_{ij} is Kronecker's delta. The *Clifford superalgebra* $Cl(a|b)$ is the associative superalgebra with generators $\{\xi_i, \eta_i \mid i = 1, \dots, a; -1, \dots, -b\}$

with generators of parity

$$\bar{\xi}_i = \bar{\eta}_i = \begin{cases} 0 & \text{if } i > 0 \\ 1 & \text{if } i < 0 \end{cases},$$

satisfying the relations

$$\{\xi_i, \xi_j\} = \{\eta_i, \eta_j\} = 0, \quad \{\eta_i, \xi_j\} = \delta_{ij},$$

where $\{u, v\} := uv + (-1)^{\bar{u}\bar{v}}vu$. In what follows, whenever $x_i, \partial_i, \xi_i, \eta_i$ are used we assume that the index i is nonzero.

We define a \mathbb{Z} -grading on $D(a|b)$ (respectively, on $Cl(a|b)$) by setting $\deg x_i := 1$, $\deg \partial_i := -1$ (respectively, $\deg \xi_i := 1$, $\deg \eta_i := -1$). If $A = D(a|b)$ or $A = Cl(a|b)$ we denote by A_{ev} the subsuperalgebra of elements of even degree, and by A_n the subsuperspace of elements of degree n . Note that $A_{\bar{0}}$, A_0 , and A_{ev} are three different subsuperalgebras of A .

For $a, b \in \mathbb{Z}_{\geq 0}$, $D(a|b)$ (respectively, $Cl(a|b)$) is naturally embedded in $D(a+1|b)$ and $D(a|b+1)$ (respectively, in $Cl(a+1|b)$ and $Cl(a|b+1)$), and

$$D(\infty|\infty) = \varinjlim D(a|b), \quad Cl(\infty|\infty) = \varinjlim Cl(a|b).$$

2.2. Connection to classical Lie superalgebras. Let $V_{2a|2b}$ be the subsuperspace of $D(a|b)$ with basis $\{x_i, \partial_i \mid -b \leq i \leq a\}$. Then $V_{2a|2b}$ has an even anti-symmetric form given by the commutator map $[V_{2a|2b}, V_{2a|2b}] \rightarrow \mathbb{C}$. The Lie superalgebra $\mathfrak{osp}(2b|2a)$ for which this form is invariant can be identified canonically with $S^2V_{2a|2b}$. The symmetrization map

$$V_{2a|2b}^{\otimes 2} \rightarrow D(a|b), \quad v \otimes w \mapsto \frac{1}{2}(v \otimes w + (-1)^{\bar{v}\bar{w}}w \otimes v),$$

factors through $S^2V_{2a|2b}$ and defines a homomorphism of Lie superalgebras $\mathfrak{osp}(2b|2a) \rightarrow D(a|b)$. This induces a homomorphism of associative superalgebras

$$\Phi_{a|b} : U(\mathfrak{osp}(2b|2a)) \rightarrow D(a|b).$$

Similarly, let $U_{2a|2b}$ be the subsuperspace of $Cl(a|b)$ with basis $\{\xi_i, \eta_i \mid -b \leq i \leq a\}$. Then $U_{2a|2b}$ has an even symmetric bilinear form given by the symmetrizer map $\{U_{2a|2b}, U_{2a|2b}\} \rightarrow \mathbb{C}$. The Lie superalgebra $\mathfrak{osp}(2a|2b)$ for which this form is invariant can be identified canonically with $\Lambda^2U_{2a|2b}$. The alternization map

$$U_{2a|2b}^{\otimes 2} \rightarrow Cl(a|b), \quad v \otimes w \mapsto \frac{1}{2}(v \otimes w - (-1)^{\bar{v}\bar{w}}w \otimes v),$$

factors through $\Lambda^2U_{2a|2b}$ and defines a homomorphism of Lie superalgebras $\mathfrak{osp}(2a|2b) \rightarrow Cl(a|b)$. This induces a homomorphism of associative superalgebras

$$\Psi_{a|b} : U(\mathfrak{osp}(2a|2b)) \rightarrow Cl(a|b).$$

The Chevalley basis vectors e_α and the respective relations of the Lie superalgebras $\mathfrak{osp}(2b|2a)$ (and also of $\mathfrak{osp}(2a|2b)$) have been computed in [FG], §3.2. Up to scalar multiples, the homomorphism $\Phi_{a|b}$ has the form

$$\begin{aligned}
e_{\varepsilon_k - \varepsilon_l} &\mapsto x_{-l} \partial_{-k}, e_{-\varepsilon_k - \varepsilon_l} \mapsto x_{-k} x_{-l}, e_{\varepsilon_k + \varepsilon_l} \mapsto \partial_{-k} \partial_{-l}, \\
e_{-\delta_i - \delta_j} &\mapsto x_i x_j, e_{-2\delta_i} \mapsto x_i^2, e_{\delta_i + \delta_j} \mapsto \partial_i \partial_j, e_{2\delta_i} \mapsto \partial_i^2, \\
e_{-\varepsilon_k + \delta_i} &\mapsto x_{-k} \partial_i, e_{\varepsilon_k - \delta_i} \mapsto x_i \partial_{-k}, e_{-\varepsilon_k - \delta_i} \mapsto x_{-k} x_i, e_{\varepsilon_k + \delta_i} \mapsto \partial_{-k} \partial_i,
\end{aligned}$$

and the homomorphism $\Psi_{a|b}$ has the form

$$\begin{aligned}
e_{\varepsilon_k - \varepsilon_l} &\mapsto \xi_l \eta_k, e_{-\varepsilon_k - \varepsilon_l} \mapsto \xi_k \xi_l, e_{\varepsilon_k + \varepsilon_l} \mapsto \eta_k \eta_l, \\
e_{-\delta_i - \delta_j} &\mapsto \xi_{-i} \xi_{-j}, e_{-2\delta_i} \mapsto \xi_{-i}^2, e_{\delta_i + \delta_j} \mapsto \eta_{-i} \eta_{-j}, e_{2\delta_i} \mapsto \eta_{-i}^2, \\
e_{-\varepsilon_k + \delta_i} &\mapsto \xi_k \eta_{-i}, e_{\varepsilon_k - \delta_i} \mapsto \eta_k \xi_{-i}, e_{-\varepsilon_k - \delta_i} \mapsto \xi_k \xi_{-i}, e_{\varepsilon_k + \delta_i} \mapsto \eta_k \eta_{-i},
\end{aligned}$$

where $k \neq l$, $i \neq j$.

Lemma 2.1. *The image of $\Phi_{a|b}$ coincides with $D(a|b)_{ev}$ and the image of $\Psi_{a|b}$ coincides with $Cl(a|b)_{ev}$.*

Proof. Let us consider $\Phi_{a|b} : U(\mathfrak{osp}(2b|2a)) \rightarrow D(a|b)$. For any $vw \in S^2 V_{2a|2b}$ we have $\Phi_{a|b}(vw) \in D(a|b)_2 \oplus D(a|b)_0 \oplus D(a|b)_{-2}$. Therefore

$$\Phi_{a|b}(U(\mathfrak{osp}(2b|2a))) \subset D(a|b)_{ev}.$$

Moreover, the above formulas for $\Phi_{a|b}$ show that $\Phi_{a|b}(\mathbb{C} \oplus \mathfrak{osp}(2b|2a))$ is the span of

$$S = \{1, x_i \partial_j, \partial_i \partial_j, x_i x_j \mid -b \leq i, j \leq a\}.$$

By a simple induction argument one shows that S generates $D(a|b)_{ev}$, and the statement follows. A similar argument applies to $\Psi_{a|b}$. \square

By $\mathbb{C}[x]$ we denote the symmetric superalgebra of the superspace with basis $\{x_i \mid -b \leq i \leq a\}$. The superspace $\mathbb{C}[x]$ is a simple faithful $D(a|b)$ -module, and we call it the *defining* $D(a|b)$ -module. Furthermore, $\mathbb{C}[x]_{ev} = \mathbb{C}[x] \cap D(a|b)_{ev}$ is a simple faithful $D(a|b)_{ev}$ -module and hence $\ker \Phi_{a|b}$ is a primitive ideal of $U(\mathfrak{osp}(2b|2a))$. The pullback of $\mathbb{C}[x]_{ev}$ to $U(\mathfrak{osp}(2b|2a))$ is a simple highest weight module of $U(\mathfrak{osp}(2b|2a))$ of highest weight $\frac{1}{2} \left(\sum_{i=1}^b \varepsilon_i - \sum_{j=1}^a \delta_j \right)$ relative to the Borel sub-superalgebra with positive roots

$$\delta_p \pm \delta_q \text{ for } p > q, 2\delta_p, \delta_p \pm \varepsilon_q, \varepsilon_p \pm \varepsilon_q \text{ for } p < q,$$

where the sum $\sum_{i=1}^b \varepsilon_i - \sum_{j=1}^a \delta_j$ is an infinite formal sum if $b = \infty$ or $a = \infty$.

Similarly, the *defining* $Cl(a|b)$ -module $\Lambda[\xi]$ is the exterior superalgebra of the superspace with basis $\{\xi_i \mid -b \leq i \leq a\}$. The module $\Lambda[\xi]$ is a simple and faithful $Cl(a|b)$ -module. Furthermore, $\Lambda[\xi]_{ev} = \Lambda[\xi] \cap Cl(a|b)_{ev}$ is a simple faithful $Cl(a|b)_{ev}$ -module and hence $\ker \Psi_{a|b}$ is a primitive ideal of $U(\mathfrak{osp}(2a|2b))$. The pullback of $\Lambda[\xi]_{ev}$ is a simple highest weight $\mathfrak{osp}(2a|2b)$ -module with highest weight $\frac{1}{2} \left(\sum_{i=1}^b \varepsilon_i - \sum_{j=1}^a \delta_j \right)$, and it is isomorphic to the pullback of $\mathbb{C}[x]_{ev}$. These two isomorphic highest weight modules have purely even highest weight spaces. Next, the pullback of the odd-degree part $\Lambda[\xi]_{odd}$ of $\Lambda[\xi]$ is a simple $\mathfrak{osp}(2a|2b)$ -module with

highest weight $\frac{1}{2} \left(\sum_{i=1}^a \varepsilon_i - \sum_{j=1}^b \delta_j \right) - \delta_1$. The pullbacks of $\Lambda[\xi]_{\text{odd}}$ and $\mathbb{C}[x]_{\text{odd}}$ are isomorphic and have purely odd highest weight spaces.

The pullbacks of $\mathbb{C}[x]_{\text{ev}}$ and $\mathbb{C}[x]_{\text{odd}}$ (equivalently, of $\Lambda[\xi]_{\text{ev}}$ and $\Lambda[\xi]_{\text{odd}}$), together with their counterparts with changed parity, are four pairwise nonisomorphic $\mathfrak{osp}(2a|2b)$ -modules, which we define to be *spinor-oscillator representations*. A general spinor-oscillator representation is the twist of some of these four modules by an automorphism of the Lie superalgebra $\mathfrak{osp}(2a|2b)$. For $b = 0$ (respectively, for $a = 0$) the spinor-oscillator representations are nothing but the spinor representations of $\mathfrak{o}(2a)$ (respectively, the oscillator or Shale-Weil representations of $\mathfrak{sp}(2b)$). (It is well known that for a fixed Borel subalgebra there are precisely two isomorphism classes of purely even spinor or, respectively, oscillator representations.)

The isomorphisms of the pullbacks of $\mathbb{C}[x]_{\text{ev}}$ and $\Lambda[\xi]_{\text{ev}}$ imply the following.

Corollary 2.2. $\ker \Phi_{b|a} = \ker \Psi_{a|b}$ and hence $Cl(a|b)_{\text{ev}}$ and $D(b|a)_{\text{ev}}$ are isomorphic associative superalgebras.

Remark 2.3. It is known that $Cl(a|b)$ is the universal enveloping algebra of the Jordan superalgebra $U_{2a|2b} \oplus \mathbb{C}1$, while $D(a|b)$ is the quotient of the universal enveloping algebra of the Heisenberg superalgebra $V_{2a|2b} \oplus \mathbb{C}z$ by the ideal $(z - 1)$. Furthermore, it is easy to see that the superalgebras $D(b|a)$ and $Cl(a|b)$ are not isomorphic unless $ab = 0$. \circ

Now, we note that $\Psi_{a|b}(\mathfrak{osp}(2a|2b)) \oplus V_{2a|2b}$ is closed under supercommutator, and the corresponding Lie superalgebra is isomorphic to $\mathfrak{osp}(2a + 1|2b)$. Hence we have a surjective homomorphism

$$\Theta_{a|b} : U(\mathfrak{osp}(2a + 1|2b)) \rightarrow Cl(a|b).$$

The explicit formulas for $\Theta_{a|b}$ are the same as those for $\Psi_{a|b}$, with the following addition:

$$e_{\varepsilon_k} \mapsto \eta_k, e_{-\varepsilon_k} \mapsto \xi_k, e_{-\delta_i} \mapsto \xi_{-i}, e_{\delta_i} \mapsto \eta_{-i}.$$

The pullback via $\Theta_{a|b}$ of the defining $Cl(a|b)$ -module $\Lambda[\xi]$ is an irreducible $\mathfrak{osp}(2a + 1|2b)$ -module with highest weight $\frac{1}{2} \left(\sum_{i=1}^a \varepsilon_i - \sum_{j=1}^b \delta_j \right)$ with respect to the Borel subsuperalgebra with positive roots

$$\delta_p \pm \delta_q \text{ for } p > q, \delta_p, 2\delta_p, \delta_p \pm \varepsilon_q, \varepsilon_p \pm \varepsilon_q \text{ for } p < q, \varepsilon_p.$$

We call this highest weight module, together with its counterpart with changed parity, a *spinor-oscillator representation* of $\mathfrak{osp}(2a + 1|2b)$. Moreover, $\ker \Theta_{a|b}$ is the primitive ideal of a spinor-oscillator representation of $\mathfrak{osp}(2a + 1|2b)$.

We note also that $\mathfrak{gl}(a|b)$ is the reductive part of a parabolic subalgebra of $\mathfrak{osp}(2a|2b)$, and by composing the injection $\mathfrak{gl}(a|b) \hookrightarrow \mathfrak{osp}(2a|2b)$ with $\Phi_{b|a}$ we obtain a surjective homomorphism

$$(2.1) \quad U(\mathfrak{gl}(a|b)) \rightarrow D(b|a)_0 \simeq Cl(a|b)_0.$$

Similarly, the embedding $\mathfrak{gl}(a|b) \hookrightarrow \mathfrak{osp}(2b|2a)$ induces a surjective homomorphism

$$(2.2) \quad U(\mathfrak{gl}(a|b)) \rightarrow D(a|b)_0 \simeq Cl(b|a)_0.$$

We denote by $\Upsilon_{a|b}^-$ the restriction of the homomorphism (2.1) to $U(\mathfrak{sl}(a|b))$, and $\Upsilon_{a|b}^+$ the restriction of the homomorphism (2.2) to $U(\mathfrak{sl}(a|b))$.

We will use the homomorphisms $\Upsilon_{a|b}^\pm$ in Section 5.

2.3. Tensor product isomorphisms. Let $Cl^\dagger(a|b)$ (respectively, $D^\dagger(a|b)$) be the superalgebra defined by the same generators and relations as $Cl(a|b)$ (respectively, $D(a|b)$), but where the generators x_i, ∂_i (respectively, ξ_i, η_i) for $i > 0$ are endowed with the opposite parity.

Then one can check that the correspondence $\xi_{-i} \mapsto x_i, \eta_{-i} \mapsto \partial_i, i = 1, \dots, b$, defines an isomorphism of superalgebras

$$(2.3) \quad Cl(0|b) \simeq D^\dagger(b|0),$$

and the correspondence $\xi_i \mapsto x_{-i}, \eta_i \mapsto \partial_{-i}, i = 1, \dots, b$, defines an isomorphism of superalgebras

$$(2.4) \quad Cl^\dagger(b|0) \simeq D(0|b).$$

Lemma 2.4. *We have the following isomorphisms of associative superalgebras*

$$(2.5) \quad D(a|b) \simeq D(a|0) \otimes D(0|b) \simeq D(a|0) \otimes Cl^\dagger(b|0),$$

$$(2.6) \quad Cl(a|b)^\dagger \simeq D(0|a) \otimes D^\dagger(b|0).$$

Proof. The isomorphisms (2.5) follow from (2.4) and from the fact that x_i, ∂_i commute with x_{-j}, ∂_{-j} for all positive i, j . Similarly, the isomorphism (2.6) follows from (2.3) and from the fact that ξ_i, η_i anticommute with ξ_{-j}, η_{-j} for all positive i, j . \square

Corollary 2.5. *We have isomorphisms of (purely even) associative algebras:*

- (a) $D(a|b)_{\bar{0}} \simeq D(a|0) \otimes Cl(b|0)_{ev}, \quad Cl(a|b)_{\bar{0}} \simeq Cl(a|0) \otimes D(b|0)_{ev};$
- (b) $(D(a|b)_{ev})_{\bar{0}} \simeq D(a|0)_{ev} \otimes D(0|b)_{ev}, \quad (Cl(a|b)_{ev})_{\bar{0}} \simeq Cl(a|0)_{ev} \otimes Cl(0|b)_{ev}.$

Proof. Part (a) is a consequence of the existence of isomorphisms $Cl^\dagger(b|0)_{\bar{0}} \simeq Cl(b|0)_{ev}$ and $D^\dagger(b|0)_{\bar{0}} \simeq D(b|0)_{ev}$. Part (b) follows straightforwardly from part (a). \square

2.4. Simple weight modules over Clifford and Weyl algebras. In the rest of the paper, A stands for $D(a|b)$ or $Cl(a|b)$ unless a restriction on A is made explicit. Set $u_i := x_i \partial_i$ ($i \neq 0$) for $A = D(a|b)$, $u_i := \xi_i \eta_i$ ($i \neq 0$) for $A = Cl(a|b)$, and define

$$\mathfrak{h}_A := \text{span}\{u_i \mid i \neq 0\}.$$

Let $\{\zeta_i \mid i \neq 0\} \subset \mathfrak{h}_A^*$ be the system dual to $\{u_i \mid i \neq 0\}$. Then $\mathfrak{h}_A^* = \prod_{i \neq 0} \mathbb{C} \zeta_i$. For convenience, we will write the elements of \mathfrak{h}_A^* as formal (possibly infinite) sums $\sum_{i \neq 0} a_i \zeta_i$. We set

$$Q_A := \bigoplus_{i \neq 0} \mathbb{Z} \zeta_i.$$

One can easily see that the abelian Lie algebra \mathfrak{h}_A acts semisimply on A via the adjoint action. In other words,

$$A = \bigoplus_{\alpha \in R_A \sqcup \{0\}} A^\alpha, \quad A^\alpha = \{x \in A \mid \text{ad}_h(x) = \alpha(h)x \text{ for every } h \in \mathfrak{h}_A\},$$

and R_A is the set of all $\alpha \in Q_A \setminus \{0\}$ such that $A^\alpha \neq 0$. If $A = Cl(a|b)$, then

$$R_A \sqcup \{0\} = \left\{ \sum_{i \neq 0} a_i \zeta_i \in Q_A \mid a_i = 0 \text{ for almost all } i, \text{ and } a_i \in \{0, 1\} \text{ for } i > 0 \right\}.$$

If $A = D(a|b)$, then

$$R_A \sqcup \{0\} = \left\{ \sum_{i \neq 0} a_i \zeta_i \in Q_A \mid a_i = 0 \text{ for almost all } i, \text{ and } a_i \in \{0, 1\} \text{ for } i < 0 \right\}.$$

Moreover, for $A = Cl(a|b)$ we have $\xi_i \in A^{\zeta_i}$, $\eta_i \in A^{-\zeta_i}$ if $i \neq 0$. For $A = D(a|b)$ we have $x_i \in A^{\zeta_i}$, $\partial_i \in A^{-\zeta_i}$ if $i \neq 0$.

Note that each superspace A^α is purely even or purely odd. Define the *parity function on Q_A* to be the homomorphism of abelian groups $p : Q_A \rightarrow \mathbb{Z}_2$ which records the parity of the superspace A^α for $\alpha \in R_A$. Explicitly, $p(\zeta_i) = 0$ for $i > 0$ and $p(\zeta_j) = 1$ for $j < 0$.

- Lemma 2.6.** (a) *The subalgebra $H_A := A^0$ is generated by \mathfrak{h}_A .*
 (b) *If $A = D(a|b)$ then H_A is isomorphic to $\mathbb{C}[u]/(u_i^2 - u_i)_{i < 0}$.*
 (c) *If $A = Cl(a|b)$ then H_A is isomorphic to $\mathbb{C}[u]/(u_i^2 - u_i)_{i > 0}$.*
 (d) *Every root space $0 \neq A^\alpha$ is a cyclic H_A -module.*

Proof. Straightforward computations. □

Set

$$\mathfrak{h}_A^\vee := \{\mu \in \mathfrak{h}_A^* \mid \mu(u_i) = 0, 1 \text{ where } i < 0 \text{ for } A = D(a|b) \text{ and } i > 0 \text{ for } A = Cl(a|b)\}.$$

In what follows, we refer to the elements of \mathfrak{h}_A^\vee as to the *weights of A* . An element μ of \mathfrak{h}_A^\vee is a formal sum

$$\mu = \sum_{i \neq 0} \mu_i \zeta_i$$

with the only restriction that $\mu_i \in \{0, 1\}$ for $i > 0$ if $A = Cl(a|b)$, and $\mu_i \in \{0, 1\}$ for $i < 0$ if $A = D(a|b)$. Note that \mathfrak{h}_A^\vee is not a vector space.

Remark 2.7. Let \mathfrak{g} be a Lie superalgebra isomorphic to $\mathfrak{osp}(2a|2b)$ (respectively, $\mathfrak{osp}(2a+1|2b)$) with fixed Cartan subalgebra \mathfrak{h} . Set $A = Cl(a|b)$ and let $F : U(\mathfrak{g}) \rightarrow A$ be the homomorphism $\Psi_{a|b}$ (respectively, $\Theta_{a|b}$). Then $F(U(\mathfrak{h})) = H_A$. We have

$$\text{Specm } H_A = \mathfrak{h}_A^*, \quad \text{Specm } U(\mathfrak{h}) = \mathfrak{h}^*,$$

where Specm denotes maximal spectrum. Set

$$\tau := \frac{1}{2} \left(\sum_{i>0} \varepsilon_i - \sum_{j>0} \delta_j \right).$$

The map $f : \mathfrak{h}_A^* \rightarrow \mathfrak{h}^*$ induced by F is not linear but affine, i.e.,

$$f(\mu + \nu) = f(\mu) + f(\nu) - f(0)$$

with $f(0) = \tau$. Moreover,

$$f(\zeta_i) = \begin{cases} \varepsilon_i - \tau, & i > 0 \\ \delta_{-i} - \tau, & i < 0 \end{cases}.$$

Similarly if $\mathfrak{g} = \mathfrak{osp}(2b|2a)$, $A = D(a|b)$ and $F := \Phi_{a|b}$, we have

$$f(\zeta_i) = \begin{cases} \varepsilon_{-i} - \tau, & i < 0 \\ \delta_i - \tau, & i > 0 \end{cases}.$$

○

Let \mathbb{C}_μ be the unique $(1|0)$ -dimensional H_A -module on which \mathfrak{h}_A acts via μ . According to Lemma 2.6(a)-(c) every simple H_A -module is one-dimensional and is isomorphic to \mathbb{C}_μ for some $\mu \in \mathfrak{h}_A^\vee$.

An A -module X is a *weight module* if X is semisimple as an H_A -module, i.e., if X has a decomposition

$$X = \bigoplus_{\mu \in \mathfrak{h}_A^\vee} X^\mu,$$

where $X^\mu := \{x \in X \mid hx = \mu(h)x \text{ for every } h \in \mathfrak{h}_A\}$ is the μ -weight space of X . The *support* of a weight module X is

$$\text{supp}X = \{\mu \in \mathfrak{h}_A^\vee \mid X^\mu \neq 0\}.$$

Lemma 2.8. *Let X be a simple weight A -module. Then the weight spaces of X are purely even or purely odd. Hence X and ΠX are never isomorphic.*

Proof. Let $0 \neq x \in (X^\mu)_\kappa$, where $\kappa \in \mathbb{Z}_2$. Then

$$X = Ax = \bigoplus_{\alpha \in Q_A} (A^\alpha x = X^{\alpha+\mu}),$$

i.e., all nonzero vectors in $X^{\alpha+\mu}$ are purely even (respectively, purely odd) if $\kappa + p(\alpha) = \bar{0}$ (respectively, if $\kappa + p(\alpha) = \bar{1}$). \square

For the remainder of the paper we fix an extension of the parity function $p : Q_A \rightarrow \mathbb{Z}_2$ to a map $p : \mathfrak{h}_A^\vee \rightarrow \mathbb{Z}_2$ satisfying $p(\mu + \alpha) = p(\mu) + p(\alpha)$ for any $\alpha \in Q_A$ and any $\mu \in \mathfrak{h}_A^\vee$. Note that such an extension is not unique.

We call a weight A -module X *preferred* if for any $\mu \in \text{supp}X$, the weight space X^μ is purely even if $p(\mu) = \bar{0}$ and the weight space X^μ is purely odd if $p(\mu) = \bar{1}$. Lemma

2.8 implies that, if X is a simple weight module then exactly one of the modules X or ΠX is preferred. Moreover, any weight A -module X decomposes uniquely into a direct sum $X_1 \oplus \Pi X_2$ for some preferred modules X_1 and X_2 .

Proposition 2.9. *The category of preferred weight A^\dagger -modules is equivalent to the category of preferred weight A -modules as an abelian category.*

Proof. The superalgebra A^\dagger has its own parity function $p^\dagger : Q_{A^\dagger} \rightarrow \mathbb{Z}_2$ with the property $p^\dagger(\alpha) = 1$ for $\alpha = \varepsilon_i, \delta_j$. We can extend this function to a map $p^\dagger : \mathfrak{h}_{A^\dagger}^\vee \rightarrow \mathbb{Z}_2$ satisfying $p^\dagger(\mu + \alpha) = p^\dagger(\mu) + p^\dagger(\alpha)$ for any $\alpha \in Q_{A^\dagger}$. Then, for a preferred weight module X we set

$$X^\dagger := \bigoplus_{\mu \in \text{supp} X} \Pi^{p^\dagger(\mu) - p(\mu)} X^\mu.$$

It is clear that \cdot^\dagger is a functor from the category of preferred weight A -modules to the category of preferred weight A^\dagger -modules. Moreover, the functor $(\cdot^\dagger)^\dagger$ is isomorphic to the identity functor. \square

In order to proceed with our study of weight A -modules, for any $\mu \in \mathfrak{h}_A^\vee$ we introduce a certain multiplicity free weight A -module $F(\mu)$ such that $\mu \in \text{supp} F(\mu)$.

First, assume $A = D(a|b)$ and fix $\mu \in \mathfrak{h}_A^\vee$. We can write $\mu = \{\mu_i\}$ with $\mu_i \in \mathbb{C}$ for $i > 0$ and $\mu_i = 0, 1$ for $i < 0$. Let B be the subalgebra in $D(0|b)$ generated by all x_i for $i < 0$ such that $\mu_i = 1$, and by all ∂_i for $i < 0$ such that $\mu_i = 0$. Then B is a local supercommutative algebra, and we denote by J its maximal ideal.

Set $R := \mathbb{C}[x_i, x_i^{-1}]_{i>0}$. Consider the $D(a|0)$ -module $F^+(\mu) := Rx^\mu$ defined by the relations $\partial_i x^\mu = \mu_i x_i^{-1} x^\mu$ and the $D(0|b)$ -module $F^-(\mu) := D(0|b) \otimes_B (B/J)$. Finally using the first isomorphism of (2.5), we define the A -module $F(\mu)$ by setting $F(\mu) := F^+(\mu) \otimes \Pi^{p(\mu)} F^-(\mu)$.

Now let $A = Cl(a|b)$. Here we use the isomorphism (2.6), and set

$$F(\mu) := \Pi^{p(\mu)} (F^-(\mu) \otimes F^+(\mu)^\dagger)^\dagger,$$

where now $F^-(\mu)$ is a $D(0|a)$ -module and $F^+(\mu)$ is a $D(b|0)$ -module.

By construction, $\mu \in \text{supp} F(\mu)$ and all weight spaces of $F(\mu)$ are 1-dimensional.

Lemma 2.10. *The A -module $F(\mu)$ is preferred, indecomposable, and has a simple socle (i.e., a simple submodule which is contained in any nonzero submodule of $F(\mu)$). Under the assumption $\mu_i \notin \mathbb{Z}$ for all $i > 0$ if $A = D(a|b)$, and $\mu_j \notin \mathbb{Z}$ for all $j < 0$ if $A = Cl(a|b)$, the module $F(\mu)$ is simple.*

Proof. Let $A = D(a|b)$. The fact that $F(\mu)$ is preferred follows directly from the definition of $F(\mu)$.

Define the weight $\tilde{\mu} \in \text{supp} F(\mu)$ by setting

$$\tilde{\mu}_i := \begin{cases} \mu_i & \text{if } i < 0 \text{ or } \mu_i \notin \mathbb{Z} \\ 0 & \text{otherwise} \end{cases}.$$

We claim that $F(\mu)^{\tilde{\mu}}$ generates a simple submodule of $F(\mu)$ which is the socle of $F(\mu)$. Indeed, note that if $\nu \in \text{supp}F(\mu)$, the construction of $F(\mu)$ shows that the map $F(\mu)^\nu \rightarrow F(\mu)^{\nu+\varepsilon_i}$ of multiplication by x_i is an isomorphism for all positive i , and that the map $F(\mu)^\nu \rightarrow F(\mu)^{\nu-\varepsilon_i}$ of application of ∂_i is an isomorphism iff $\nu_i \neq 0$. Furthermore, for $i < 0$ the map $F(\mu)^\nu \rightarrow F(\mu)^{\nu+\delta_i}$ of multiplication by x_i is an isomorphism iff $\nu + \delta_i \in \text{supp}F(\mu)$, and similarly the map $F(\mu)^\nu \rightarrow F(\mu)^{\nu-\delta_i}$ of application of ∂_i is an isomorphism iff $\nu - \delta_i \in \text{supp}F(\mu)$. Consequently, the cyclic submodule of $F(\mu)$ generated by any nonzero weight vector contains the weight space $F(\mu)^{\tilde{\mu}}$. This proves our claim, and we see that $F(\mu)$ is indecomposable as it has a simple socle.

Finally, if $\mu_i \notin \mathbb{Z}$ for all $i > 0$ then $\mu = \tilde{\mu}$ and $F(\mu)$ is simple.

The case of $A = Cl(a|b)$ is handled in a similar manner. \square

For $\mu, \nu \in \mathfrak{h}_A^\vee$ we write $\mu \approx \nu$ if $\mu - \nu \in Q_A$ and the respective sets of indices i for which $\mu_i \in \mathbb{Z}_{\geq 0}$ and $\nu_i \in \mathbb{Z}_{\geq 0}$ coincide.

Theorem 2.11. (a) *Every simple weight A -module is multiplicity free.*

(b) *For every $\mu \in \mathfrak{h}_A^\vee$, up to isomorphism, there exist precisely two simple A -modules $X(\mu)$ and $\Pi X(\mu)$ whose supports contain μ , and such that $X(\mu)$ is preferred.*

(c) $\text{supp}X(\mu) = \{\lambda \in \mathfrak{h}_A^\vee \mid \lambda \approx \mu\}$.

(d) *Let $\mu - \nu \in Q_A$. The modules $X(\mu)$ and $X(\nu)$ are isomorphic if and only if $\mu \approx \nu$.*

Proof. Set $P(\mu) := A \otimes_{H_A} (\Pi^{p(\mu)} \mathbb{C}_\mu)$ for $\mu \in \mathfrak{h}_A^\vee$. Then by Frobenius reciprocity $\text{Hom}_A(P(\mu), F(\mu)) \neq 0$. Hence the weight space $P(\mu)^\mu$ is nonzero and generates $P(\mu)$. Since each weight space of $P(\mu)$ is a cyclic H_A -module (Lemma 2.6(d)), the A -module $P(\mu)$ is multiplicity free.

Therefore the sum N of all submodules Z of $P(\mu)$ with $Z^\mu = 0$ constitutes the unique maximal proper submodule of $P(\mu)$. Since $P(\mu)$ is multiplicity free, the quotient $X(\mu) := P(\mu)/N$ and the module $\Pi X(\mu)$ are (up to isomorphism) the only two simple A -modules whose supports contain μ . Note that $X(\mu)$ is preferred, while $\Pi X(\mu)$ is not. This proves (a) and (b).

(c). It follows from (b) that the supports of non-isomorphic simple modules are disjoint. It remains to check that $\text{supp}X(\mu)$ is exactly the equivalence class of μ . Let $A = D(a|b)$ and $v \in X(\mu)^\nu$. Then $x_i v \neq 0$ (respectively, $\partial_i v \neq 0$) iff $\nu + \varepsilon_i \approx \nu$ (respectively, $\nu + \varepsilon_i \approx \nu$). Therefore, $X(\mu)^\nu \neq 0$ iff $\nu \approx \mu$. The case $A = D(a|b)$ is similar.

(d). Direct corollary of (c). \square

Next we would like to decompose the simple weight A -modules in accordance with the isomorphisms (2.5) and (2.6). We start by discussing weight modules of $A = Cl(b|0)$ and $A = D(0|b)$. In these cases we identify the subsets \mathbb{A} of $\mathbb{Z} \cap [1, b]$ (where $b = \infty$ is possible) with the weights of A via the map

$$\mathbb{A} \mapsto \zeta_{\mathbb{A}},$$

where $\zeta_{\mathbb{A}} = \sum_{i \in \mathbb{A}} \zeta_i$ for $A = Cl(b|0)$ and $\zeta_{\mathbb{A}} = \sum_{i \in \mathbb{A}} \zeta_{-i}$ for $A = D(0|b)$. Accordingly, we write $X(\mathbb{A})$ instead of $X(\zeta_{\mathbb{A}})$.

Lemma 2.12. *Let $A(b) = Cl(b|0)$ or $A(b) = D(0|b)$.*

(a) *If $b < \infty$, then the category of preferred weight $A(b)$ -modules is semisimple and has, up to isomorphism, one simple object $X(\emptyset)$.*

(b) *If $b = \infty$ then, up to isomorphism, the simple preferred weight $A(b)$ -modules can be enumerated by equivalence classes of subsets of $\mathbb{Z}_{>0}$ with respect to the following equivalence relation: \mathbb{A} is equivalent to \mathbb{B} if the symmetric difference $\mathbb{A} \Delta \mathbb{B}$ is finite. In other words, up to isomorphism, there is exactly one simple weight $A(b)$ -module $X(\mathbb{A})$ corresponding to \mathbb{A} .*

(c) *We have $X(\mathbb{A}) \simeq X(\mathbb{B})$ if and only if $\mathbb{A} \Delta \mathbb{B}$ is finite.*

Proof. Claim (a) for $A(b) = Cl(b|0)$ is an immediate consequence of the fact that $A(b)$ is a matrix algebra. The case $A(b) = D(0|b)$ with $b < \infty$ follows from Proposition 2.9.

Claim (b) follows from Theorem 2.11(b).

For part (c), we note that $\zeta_{\mathbb{A}} \approx \zeta_{\mathbb{B}}$ if and only if $\mathbb{A} \Delta \mathbb{B}$ is finite. \square

Proposition 2.13. (a) *Every simple preferred weight $D(a|b)$ -module X is isomorphic to $X^+ \otimes (X^-)^\dagger$ for some simple preferred weight $D(a|0)$ -module X^+ and some simple preferred weight $Cl(b|0)$ -module X^- .*

(b) *Every simple preferred weight $Cl(a|b)$ -module X is isomorphic to $((X^+)^\dagger \otimes (X^-)^\dagger)^\dagger$ for some simple preferred weight $Cl(a|0)$ -module X^+ and some simple preferred weight $D(b|0)$ -module X^- .*

Proof. We prove (a) since (b) is similar. For any weight $\mu \in \text{supp} X$ we can choose a simple preferred weight $D(a|0)$ -module X^+ and a simple preferred weight $Cl(b|0)$ -module X^- so that $\mu \in \text{supp} (X^+ \otimes (X^-)^\dagger)$. Moreover, it is clear from the construction that the module $X^+ \otimes (X^-)^\dagger$ is simple. Therefore Theorem 2.11 implies the claim. \square

2.5. Categories of weight modules over Clifford and Weyl algebras. Let \mathcal{W}_A denote the category of preferred weight A -modules. To study the category of all weight A -modules, it suffices to study the category \mathcal{W}_A . Indeed, since every weight A -module decomposes canonically as $X_1 \oplus \Pi X_2$ where X_1 and X_2 are preferred, the morphisms in the category of all weight A -modules are recovered by the morphisms in the category \mathcal{W}_A (the latter morphisms necessarily preserve the \mathbb{Z}_2 -grading).

Recall the A -module $P(\mu)$ introduced in the proof of Theorem 2.11.

Lemma 2.14. *The A -module $P(\mu)$ is an indecomposable projective object in the category \mathcal{W}_A . The category \mathcal{W}_A has enough projectives.*

Proof. By Frobenius reciprocity we have

$$\text{Hom}_A(P(\mu), X) \simeq \text{Hom}_{\mathfrak{h}_A}(\Pi^{p(\mu)} \mathbb{C}_\mu, X) \simeq X^\mu$$

for any preferred module X in \mathcal{W}_A . This implies the projectivity of $P(\mu)$. The indecomposability of $P(\mu)$ follows from the fact that $P(\mu)$ has a unique maximal proper submodule.

Noting that any $X \in \mathcal{W}_A$ is a quotient of $\bigoplus_{\mu \in \text{supp}X} P(\mu) \otimes X^\mu$, we see that \mathcal{W}_A has enough projectives. \square

We introduce the following equivalence relation on the set of weights \mathfrak{h}_A^\vee : $\mu \sim \nu \iff \mu \in \nu + Q_A$. Note that the relation \sim is weaker than the relation \approx , i.e., $\mu \approx \nu$ implies $\mu \sim \nu$. Let Γ denote a \sim -equivalence class in \mathfrak{h}_A^\vee , and let \mathcal{W}_A^Γ be the full subcategory of \mathcal{W}_A with objects X satisfying $\text{supp}X \subset \Gamma$. Since the support of every indecomposable weight A -module X belongs to Γ for some class Γ , we have a decomposition

$$\mathcal{W}_A = \prod \mathcal{W}_A^\Gamma.$$

Proposition 2.15. *The subcategories \mathcal{W}_A^Γ are blocks of \mathcal{W}_A .*

Proof. If X and X' are two simple weight A -modules from \mathcal{W}_A satisfying $\mu \sim \nu$ for some $\mu \in \text{supp}X$ and $\nu \in \text{supp}X'$, then the modules X and X' occur as simple constituents in the A -module $F(\mu)$. We know from Lemma 2.10 that $F(\mu)$ is a preferred indecomposable module. This implies the assertion. \square

Lemma 2.16. *If $A = Cl(a|0)$ or $A = D(0|b)$ then \mathcal{W}_A is a semisimple category.*

Proof. It suffices to prove that every indecomposable projective module $P \in \mathcal{W}_A$ is simple. For this, note that P is an object of \mathcal{W}_A^Γ for some Γ , and let $\mu \in \mathfrak{h}_A^\vee$ belong to $\text{supp}P$. Then $\text{Hom}_{H_A}(\Pi^{p(\mu)}\mathbb{C}_\mu, P) \neq 0$ and Frobenius reciprocity yields a nonzero homomorphism $P \rightarrow P(\mu) = A \otimes_{H_A} \Pi^{p(\mu)}\mathbb{C}_\mu$. The key observation is that under the assumption $A = Cl(a|0)$ or $A = D(0|b)$, the A -module $P(\mu)$ is simple. This together with the projectivity of $P(\mu)$ allows us to conclude that $P \simeq P(\mu)$. \square

The following proposition extends Proposition 2.13 to indecomposable modules.

Proposition 2.17. (a) *If X is an indecomposable module from $\mathcal{W}_{D(a|b)}$, then X is isomorphic to $X^+ \otimes (X^-)^\dagger$ for some indecomposable module $D(a|0)$ -module X^+ from $\mathcal{W}_{D(a|0)}$ and some simple module X^- from $\mathcal{W}_{Cl(b|0)}$.*

(b) *If X is an indecomposable module from $\mathcal{W}_{Cl(a|b)}$, then X is isomorphic to $((X^+)^\dagger \otimes (X^-)^\dagger)^\dagger$ for some simple module X^+ from $\mathcal{W}_{Cl(a|0)}$ and some indecomposable module X^- from $\mathcal{W}_{D(b|0)}$.*

Proof. Let us prove (a). Set $A = D(a|b)$. The indecomposability of X implies $\text{supp}X \subset \mu + Q_A$ for some $\mu \in \mathfrak{h}_A^\vee$. If S and S' are simple subquotients of X then $\text{supp}S \subset \text{supp}S' + Q_A$, and therefore S and S' have the same support when restricted to $D(0|b)$. This, together with Proposition 2.13(a), implies the existence of isomorphisms $S \simeq Y \otimes (X^-)^\dagger$ and $S' \simeq Z \otimes (X^-)^\dagger$ for some simple module

$X^- \in \mathcal{W}_{Cl(b|0)}$ and some simple modules $Y, Z \in \mathcal{W}_{D(a|0)}$. Moreover, according to Lemma 2.16, the restriction of X to $D(0|b)$ is a semisimple $D(0|b)$ -module. Hence this restriction is isomorphic to an isotypic component of the simple $D(0|b)$ -module $(X^-)^\dagger$. This allows us to conclude that the map

$$\mathrm{Hom}_{D(0|b)}((X^-)^\dagger, X) \otimes (X^-)^\dagger \rightarrow X$$

is an isomorphism.

Therefore we can set $X^+ := \mathrm{Hom}_{D(0|b)}((X^-)^\dagger, X)$. Finally, the indecomposability of X implies the indecomposability of X^+ .

The proof of (b) is similar, but instead of Proposition 2.13(a) one uses Proposition 2.13(b). \square

Corollary 2.18. (a) *If $b < \infty$ then the category $\mathcal{W}_{D(a|b)}$ is equivalent to the category $\mathcal{W}_{D(a|0)}$. The category $\mathcal{W}_{D(a|\infty)}$ decomposes into a direct product of subcategories $\mathcal{W}_{[\mathbb{A}]}$ where $[\mathbb{A}]$ runs over equivalence classes of subsets of $\mathbb{Z}_{>0}$ as in Lemma 2.12, and each subcategory $\mathcal{W}_{[\mathbb{A}]}$ is equivalent to the category $\mathcal{W}_{D(a|0)}$.*

(b) *If $a < \infty$ then the category $\mathcal{W}_{Cl(a|b)}$ is equivalent to the category $\mathcal{W}_{D(b|0)}$. The category $\mathcal{W}_{Cl(\infty|b)}$ decomposes into a direct product of subcategories $\mathcal{W}_{[\mathbb{A}]}$ where $[\mathbb{A}]$ runs over equivalence classes of subsets of $\mathbb{Z}_{>0}$ as in Lemma 2.12, and each subcategory $\mathcal{W}_{[\mathbb{A}]}$ is equivalent to the category $\mathcal{W}_{D(b|0)}$.*

(c) *Every block of $\mathcal{W}_{D(a|b)}$ and of $\mathcal{W}_{Cl(a|b)}$ is equivalent to the block $\mathcal{W}_{D(c|0)}^\Gamma$ of $\mathcal{W}_{D(c|0)}$ for some $c \leq \infty$ and $\Gamma = Q_{D(c|0)}$.*

(d) *Two blocks \mathfrak{B}_1 and \mathfrak{B}_2 of $\mathcal{W}_{D(c|0)}$ are equivalent if and only if $c(\mathfrak{B}_1) = c(\mathfrak{B}_2)$ where $c(\mathfrak{B})$ denotes the cardinality of the set of isomorphism classes of simple objects in \mathfrak{B} .*

Proof. Again we prove just (a) since (b) is similar. Let X^- be a preferred simple $Cl(b|0)$ -module and $\mathcal{W}_A(X^-)$ be the full subcategory of \mathcal{W}_A with objects of the form $X^+ \otimes (X^-)^\dagger$ for preferred weight $D(a|0)$ -modules X^+ . It follows from Proposition 2.17 that \mathcal{W}_A is the direct product of its subcategories $\mathcal{W}_A(X^-)$ where X^- runs over the set of isomorphism classes of $Cl(b|0)$ -modules. Each category $\mathcal{W}_A(X^-)$ is equivalent to the category of preferred weight $D(a|0)$ -modules via the functors $\cdot \otimes (X^-)^\dagger$ and $\mathrm{Hom}_{D(0|b)}((X^-)^\dagger, \cdot)$. If $b < \infty$, there is a single isomorphism class to which X^- belongs. If $b = \infty$, the isomorphism classes of modules in $\mathcal{W}_{Cl(b|0)}$ are enumerated by the equivalence classes of subsets of $\mathbb{Z}_{>0}$ from Lemma 2.12.

Parts (c) and (d) follow from parts (a) and (b) and from the classification of blocks in $\mathcal{W}_{D(c|0)}$ for $c < \infty$ in [GS], and in $\mathcal{W}_{D(\infty|0)}$ in [FGM]. \square

We conclude this section by a structural result on indecomposable weight A -modules with finite-dimensional weight spaces.

Theorem 2.19. *Any indecomposable A -module X in \mathcal{W}_A with finite-dimensional weight spaces has a strict A -module filtration $X = \cup_{n \in \mathbb{R}} X_n$ (i.e., $X_n \subsetneq X_m$ for*

$n < m$) for some interval R in \mathbb{Z} , satisfying $\bigcap_{n \in R} X_n = \{0\}$ and such that X_n/X_{n-1} is a simple A -module for any $n, n-1 \in R$.

Proof. Due to Corollary 2.18 we can reduce this statement to the case $A = D(a|0)$. If a is finite then X has finite length and the statement is trivial. For any a , a preferred simple weight $D(a|0)$ -module is determined up to isomorphism by its support. Therefore, if X belongs to a block \mathfrak{B} with $c(\mathfrak{B}) < \infty$, the statement is trivial since X necessarily has finite length.

We can thus assume that X belongs to a block \mathfrak{B} with $c(\mathfrak{B}) = \infty$, and by Corollary 2.18 (c) we can assume further that $\Gamma = Q_A$. Then, simple objects in \mathfrak{B} are enumerated (up to isomorphism) by finite subsets \mathbb{A} of $\mathbb{Z}_{>0}$. For a subset \mathbb{A} , we set $\zeta_{\mathbb{A}} := -\sum_{i \in \mathbb{A}} \zeta_i$ and choose a basis $\{v_i^{\mathbb{A}}\}$ of the weight subspace $X^{\zeta_{\mathbb{A}}}$. Let U be the union of these bases. Note that every cyclic A -module is multiplicity free and has at most countably many cyclic submodules generated by vectors of weights of the form $\zeta_{\mathbb{A}}$. Consider the set \mathcal{X} of cyclic submodules of X consisting of all modules Au for $u \in U$ and all cyclic submodules of Au generated by weight vectors (the weights necessarily having the form $\zeta_{\mathbb{B}}$ for finite subsets \mathbb{B} of $\mathbb{Z}_{>0}$). Then \mathcal{X} is a partially ordered set with respect to the inclusion order. Clearly, $X = \sum_{Y \in \mathcal{X}} Y$.

We claim that any interval in this partial order is finite. To prove this, it suffices to consider an interval of the form $[Av, Aw]$ where $Av \subset Aw$. Let $v \in X^{\zeta_{\mathbb{A}}}$ and $w \in X^{\zeta_{\mathbb{B}}}$ for some finite sets \mathbb{A}, \mathbb{B} . Note that Aw is a quotient of the indecomposable projective A -module $P(\zeta_{\mathbb{B}})$. Therefore

$$v = d \prod_{i \in I} x_i \prod_{j \in J} \partial_j w$$

for some $d \in \mathbb{C}^*$ and some finite subsets $J \subset \mathbb{A}$, $I \subset \mathbb{Z}_{>0} \setminus \mathbb{B}$. If $Av \subset Au \subset Aw$ then

$$u = d' \prod_{i \in I'} x_i \prod_{j \in J'} \partial_j w, \quad v = d'' \prod_{i \in I''} x_i \prod_{j \in J''} \partial_j v.$$

Note that $I = I' \sqcup I''$, $J = J' \sqcup J''$. Since for fixed I, J there exist finitely many choices for I', I'', J', J'' , the claim follows.

Recall that by the Szpilrajn theorem [Mar] any partial order can be extended to a total order. Moreover, we claim that any interval-finite partial order on a countable set I can be extended to an interval-finite total order. Indeed, assume that I does not have a smallest or greatest element. (If I is bounded above or below, the proof is similar). We can choose a sequence of distinct elements $\{x_i \mid i \in \mathbb{Z}\}$ such that if $x_i < x_j$ then $i < j$, and also $I = \cup [x_i, x_{i+1}]$. Let $U_n = \cup_{i=-n}^{n-1} [x_i, x_{i+1}]$ for $n > 0$. Using induction we can define a total order on U_n as required. Indeed, one can see that $U_{n+1} \setminus U_n = Y \cup Z$ where all elements of Z are not less than elements of U_n and all elements of Y are not greater than the elements of U_n . On the other hand, both Y and Z are finite and therefore one can clearly define a suitable total order on them.

This argument endows \mathcal{X} with a total order \prec such that the ordered set (\mathcal{X}, \prec) is isomorphic to $(\mathbb{Z}, <)$, $(\mathbb{Z}_{<0}, <)$, $(\mathbb{Z}_{>0}, <)$, or some finite interval of \mathbb{Z} . We enumerate the elements of \mathcal{X} using this isomorphism. Set $X_n := \sum_{i < n} Y_i$ for $Y_i \in \mathcal{X}$. Let us prove that the A -module X_n/X_{n-1} is simple for any n . Indeed, $X_n/X_{n-1} \simeq Y_n/(Y_n \cap X_{n-1})$. Since $Y_n \cap X_{n-1}$ contains all proper cyclic submodules of Y_n , the submodule $Y_n \cap X_{n-1}$ is the unique maximal submodule of Y_n and the quotient $Y_n/(Y_n \cap X_{n-1})$ is simple. If (\mathcal{X}, \prec) has no minimal element then clearly $\bigcap_n X_n = \{0\}$. If $X_1 = Y_1$ is the minimal element of (\mathcal{X}, \prec) , then X_1 is simple and we add $X_0 := \{0\}$. \square

Example 2.20. Let $A = D(\infty|0)$, $\mu \in Q_A$, and let X be an indecomposable A -module of infinite length with finite weight multiplicities.

- (a) Theorem 2.19 implies that X admits a $\mathbb{Z}_{>0}$ -filtration with simple quotients whenever X has a simple submodule contained in any nonzero submodule of X . Therefore the A -module $F(\mu)$ has such a filtration by Lemma 2.10.
- (b) Similarly, if X has a unique maximal submodule then X admits a $\mathbb{Z}_{<0}$ -filtration with simple quotients. In particular, this applies to the A -module $P(\mu)$.
- (c) Fix an isomorphism $A \simeq A \otimes A$ of associative algebras and consider $X := F(\mu) \otimes P(\mu)$ as an A -module via this isomorphism. One can see that X has neither a simple submodule nor a simple quotient. Nevertheless, by Theorem 2.19 the module X admits a \mathbb{Z} -filtration with simple quotients.

2.6. Weight modules over A_{ev} and $A_{\bar{0}}$ for $A = D(a|b)$ or $A = Cl(a|b)$. Let $\tau : Q_A \rightarrow \mathbb{Z}/2\mathbb{Z}$ be a surjective homomorphism of abelian groups. We define

$$B := \bigoplus_{\tau(\mu)=\bar{0}} A^\mu, \quad B' := \bigoplus_{\tau(\mu)=\bar{1}} A^\mu.$$

Then B is a subsuperalgebra of A containing H_A , and the decomposition $A = B \oplus B'$ defines a $\mathbb{Z}/2\mathbb{Z}$ -grading.

In this subsection we establish an equivalence between the category \mathcal{W}_A and the category \mathcal{W}_B of preferred weight B -modules. This result applies to the particular cases $B = A_{\bar{0}}$ and $B = A_{ev}$. (For $B = A_{\bar{0}}$ preferred B -modules are purely even B -modules.)

The root lattice $Q_B = \ker \tau$ is an index-two subgroup in Q_A . Consider the block \mathcal{W}_A^Γ for some equivalence class $\Gamma \subset \mathfrak{h}_A^\vee$. Note that $\Gamma = \Gamma' \sqcup \Gamma''$, where $\Gamma' := (\mu + Q_B) \cap \Gamma$ for some $\mu \in \Gamma$. This decomposition depends on the choice of μ but only up to swapping Γ' and Γ'' . By $\mathcal{W}_B^{\Gamma'}$ we denote the subcategory of \mathcal{W}_B of B -modules with support in Γ' .

Theorem 2.21. *The abelian categories \mathcal{W}_A^Γ and $\mathcal{W}_B^{\Gamma'}$ are equivalent.*

Proof. We define functors $R : \mathcal{W}_A^\Gamma \rightarrow \mathcal{W}_B^{\Gamma'}$ and $I : \mathcal{W}_B^{\Gamma'} \rightarrow \mathcal{W}_A^\Gamma$ by setting

$$R(X) := \bigoplus_{\mu \in \Gamma'} X^\mu, \quad I(Y) := A \otimes_B Y.$$

We observe that R is exact, I is right exact, and I is left adjoint to R . Therefore, there are canonical morphisms of functors $\phi : IR \rightarrow \text{Id}_{\mathcal{W}_A}$ and $\psi : \text{Id}_{\mathcal{W}_B} \rightarrow RI$. It remains to check that both functors are isomorphisms on objects. Recall that for any $\mu \in \Gamma$ the induced module $P(\mu) = A \otimes_{H_A} (\Pi^{p(\mu)} \mathbb{C}_\mu)$ is projective in \mathcal{W}_A . Similarly, the B -module $Q(\mu) := B \otimes_{H_A} (\Pi^{p(\mu)} \mathbb{C}_\mu)$ is projective in \mathcal{W}_B . By construction we have $I(Q(\mu)) \simeq P(\mu)$ and $R(P(\mu)) \simeq Q(\mu)$. Thus $\phi(P(\mu)) \simeq P(\mu)$ and $\psi(Q(\mu)) \simeq Q(\mu)$. Every object in \mathcal{W}_A (respectively, \mathcal{W}_B) has a resolution with terms given by direct sums of $P(\mu)$ -s (respectively, $Q(\mu)$ -s). Hence ϕ and ψ are isomorphisms on objects. \square

2.7. Weight modules over A_0 . Here we classify simple bounded A_0 -modules. We note that $\mathfrak{h}_A \subset A_0$ and that the root lattice Q_{A_0} is the sublattice of Q_A generated by $\zeta_i - \zeta_j$ for $i, j \neq 0, i \neq j$. As before, we can work with preferred modules only. We introduce a new equivalence relation on \mathfrak{h}^\vee by setting $\mu \approx_0 \nu$ iff $\mu \approx \nu$ and $\mu - \nu \in Q_{A_0}$.

Theorem 2.22. (a) For every $\mu \in \mathfrak{h}_A^\vee$ there exists a unique (up to isomorphism) preferred simple weight module $Y(\mu)$ such that $\mu \in \text{supp}Y(\mu)$.

(b) Two simple preferred A_0 -modules $Y(\mu)$ and $Y(\nu)$ are isomorphic if and only if $\mu \approx_0 \nu$.

Proof. (a) We define the A_0 -module $Y(\mu)$ to be the A_0 -submodule of $X(\mu)$ generated by the weight space $X(\mu)^\mu$. It is simple since for every proper submodule Z of $Y(\mu)$ we have $AZ \cap Y(\mu) = Z$. Furthermore any simple weight A_0 -module, whose support contains μ , is isomorphic to the unique simple quotient of the induced module $A_0 \otimes_{H_A} \mathbb{C}_\mu$. This proves (a).

(b) It follows from (a) that $Y(\mu)$ and $Y(\nu)$ are isomorphic if and only if $\mu \in \text{supp}Y(\nu)$. On the other hand,

$$\text{supp}Y(\nu) = \text{supp}X(\nu) \cap (\nu + Q_{A_0}).$$

This implies the statement. \square

Let $A = D(a|b)$. Any A_0 -module M is also a module over $\text{Lie } A_0$, the Lie superalgebra associated to A_0 . We call M *integrable* if M is integrable as an $\mathfrak{sl}(a|b)$ -module.

Proposition 2.23. (a) A simple weight A_0 -module $Y(\mu)$ is integrable if and only if $\mu_i \in \mathbb{Z}_{\geq 0}$ for all $i > 0$ or $\mu_i \in \mathbb{Z}_{< 0}$ for all $i > 0$.

(b) Every simple weight A_0 -module is integrable as a $D(0|b)_0$ -module.

Proof. (a) By a direct inspection of $\text{supp}Y(\mu)$ one sees that, if μ satisfies the condition of the proposition, then any $\nu \in \text{supp}Y(\mu)$ satisfies the same condition. Therefore the set $(\nu + \mathbb{Z}\alpha) \cap \text{supp}Y(\mu)$ is finite for any $\nu \in \text{supp}Y(\mu)$ and any root α of $\mathfrak{sl}(a|b)$. This implies that $Y(\mu)$ is integrable whenever μ satisfies the condition of the proposition.

On the other hand, if there exist $i, j > 0, i \neq j$, such that μ_j is not an integer or $\mu_i \in \mathbb{Z}_{\geq 0}$ and $\mu_j \in \mathbb{Z}_{< 0}$, then $x_i \partial_j$ acts freely on $Y(\mu)^\mu$.

(b) For any $\mu \in \mathfrak{h}_A^\vee$ and any $\alpha = \zeta_i - \zeta_j$ for $i, j < 0$, at most one of $\mu + \alpha$ and $\mu - \alpha$ lies in \mathfrak{h}_A^\vee . Since the support of any weight A_0 -module is a subset of \mathfrak{h}_A^\vee , the statement follows. \square

Proposition 2.24. *Suppose $A = D(\infty|\infty)$. A simple A_0 -module $Y(\mu)$ is faithful if and only if the set of values*

$$S_i = \{\nu_i \mid \nu \in \text{supp}Y(\mu)\}$$

is infinite at least for one i .

Proof. We first note that (by definition) $\nu_i \in \{0, 1\}$ for all $\nu \in \text{supp}Y(\mu)$ and $i < 0$. If $i > 0$ and $\nu_i \geq m$ for some $m \in \mathbb{Z}_{>0}$, then $(x_j \partial_i)^m Y(\mu)^\nu \neq 0$ for any $j > 0$. Similarly, if $i > 0$ and $\nu_i < -m$ for some $m \in \mathbb{Z}_{>0}$, then $(x_i \partial_j)^m Y(\mu)^\nu \neq 0$ for any $j > 0$. Thus the sets S_i are infinite for all $i > 0$ whenever S_i is infinite for some $i > 0$. Moreover, we observe that the set $T_k := \{(\nu_1, \dots, \nu_k) \mid \nu \in \text{supp}Y(\mu)\}$ is Zariski dense in \mathbb{C}^k for every $k > 0$.

Next we notice that if S_i is finite for some $i > 0$ then $\prod_{s \in S_i} (u_i - s) \in \text{Ann}_{A_0} Y(\mu)$, where $u_i = x_i \partial_i$.

It remains to show that if S_i is infinite for any positive i then $\text{Ann}_{A_0} Y(\mu) = 0$. Clearly, $\text{Ann}_{A_0} Y(\mu)$ is a weight \mathfrak{h} -module with respect to the adjoint action of \mathfrak{h} . Furthermore, for any $u \in A_0^\gamma$ there exists $v \in A_0^{-\gamma}$ such that $uv \neq 0$. Thus, it suffices to prove that $\text{Ann}_{A_0} Y(\mu) \cap H_A = \{0\}$. Any $u \in H_A$ can be written in the form

$$u = p_0(u_1, \dots, u_k) + \sum_{i=1}^l p_i(u_1, \dots, u_k) u_{-i}$$

for some $k, l \in \mathbb{Z}_{>0}$ and polynomials p_0, p_1, \dots, p_l . The condition $u \in \text{Ann}_{A_0} Y(\mu)$ implies $p_i(\nu_1, \dots, \nu_k) = 0$ for all $\nu \in \text{supp}Y(\mu)$ and $i = 1, \dots, l$. Therefore $p_i(T_k) = 0$, and hence $u = 0$. \square

Corollary 2.25. *The ideals $\ker \Upsilon^\pm$ are primitive ideals of $U(\mathfrak{sl}(\infty|\infty))$.*

3. CLASSIFICATION OF SIMPLE BOUNDED WEIGHT \mathfrak{osp} -MODULES AT INFINITY

We are now ready to describe the category of bounded weight \mathfrak{g} -modules for $\mathfrak{g} = \mathfrak{osp}(2a|2b)$, $\mathfrak{osp}(2a+1|2b)$. In what follows we assume that \mathfrak{g} is infinite dimensional, i.e., that at least one of a, b equals ∞ . We fix an exhaustion of \mathfrak{g} as $\varinjlim \mathfrak{g}_k$, where $\mathfrak{g}_k \simeq \mathfrak{osp}(2a_k|2b_k)$ or $\mathfrak{g}_k \simeq \mathfrak{osp}(2a_k+1|2b_k)$, and $a_k, b_k \in \mathbb{Z}_{>0}$ satisfy $a_k = a$ for $a < \infty$ and $b_k = b$ for $b < \infty$.

We start with the following observation.

Proposition 3.1. *If M is a bounded \mathfrak{g} -module, then the restriction of M to $\mathfrak{o}(2a)$ or $\mathfrak{o}(2a+1)$ is integrable and semisimple.*

Proof. M is a bounded semisimple \mathfrak{h} -module, and hence M is a bounded weight $(\mathfrak{o}(2a) + \mathfrak{h})$ - or $(\mathfrak{o}(2a+1) + \mathfrak{h})$ -module. Therefore, as an $\mathfrak{o}(2a)$ - or $\mathfrak{o}(2a+1)$ -module,

M is isomorphic to a direct sum of bounded weight $\mathfrak{o}(2a)$ - or $\mathfrak{o}(2a + 1)$ -modules. As mentioned in Section 1, a bounded weight $\mathfrak{o}(2a)$ - or $\mathfrak{o}(2a + 1)$ -module is integrable for $a = \infty$, and is a sum of finite-dimensional modules for $a < \infty$. Therefore the semisimplicity claim holds trivially for $a < \infty$. For $a = \infty$ the semisimplicity claim follows from Theorem 3.7 in [PSer2]. \square

Recall that an *odd reflection* is the replacement of a Borel subsuperalgebra \mathfrak{b} of \mathfrak{g} by a Borel subsuperalgebra \mathfrak{b}' of \mathfrak{g} such that exactly one odd root α of \mathfrak{b} is not a root of \mathfrak{b}' (and hence $-\alpha$ is a root of \mathfrak{b}'). If $L_{\mathfrak{b}}(\lambda)$ denotes an irreducible \mathfrak{g} -module with \mathfrak{b} -highest weight λ and purely even highest-weight vector, then $L_{\mathfrak{b}}(\lambda)$ is isomorphic either to $L_{\mathfrak{b}'}(\lambda)$ or to $\Pi L_{\mathfrak{b}'}(\lambda - \alpha)$. The latter case, called a *typical* reflection, occurs precisely when $(\lambda, \alpha) \neq 0$, while the former case, called an *atypical* reflection, occurs when $(\lambda, \alpha) = 0$.

By $J_{\mathfrak{g}}$ we denote the kernel of $\Psi_{a|b}$ if $\mathfrak{g} = \mathfrak{osp}(2a|2b)$, and respectively of $\Theta_{a|b}$ if $\mathfrak{g} = \mathfrak{osp}(2a + 1|2b)$. Recall that $J_{\mathfrak{g}}$ is the annihilator of any spinor-oscillator representation. Moreover, it is obvious that $J_{\mathfrak{g}} = \varinjlim J_{\mathfrak{g}_k}$ whenever $\mathfrak{g} = \varinjlim \mathfrak{g}_k$ for an inductive system of finite-dimensional Lie superalgebras \mathfrak{g}_k of type \mathfrak{osp} .

Lemma 3.2. *Let $\mathfrak{q} = \mathfrak{osp}(m|2n)$ for $m, n \in \mathbb{Z}_{\geq 0}$, and $I \subset U(\mathfrak{q})$ be a bounded primitive ideal of degree d . Assume that at least one of the simple ideals of $\mathfrak{q}_{\bar{0}}$ has rank greater than d . Then $d = 1$. Moreover $I = J_{\mathfrak{q}}$, unless I is the augmentation ideal or the annihilator of a defining module.*

Proof. For $m \leq 1$ the statement follows directly from Lemmas 1.5 and 1.6. Therefore in the rest of the proof we assume that $m \geq 2$.

By Musson's Theorem [M], $I = \text{Ann}_{U(\mathfrak{q})} L_{\mathfrak{b}}(\lambda)$ for some Borel subsuperalgebra \mathfrak{b} and some weight λ . For $\lambda = 0$, the ideal I is the augmentation ideal. For the rest of the proof we assume $\lambda \neq 0$. Let \mathfrak{s} be a simple ideal of $\mathfrak{q}_{\bar{0}}$ of rank greater than $d + 1$. We can choose the Borel subsuperalgebra \mathfrak{b} so that its base of simple roots contains a base of simple roots for \mathfrak{s} . By μ we denote the weight of \mathfrak{s} obtained from λ by restriction.

In order to study the annihilator I of the simple highest weight \mathfrak{q} -module $L_{\mathfrak{b}}(\lambda)$, we will consider $L_{\mathfrak{b}}(\lambda)$ as a highest weight module over a variable Borel subalgebra \mathfrak{b}' obtained from \mathfrak{b} by some sequence of odd reflections. Then λ' will denote the corresponding highest weight, and μ' will be its restriction to \mathfrak{s} . Lemma 1.4 implies that the simple \mathfrak{s} -modules with highest weights μ and μ' are necessarily multiplicity free.

We may assume that \mathfrak{b}' is obtained from \mathfrak{b} by odd reflections with respect to some isotropic odd roots $\alpha_1, \dots, \alpha_r$. It is essential to note that there are at most four nonisomorphic multiplicity free simple weight \mathfrak{s} -modules which have a highest weight with respect to a fixed Borel subalgebra of \mathfrak{s} . (Indeed, these are the trivial, natural, and spinor modules for $\mathfrak{s} \simeq \mathfrak{o}(m)$, and the trivial, natural, and oscillator modules for $\mathfrak{s} \simeq \mathfrak{sp}(2n)$.) This shows that each of the weights μ and μ' can take at most four

different values. Moreover, since λ, λ' have the same image modulo the root lattice of \mathfrak{q} , it is easy to check that for a given μ there is a unique μ' with $\mu' \neq \mu$. Therefore in a shortest chain of odd reflections connecting \mathfrak{b} and \mathfrak{b}' there can be at most one typical reflection.

Assume $\mathfrak{s} = \mathfrak{sp}(2n)$. If $m = 2\ell + 1$ we fix the simple roots

$$\varepsilon_1 - \varepsilon_2, \dots, \varepsilon_\ell - \delta_1, \dots, \delta_{n-1} - \delta_n, \delta_n,$$

and if $m = 2\ell$ we take the simple roots

$$\varepsilon_1 - \varepsilon_2, \dots, \varepsilon_\ell - \delta_1, \dots, \delta_{n-1} - \delta_n, 2\delta_n.$$

Set $\lambda = a_1\varepsilon_1 + \dots + a_\ell\varepsilon_\ell + \mu$. The above conditions and Lemma 1.6 show that for $\mu \neq \mu'$ one of the following holds:

- (1) $\mu = 0, \mu' = \delta_1,$
- (2) $\mu = \delta_1, \mu' = 0,$
- (3) $\mu = -\frac{1}{2}(\delta_1 + \dots + \delta_n), \mu' = -\frac{1}{2}(\delta_1 + \dots + \delta_{n-1}) - \frac{3}{2}\delta_n,$
- (4) $\mu = -\frac{1}{2}(\delta_1 + \dots + \delta_{n-1}) - \frac{3}{2}\delta_n, \mu' = -\frac{1}{2}(\delta_1 + \dots + \delta_n).$

Consider the first case. We start by applying the odd reflections corresponding to the sequence of odd roots $\varepsilon_\ell - \delta_1, \dots, \varepsilon_1 - \delta_1$. Since $\lambda \neq 0$, exactly one of these reflections must be typical, say with respect to $\varepsilon_p - \delta_1$. This implies $a_{p+1} = \dots = a_\ell = 0, a_1 = \dots = a_{p-1} = -1$. Next, an application of the reflections corresponding to $\varepsilon_\ell - \delta_2, \dots, \varepsilon_1 - \delta_2$ cannot change λ' . This is only possible for $p = 1$ and $\lambda = \varepsilon_1$, and then $L_{\mathfrak{b}}(\lambda)$ is a defining representation.

Let us deal with the second case. The odd reflections with respect to the roots $\varepsilon_\ell - \delta_1, \dots, \varepsilon_1 - \delta_1$ do not change λ , i.e., they are all atypical. Therefore $a_1 = \dots = a_\ell = -1$, but then the reflection with respect to $\varepsilon_\ell - \delta_2$ is typical and $\mu' = \delta_1 + \delta_2$. This proves that the second case is impossible.

Now, consider the third case. Here we perform in some order all odd reflections with roots $\varepsilon_i - \delta_j, i = 1, \dots, \ell, j = 1, \dots, n-1$, and check that all these reflections do not change λ . This forces $a_1 = \dots = a_\ell = \frac{1}{2}$. Hence $\lambda = \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_\ell) - \frac{1}{2}(\delta_1 + \dots + \delta_n)$ and $L_{\mathfrak{b}}(\lambda)$ is a spinor-oscillator representation.

Finally, let us look at the fourth case. We can show that $a_1 = \dots = a_\ell = \frac{1}{2}$ in the same way as in the third case. Therefore, if m is even we have $\lambda = \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_\ell) - \frac{1}{2}(\delta_1 + \dots + \delta_{n-1}) - \frac{3}{2}\delta_n$, and $L_{\mathfrak{b}}(\lambda)$ is a spinor-oscillator representation not isomorphic up to parity change to a spinor-oscillator representation that occurred in the third case. If m is odd, then by Lemma 1.5 the restriction of λ to $\mathfrak{osp}(1|2n)$ with roots $\pm\delta_i \pm \delta_j, \delta_r - \delta_s, \pm\delta_i$ for $r \neq s$, must equal $-\frac{1}{2}(\delta_1 + \dots + \delta_n)$. This contradicts our assumption for μ , therefore the fourth case forces m to be even.

This proves our claim for $\mathfrak{s} = \mathfrak{sp}(2n)$ since in case (1) I is the annihilator of a defining representation, while in cases (3) and (4) I is the annihilator of a spinor-oscillator representation.

We conclude the proof by essentially repeating the above argument for $\mathfrak{s} = \mathfrak{o}(m)$. For $m = 2\ell$ we fix the simple roots

$$\delta_1 - \delta_2, \dots, \delta_{n-1} - \delta_n, \delta_n - \varepsilon_1, \varepsilon_1 - \varepsilon_2, \dots, \varepsilon_{\ell-1} - \varepsilon_\ell, \varepsilon_{\ell-1} + \varepsilon_\ell,$$

and for $m = 2\ell + 1$ we choose the simple roots

$$\delta_1 - \delta_2, \dots, \delta_{n-1} - \delta_n, \delta_n - \varepsilon_1, \varepsilon_1 - \varepsilon_2, \dots, \varepsilon_{\ell-1} - \varepsilon_\ell, \varepsilon_\ell.$$

A priori there are the following cases for $\mu \neq \mu'$:

- (1) $\mu = 0, \mu' = \varepsilon_1,$
- (2) $\mu = \varepsilon_1, \mu' = 0,$
- (3) $\mu = \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_\ell), \mu' = \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_{\ell-1} - \varepsilon_\ell),$
- (4) $\mu = \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_{\ell-1} - \varepsilon_\ell), \mu' = \frac{1}{2}(\varepsilon_1 + \dots + \varepsilon_\ell).$

All these cases can be treated in the same way as above. \square

Corollary 3.3. *Let \mathfrak{q} and I be as in the previous lemma. Then the superalgebra of \mathfrak{h} -invariants $(U(\mathfrak{q})/I)^{\mathfrak{h}}$ is abelian. Hence any weight \mathfrak{q} -module generated by a single weight vector is multiplicity free.*

We are now ready to prove the following.

Proposition 3.4. *Let M be a simple bounded \mathfrak{g} -module. Then M is multiplicity free. Moreover, M satisfies $\text{Ann}_{U(\mathfrak{g})}M = J_{\mathfrak{g}}$ or M is a trivial or a natural module.*

Proof. Let $I := \text{Ann}M$, $\bar{U} := U(\mathfrak{g})/I$, and let λ be a weight of M . Then a standard argument shows that M^λ is a simple $\bar{U}^{\mathfrak{h}}$ -module. Next, set

$$\mathfrak{h}_k := \mathfrak{g}_k \cap \mathfrak{h}, \quad \bar{U}_k := U(\mathfrak{g}_k)/(U(\mathfrak{g}_k) \cap I).$$

We have $\bar{U}^{\mathfrak{h}} = \varinjlim \bar{U}_k^{\mathfrak{h}_k}$. Since \mathfrak{g} is infinite dimensional, Lemma 3.2 implies that for sufficiently large k the simple $\bar{U}_k^{\mathfrak{h}_k}$ -constituents of the module M^λ are one-dimensional. By passing to the direct limit we obtain $\dim M^\lambda = 1$. Furthermore, again by Lemma 3.2 we see that the annihilator of $U(\mathfrak{g}_k)M^\lambda$ equals $J_{\mathfrak{g}_k}$, unless $U(\mathfrak{g}_k)M^\lambda$ is a trivial representation or a defining representation. The statement follows by passing to the direct limit. \square

Remark 3.5. The claim of Proposition 3.4 is proved in [GP] in the case where $\mathfrak{g} = \mathfrak{g}_{\bar{0}}$, i.e., for $\mathfrak{g} = \mathfrak{sp}(\infty), \mathfrak{o}(\infty)$. \circ

We say that a simple weight \mathfrak{g} -module M is of *spinor-oscillator type* if it is annihilated by $J_{\mathfrak{g}}$, i.e., M is obtained by pullback along the homomorphism $\Theta_{a,b}$ from a weight $Cl(a|b)$ -module or, respectively, along the homomorphism $\Psi_{a,b}$ from a weight $Cl(a|b)_{ev}$ -module. Proposition 3.4 implies the following.

Corollary 3.6. *Let M be a simple bounded weight \mathfrak{g} -module such that $M \not\cong V, \text{IV}, \mathbb{C}, \text{II}\mathbb{C}$. Then M is of spinor-oscillator type.*

Note that every simple weight $\mathfrak{sp}(2b)$ -module T of oscillator type (as defined in Section 1) is the pullback of a (unique, up to isomorphism) simple weight $Cl(0|b)_{ev}$ -module \tilde{T} . This follows from the fact that the ideal $\ker \Psi_{0,b}$ of $U(\mathfrak{sp}(2b))$ is the primitive ideal not only of the oscillator representations but of any simple multiplicity free weight module of $\mathfrak{sp}(2b)$. For $b < \infty$ this is well known, and for $b = \infty$ see [GP].

Given T as above, the module \tilde{T} generates a unique simple weight $Cl(0|b)$ -module which has the form $\tilde{T} \oplus \tilde{T}'$ as a $Cl(0|b)_{ev}$ -module. The pullback of \tilde{T}' to $\mathfrak{sp}(2b)$ is by definition the *twin* of T and is a simple module. Similarly, any spinor $\mathfrak{o}(2a)$ -module is the pullback of a simple weight $Cl(a|0)_{ev}$ -module, and we call two spinor $\mathfrak{o}(2a)$ -modules *twins* if they are pullbacks of the two simple $Cl(a|0)_{ev}$ -constituents of a simple $Cl(a|0)$ -module. For $\mathfrak{o}(2a+1)$ we declare two spinor $\mathfrak{o}(2a+1)$ -modules to be *twins* if they are isomorphic.

We are ready to state our explicit description of simple bounded weight \mathfrak{g} -modules.

Theorem 3.7. *Let M be a simple bounded weight \mathfrak{g} -module of spinor-oscillator type. Then the following statements hold.*

- (a) $M_{\bar{0}}$ and $M_{\bar{1}}$ are simple $\mathfrak{g}_{\bar{0}}$ -modules.
- (b) There exist twin spinor $\mathfrak{o}(2a)$ - or $\mathfrak{o}(2a+1)$ -modules S and S' , and twin simple $\mathfrak{sp}(2b)$ -modules T and T' of oscillator type, such that

$$(3.1) \quad M_{\bar{0}} \simeq S \otimes T, \quad M_{\bar{1}} \simeq \Pi(S' \otimes T').$$

The modules S, S', T, T' are unique up to isomorphism and determine M up to isomorphism.

- (c) Any pair (S, T) where S is a spinor $\mathfrak{o}(2a)$ - or $\mathfrak{o}(2a+1)$ -module and T is a simple $\mathfrak{sp}(2b)$ -module of oscillator type determines a simple bounded weight \mathfrak{g} -module M of spinor-oscillator type for which (3.1) holds.

Proof. Let $A = Cl(a|b)$. Claim (a) follows directly from Lemma 1.1 since the map $\Psi_{a|b} : U(\mathfrak{g}_{\bar{0}}) \rightarrow (A_{ev})_{\bar{0}}$ (respectively, $\Theta_{a|b} : U(\mathfrak{g}_{\bar{0}}) \rightarrow A_{\bar{0}}$) is surjective.

(b) Note that if the statement holds for M then it holds for ΠM .

If $\mathfrak{g} = \mathfrak{osp}(2a+1|2b)$ then we can assume that M is the pullback of a simple preferred weight $Cl(a|b)$ -module X . By Proposition 2.13(b) there is an isomorphism $X \simeq ((X^+)^\dagger \otimes (X^-)^\dagger)^\dagger$ for some simple preferred weight $Cl(a|0)$ -module X^+ and some simple preferred weight $D(b|0)$ -module X^- . Next, using the isomorphism $Cl(a|b)_{\bar{0}} \simeq Cl(a|0) \otimes Cl(0|b)_{ev}$ from Corollary 2.5 we see that $X_{\bar{0}} \simeq X^+ \otimes R(X^-)$ and $X_{\bar{1}} \simeq X^+ \otimes (X^-/R(X^-))$ where the functor R is defined in Section 2.6. Thus, $S = S'$ is isomorphic to the pullback to $\mathfrak{o}(2a+1)$ of X^+ while T and T' are isomorphic to the pullbacks to $\mathfrak{sp}(2b)$ of $R(X^-)$ and $X^-/R(X^-)$, respectively.

Now let $\mathfrak{g} = \mathfrak{osp}(2a|2b)$. We can assume that M is the pullback of $R(X)$ for a simple preferred weight $Cl(a|b)_{ev}$ -module X . Then

$$R(X)_{\bar{0}} \simeq R(X^+) \otimes R(X^-), \quad R(X)_{\bar{1}} \simeq (R(X^+)/X^+) \otimes (X^-/R(X^-)).$$

Therefore S and S' are isomorphic to the respective pullbacks to $\mathfrak{o}(2a)$ of $R(X^+)$ and $(R(X^+)/X^+)$, and T and T' are the same as in the case of $\mathfrak{osp}(2a+1|2b)$.

The uniqueness of S and T , and hence also of S' and T' , is clear from the isomorphism of $\mathfrak{g}_{\bar{0}}$ -modules $M_{\bar{0}} \simeq S \otimes T$. The fact that S, S', T, T' determine M up to isomorphism is a consequence of the observation that $M_{\bar{0}}$ determines $R(X)_{\bar{0}}$, which in turn determines X^+ and $R(X^-)$ for $\mathfrak{g} = \mathfrak{osp}(2a+1|2b)$ (respectively, $R(X^+)$ and $R(X^-)$ for $\mathfrak{g} = \mathfrak{osp}(2a|2b)$), and ultimately X^+ and X^- since R is an equivalence of categories. Then M is the pullback of $((X^+)^\dagger \otimes (X^-)^\dagger)^\dagger$ for $\mathfrak{g} = \mathfrak{osp}(2a+1|2b)$ and of $R((X^+)^\dagger \otimes (X^-)^\dagger)^\dagger$ for $\mathfrak{g} = \mathfrak{osp}(2a|2b)$.

(c) The given pair (S, T) determines a pair (X^+, X^-) , where S is the pullback of a simple weight $Cl(a|0)$ -module X^+ and T is the pullback of a simple weight $Cl(0|b)$ -module $(X^-)^\dagger$ for a simple weight $D(b|0)$ -module X^- . Then M is recovered from X^+ and X^- as in the proof of part (b). \square

Remark 3.8. There is an alternative definition of pairs of twins (S, S') or (T, T') in terms of the supports of the weight modules S and T . Recall that in [GP] the supports of all simple bounded (equivalently, multiplicity free) weight $\mathfrak{o}(\infty)$ - and $\mathfrak{sp}(\infty)$ -modules are described explicitly, and moreover a given such module is determined up to isomorphism by its support. For a finite-dimensional orthogonal or symplectic Lie algebra it is well known that a simple multiplicity free weight module is determined by its support as well. Both if $a < \infty$ or $a = \infty$, for any spinor $\mathfrak{o}(2a)$ -module S there exists a unique (up to isomorphism) spinor module S' such that for every $i \in \mathbb{Z}_{>0}$, $\mu + \varepsilon_i \in \text{supp}S'$ for some $\mu \in \text{supp}S$. Similarly, if $b < \infty$ or $b = \infty$, for every $\mathfrak{sp}(2b)$ -module T of oscillator type there exists a unique (up to isomorphism) module T' of oscillator type such that for every $i \in \mathbb{Z}_{>0}$, $\nu + \varepsilon_i \in \text{supp}S'$ for some $\nu \in \text{supp}S$. It is straightforward to show that the pairs (S, S') and (T, T') are precisely the pairs of twins defined above. This observation leads to another proof of Theorem 3.7(b) based on analyzing the supports of the $\mathfrak{g}_{\bar{0}}$ -modules $M_{\bar{0}}$ and $M_{\bar{1}}$. \circ

Consider the decomposition $\mathfrak{g}_{\bar{0}} = \mathfrak{g}_{\mathfrak{o}} \oplus \mathfrak{g}_{\mathfrak{sp}}$, where $\mathfrak{g}_{\mathfrak{o}} \simeq \mathfrak{o}(2a)$ or $\mathfrak{g}_{\mathfrak{o}} \simeq \mathfrak{o}(2a+1)$ and $\mathfrak{g}_{\mathfrak{sp}} \simeq \mathfrak{sp}(2b)$. Set $\mathfrak{h}_{\mathfrak{o}} = \mathfrak{h} \cap \mathfrak{g}_{\mathfrak{o}}$ and $\mathfrak{h}_{\mathfrak{sp}} = \mathfrak{h} \cap \mathfrak{g}_{\mathfrak{sp}}$. Then $\mathfrak{h}^* = \mathfrak{h}_{\mathfrak{o}}^* \oplus \mathfrak{h}_{\mathfrak{sp}}^*$. Moreover, if $\Gamma_{\mathfrak{o}} \subset \mathfrak{h}_{\mathfrak{o}}^*$ and $\Gamma_{\mathfrak{sp}} \subset \mathfrak{h}_{\mathfrak{sp}}^*$ we put $\Gamma_{\mathfrak{o}} + \Gamma_{\mathfrak{sp}} := \{\gamma_1 + \gamma_2 \mid \gamma_1 \in \Gamma_{\mathfrak{o}}, \gamma_2 \in \Gamma_{\mathfrak{sp}}\}$.

Corollary 3.9. *Let M be as in Theorem 3.7. Then*

$$\text{supp}M = (\text{supp}S \sqcup \text{supp}S') + (\text{supp}T \sqcup \text{supp}T') \subset \mathfrak{h}_{\mathfrak{o}}^* \oplus \mathfrak{h}_{\mathfrak{sp}}^*.$$

Moreover M is never isomorphic to ΠM , and $\text{supp}M$ determines the isomorphism class of M up to application of Π .

Remark 3.10. The pairs $(M, \Pi M)$ for \mathfrak{g} are appropriate superanalogs of twin pairs for $\mathfrak{o}(2a)$ or $\mathfrak{sp}(2b)$. \circ

4. ON THE CATEGORY OF BOUNDED WEIGHT \mathfrak{osp} -MODULES

Now we turn our attention to the category $\mathcal{B}_{\mathfrak{g}}$ of bounded \mathfrak{g} -modules. In this section, \mathfrak{g} stands for $\mathfrak{osp}(2a+1|2b)$ or $\mathfrak{osp}(2a|2b)$ for all, possibly finite, a and b .

Let $\mathcal{B}_{\mathfrak{g}}^{osc}$ denote the full subcategory of $\mathcal{B}_{\mathfrak{g}}$ with simple objects of spinor-oscillator type. Every $M \in \mathcal{B}_{\mathfrak{g}}$ decomposes uniquely into a direct sum $M' \oplus M''$ with $M' \in \mathcal{B}_{\mathfrak{g}}^{osc}$ and M'' being a direct sum of finitely many copies of trivial and defining modules. This follows from a simple inspection of supports which shows that any simple subquotient of M isomorphic to $V, \Pi V, \mathbb{C}, \Pi \mathbb{C}$ splits as a direct summand of M . By $\widetilde{\mathcal{W}}_A^{fin}$ for $\mathfrak{g} = \mathfrak{osp}(2a+1|2b)$ (respectively, $\widetilde{\mathcal{W}}_{A_{ev}}^{fin}$ for $\mathfrak{g} = \mathfrak{osp}(2a|2b)$) we denote the category of all weight A -modules (respectively, A_{ev} -modules) with finite weight multiplicities. Note that the objects of $\widetilde{\mathcal{W}}_A^{fin}$ (respectively, $\widetilde{\mathcal{W}}_{A_{ev}}^{fin}$) are not necessarily preferred A -modules (respectively, A_{ev} -modules).

Observe that, if a and b are finite then the indecomposable modules in $\mathcal{B}_{\mathfrak{g}}^{osc}$ have finite length. Indeed, the support of every such module M lies in a single coset of the root lattice of \mathfrak{g} . Since the root lattice of \mathfrak{g}_0 has index 2 in the root lattice of \mathfrak{g} , the support of M over \mathfrak{g}_0 lies in at most 2 cosets of the root lattice of \mathfrak{g}_0 . As a consequence, M has finite length as a \mathfrak{g}_0 -module by Lemma 3.3 in [Mat].

The following is our first main result about the category $\mathcal{B}_{\mathfrak{g}}^{osc}$.

Theorem 4.1. *Let $A = Cl(a|b)$ for $b \neq 1$. If $\mathfrak{g} = \mathfrak{osp}(2a+1|2b)$ then the category $\mathcal{B}_{\mathfrak{g}}^{osc}$ is equivalent to the category $\widetilde{\mathcal{W}}_A^{fin}$. If $\mathfrak{g} = \mathfrak{osp}(2a|2b)$ then the category $\mathcal{B}_{\mathfrak{g}}^{osc}$ is equivalent to the category $\widetilde{\mathcal{W}}_{A_{ev}}^{fin}$.*

As a first step we prove Theorem 4.1 for finite a and b .

Lemma 4.2. *Let $\dim \mathfrak{g} < \infty$. Then the restriction map $\text{Ext}_{\mathfrak{g}, \mathfrak{h}}^1(M, N) \rightarrow \text{Ext}_{\mathfrak{g}_0, \mathfrak{h}}^1(M, N)$ is injective.*

Proof. We have to show that any exact sequence in $\mathcal{B}_{\mathfrak{g}}^{osc}$

$$0 \rightarrow N \rightarrow R \rightarrow M \rightarrow 0$$

which splits over \mathfrak{g}_0 splits also over \mathfrak{g} . It suffices to show that $H^1(\mathfrak{g}, \mathfrak{g}_0; \text{Hom}(M, N))_{\bar{0}} = 0$, where Hom stands for the homomorphisms of vector spaces disregarding the \mathbb{Z}_2 -grading, see §3.1 and §4.5 of [Fuks]. Any indecomposable object in $\mathcal{B}_{\mathfrak{g}}^{osc}$ has finite length and therefore it is enough to prove that this cohomology vanishes for simple M and N . Writing down the first three terms of the complex computing relative cohomology, we have

$$0 \rightarrow \text{Hom}_{\mathfrak{g}_0}^0(M, N) \xrightarrow{d} \text{Hom}_{\mathfrak{g}_0}^0(\mathfrak{g}_{\bar{1}} \otimes M, N) \rightarrow \text{Hom}_{\mathfrak{g}_0}^0(\Lambda^2 \mathfrak{g}_{\bar{1}} \otimes M, N) \rightarrow \dots,$$

where $\text{Hom}_{\mathfrak{g}_0}^0$ denotes homomorphisms of \mathfrak{g}_0 -modules preserving the \mathbb{Z}_2 -grading. Note that the second term of the complex does not vanish if and only if $\text{supp} M_{\bar{1}} \cap (\text{supp} N_{\bar{0}} + \Delta_{\bar{1}})$ or $\text{supp} M_{\bar{0}} \cap (\text{supp} N_{\bar{1}} + \Delta_{\bar{1}})$ is non-empty. Using Theorem 3.7 we see that this

can happen if and only if $M \simeq N$. In the latter case

$$\mathrm{Hom}_{\mathfrak{g}_0}^0(\mathfrak{g}_1 \otimes M, M) = \mathrm{Hom}_{\mathfrak{g}_0}^0(\mathfrak{g}_1 \otimes M_{\bar{0}}, M_{\bar{1}}) \oplus \mathrm{Hom}_{\mathfrak{g}_0}^0(\mathfrak{g}_1 \otimes M_{\bar{1}}, M_{\bar{0}}) = \mathbb{C}^2$$

and

$$\mathrm{End}_{\mathfrak{g}_0}^0(M) = \mathrm{End}_{\mathfrak{g}_0}^0(M_{\bar{0}}) \oplus \mathrm{End}_{\mathfrak{g}_0}^0(M_{\bar{1}}) = \mathbb{C}^2.$$

Consider $\varphi_0 \in \mathrm{Hom}_{\mathfrak{g}_0}^0(\mathfrak{g}_1 \otimes M_{\bar{0}}, M_{\bar{1}})$ and $\varphi_1 \in \mathrm{Hom}_{\mathfrak{g}_0}^0(\mathfrak{g}_1 \otimes M_{\bar{1}}, M_{\bar{0}})$ defined by the formula $\varphi_i(g \otimes m) = gm$ where $g \in \mathfrak{g}_1$ and $m \in M_i$. Set

$$\psi_i(m) := \begin{cases} m & \text{if } m \in M_i \\ 0 & \text{if } m \notin M_i \end{cases}.$$

Then $\varphi_i = d(\psi_i)$. Hence $H^1(\mathfrak{g}, \mathfrak{g}_0; \mathrm{End}(M))_{\bar{0}} = 0$. \square

For $\mathfrak{g}_0 = \mathfrak{g}_o \oplus \mathfrak{g}_{sp}$ we say that a simple module Z has *spinor-oscillator type* if $Z \simeq S \otimes T$ for some spinor \mathfrak{g}_o -module S and some \mathfrak{g}_{sp} -module T of oscillator type. By $\mathcal{B}_{\mathfrak{g}_0}^{osc}$ we denote the category of \mathbb{Z}_2 -graded bounded weight \mathfrak{g}_0 -modules with simple constituents of spinor-oscillator type.

Corollary 4.3. *Theorem 4.1 holds in the case $\dim \mathfrak{g} < \infty$ (and $b \neq 1$).*

Proof. Note that if $b = 0$ the statement is trivial since $\mathcal{B}_{\mathfrak{g}}^{osc}$ is a semisimple category with objects that are finite direct sums of (finite-dimensional) spinor modules. Next, for $\mathfrak{g} = \mathfrak{sp}(2b)$ with $1 < b < \infty$ the statement is proven in [GS]. Therefore, if $\mathfrak{g} = \mathfrak{osp}(2a + 1|2b)$ (respectively, $\mathfrak{g} = \mathfrak{osp}(2a|2b)$), we have an equivalence of the categories $\mathcal{B}_{\mathfrak{g}_0}^{osc}$ and $\widetilde{\mathcal{W}}_{A_{\bar{0}}}^{fin}$ (respectively, $\widetilde{\mathcal{W}}_{(A_{ev})_{\bar{0}}}^{fin}$), where $\widetilde{\mathcal{W}}_{A_{\bar{0}}}^{fin}$ (respectively, $\widetilde{\mathcal{W}}_{(A_{ev})_{\bar{0}}}^{fin}$) is the category of \mathbb{Z}_2 -graded weight $A_{\bar{0}}$ -modules (respectively, $(A_{ev})_{\bar{0}}$ -modules) with finite weight multiplicities.

Let us prove that the pullback a projective object P in $\widetilde{\mathcal{W}}_A^{fin}$ (respectively, $\widetilde{\mathcal{W}}_{A_{ev}}^{fin}$) is projective in $\mathcal{B}_{\mathfrak{g}}^{osc}$. Since P is induced from a finite-dimensional H_A -module, P is projective in $\widetilde{\mathcal{W}}_{A_{\bar{0}}}^{fin}$ (respectively, $\widetilde{\mathcal{W}}_{(A_{ev})_{\bar{0}}}^{fin}$). By the above equivalence, the pullback of P is projective in $\mathcal{B}_{\mathfrak{g}_0}^{osc}$. Now Lemma 4.2 implies that P is projective in $\mathcal{B}_{\mathfrak{g}}^{osc}$.

Since any object M in $\mathcal{B}_{\mathfrak{g}}^{osc}$ is a quotient of a projective module, M is obtained by pullback from $\widetilde{\mathcal{W}}_A^{fin}$ (respectively, $\widetilde{\mathcal{W}}_{A_{ev}}^{fin}$)-module. The statement follows. \square

Next we recall the following statement.

Proposition 4.4 ([CP], Corollary A.3). *Let $\mathfrak{g} = \varinjlim \mathfrak{g}_k$ be a direct limit of Lie superalgebras. Let $Q = \varinjlim Q_k$ and $R = \varinjlim R_k$ be weight \mathfrak{g} -modules. Assume that R has finite-dimensional weight spaces. Then $\mathrm{Ext}_{\mathfrak{g}_k, \mathfrak{h}_k}^1(Q_k, R_k) = 0$ for all $k \gg 0$ implies $\mathrm{Ext}_{\mathfrak{g}, \mathfrak{h}}^1(Q, R) = 0$.*

We are now ready for

Proof of Theorem 4.1. We only need to consider the case $\mathfrak{g} = \mathfrak{osp}(2a+1|2b)$ or $\mathfrak{g} = \mathfrak{osp}(2a|2b)$ where $\dim \mathfrak{g} = \infty$. We fix an exhaustion $\mathfrak{g} = \varinjlim \mathfrak{g}_k$ for $\mathfrak{g}_k = \mathfrak{osp}(2a_k+1|2b_k)$ or $\mathfrak{g}_k = \mathfrak{osp}(2a_k|2b_k)$, where one of the sequences a_k or b_k may stabilize.

Since our desired equivalence will be obtained simply by pullback via the homomorphisms $\Theta_{a|b}$ or $\Psi_{a|b}$, it suffices to show that every object in $\mathcal{B}_{\mathfrak{g}}^{\text{osc}}$ is the pullback of some weight A -module (respectively, A_{ev} -module). For this, notice that Proposition 4.4 implies that if $P = \varinjlim P_k$ is a direct limit of projective objects in $\mathcal{B}_{\mathfrak{g}_k}^{\text{osc}}$, then P is projective object in $\mathcal{B}_{\mathfrak{g}}^{\text{osc}}$. Next, Theorem 4.1 holds for \mathfrak{g}_k and thus every P_k is the pullback of a projective object in $\widetilde{\mathcal{W}}_{A_k}^{\text{fin}}$ for $\mathfrak{g} = \mathfrak{osp}(2a+1|2b)$ (respectively, $\widetilde{\mathcal{W}}_{(A_k)_{\text{ev}}}^{\text{fin}}$ for $\mathfrak{g} = \mathfrak{osp}(2a|2b)$). Since every object of $\mathcal{B}_{\mathfrak{g}}^{\text{osc}}$ is a quotient of some P as above, we conclude that every object of $\mathcal{B}_{\mathfrak{g}}^{\text{osc}}$ is the pullback of some object of $\widetilde{\mathcal{W}}_A^{\text{fin}}$ (respectively, $\widetilde{\mathcal{W}}_{A_{\text{ev}}}^{\text{fin}}$). \square

Corollary 4.5. *Theorem 4.1 shows that for $b \neq 1$ any indecomposable object M of $\mathcal{B}_{\mathfrak{g}}^{\text{osc}}$ has a filtration similar to the filtration which exists on an indecomposable weight A -module with finite-dimensional weight spaces according to Theorem 2.19.*

Remark 4.6. For $b = 1$ every module in $\mathcal{B}_{\mathfrak{g}}^{\text{osc}}$ has finite length. \circ

The following is our second main result about the category $\mathcal{B}_{\mathfrak{g}}^{\text{osc}}$. Let $(\mathcal{B}_{\mathfrak{g}_0}^{\text{osc}})_{\bar{0}}$ be the category of purely even bounded weight \mathfrak{g}_0 -modules with simple constituents of spinor-oscillator type.

Corollary 4.7. *If $b > 1$ then the category $\mathcal{B}_{\mathfrak{g}}^{\text{osc}}$ is equivalent to the category $(\mathcal{B}_{\mathfrak{g}_0}^{\text{osc}})_{\bar{0}}$. The functor $\mathcal{E} : M \mapsto M_{\bar{0}}$ establishes an equivalence.*

Proof. The statement follows from Theorem 4.1 and from the equivalence of categories established in Theorem 2.21 for $B = A_{\bar{0}}$. \square

Corollary 4.8. *For $b \geq 2$ every non-semisimple block of the category of bounded \mathfrak{g} -modules is equivalent to a block of bounded $D(k|0)$ - or $D(\infty|0)$ -modules with integral weights.*

The category of bounded weight $D(k|0)$ -modules for finite k is described, for example, in [GS]. For the case of $D(\infty|0)$ see [FGM].

5. SIMPLE BOUNDED WEIGHT $\mathfrak{sl}(\infty|\infty)$ -MODULES

We start by two lemmas concerning $\mathfrak{sl}(m|n)$ -modules for $m, n \in \mathbb{Z}_{\geq 0}$. Given a Lie superalgebra $\mathfrak{q} \simeq \mathfrak{sl}(m|n)$ we fix the simple roots of \mathfrak{q} as

$$\varepsilon_1 - \varepsilon_2, \dots, \varepsilon_{m-1} - \varepsilon_m, \varepsilon_m - \delta_1, \delta_1 - \delta_2, \dots, \delta_{n-1} - \delta_n,$$

and let $\omega_1, \dots, \omega_{m-1}, \omega_m, \omega_{m+1}, \dots, \omega_{m+n-1}$ denote the dual basis (fundamental weights). There is an obvious embedding $\mathfrak{sl}(m) \subset \mathfrak{q}_{\bar{0}}$, and we consider $\omega_1, \dots, \omega_{m-1}$ also as fundamental weights of $\mathfrak{sl}(m)$.

Lemma 5.1. *Let $\mathfrak{q} = \mathfrak{sl}(m|n)$ for $m \geq 3$. Let M be a simple bounded highest weight \mathfrak{q} -module with highest weight λ and such that $d(M) < m - 1$. Assume that M is not integrable over the simple ideal $\mathfrak{sl}(m) \subset \mathfrak{q}_0$. Then $\lambda = a\omega_1$ with $a \notin \mathbb{Z}_{\geq 0}$, or $\lambda = -(1+a)\omega_{k-1} + a\omega_k$ for $2 \leq k \leq m$.*

Proof. Denote by μ the weight of $\mathfrak{sl}(m)$ obtained from λ by restriction. By Lemma 1.6, $\text{Ann}_{U(\mathfrak{sl}(m))}L(\mu) = \text{Ann}_{U(\mathfrak{sl}(m))}L(a\omega_1)$ or $\text{Ann}_{U(\mathfrak{sl}(m))}L(\mu) = \text{Ann}_{U(\mathfrak{sl}(m))}L(a\omega_{m-1})$ for some $a \notin \mathbb{Z}_{\geq 0}$. Since the primitive ideals $\text{Ann}_{U(\mathfrak{sl}(m))}L(a\omega_1)$ and $\text{Ann}_{U(\mathfrak{sl}(m))}L(a\omega_{m-1})$ have degree 1, the result of [PSer3] mentioned before Lemma 1.6 shows that also $d(L(\mu)) = 1$. Therefore Proposition 3.4 of [BBL] implies that μ is one of the following weights:

- (1) $a\omega_1$ for $a \notin \mathbb{Z}_{\geq 0}$,
- (2) $b\omega_{m-1}$ for $b \notin \mathbb{Z}_{\geq 0}$,
- (3) $-(1+a)\omega_{k-1} + a\omega_k$ for some $2 \leq k \leq m - 1$ and arbitrary a .

Let us deal first with the cases (1) and (3). Consider the odd reflections with respect to the roots $\varepsilon_m - \delta_1, \dots, \varepsilon_m - \delta_n$ of \mathfrak{q} . Since the restriction to $\mathfrak{sl}(m)$ of the highest weight of M with respect to any reflected Borel subsuperalgebra must satisfy the same respective condition (1) or (3), all these reflections must be atypical. This is only possible if the restriction of λ to the Cartan subalgebra of $\mathfrak{sl}(n)$ equals zero. Furthermore, we have $(\lambda, \varepsilon_m - \delta_1) = 0$. This implies $\lambda = a\omega_1$ or $\lambda = -(1+a)\omega_{k-1} + a\omega_k$ for $2 \leq k \leq m - 1$, respectively.

Now let $\mu = b\omega_{m-1}$ as in (2). After performing all odd reflections with respect to the roots $\varepsilon_m - \delta_1, \dots, \varepsilon_m - \delta_n$, we obtain a highest weight λ' of M such that its restriction to $\mathfrak{sl}(m)$ equals $c\omega_{m-1}$ and $b - c \in \mathbb{Z}_{\geq 0}$. Next, we perform odd reflections with respect to the roots $\varepsilon_{m-1} - \delta_1, \dots, \varepsilon_{m-1} - \delta_n$. By the same argument as in cases (1) and (3), these latter reflections must be atypical. Therefore the restriction of λ' to $\mathfrak{sl}(m)$ equals zero and $(\lambda', \varepsilon_{m-1} - \delta_1) = 0$. In other words, $\lambda' = c\omega_{m-1}$ for some $c \notin \mathbb{Z}_{\geq 0}$. Finally, passing via odd reflections to the original Borel subsuperalgebra yields $\lambda = b\omega_{m-1} + (1-b)\omega_m$. To finish the proof we set $a = 1 - b$. \square

Recall the homomorphisms $\Upsilon_{m|n}^+ : U(\mathfrak{sl}(m|n)) \rightarrow D(m|n)_0$ and $\Upsilon_{m|n}^- : U(\mathfrak{sl}(m|n)) \rightarrow D(n|m)_0$ from Section 2.2. Note that those homomorphisms map the Cartan algebra of $\mathfrak{sl}(m|n)$ to the subalgebra spanned by $u_i - u_j$ for all $i, j \neq 0, i \neq j$. Moreover, the map f induced by $\Upsilon_{m|n}^+$ (respectively, $\Upsilon_{m|n}^-$) from $(\text{Span}\{u_i - u_j \mid i \neq j\})^*$ to \mathfrak{h}^* is linear, and is determined by the correspondence $\zeta_i \mapsto \varepsilon_i$, $\zeta_{-j} \mapsto \delta_j$ (respectively, $\zeta_{-i} \mapsto \varepsilon_i$, $\zeta_j \mapsto \delta_j$).

Corollary 5.2. *Let M be a bounded simple non-integrable $\mathfrak{q} = \mathfrak{sl}(m|n)$ -module with $d(M) < \min(m, n) - 1$. Then $d = 1$ and $\text{Ann}_{U(\mathfrak{q})}M$ contains $\ker \Upsilon_{m|n}^-$ or $\ker \Upsilon_{m|n}^+$.*

Proof. Without loss of generality we can assume that M is not integrable over $\mathfrak{sl}(m)$. Then $\text{Ann}_{U(\mathfrak{q})}M = \text{Ann}_{U(\mathfrak{q})}L(\lambda)$ where λ is one of the weights in Lemma 5.1. It suffices to show that $L(\lambda)$ is obtained by pullback from a weight $D(m|n)_0$ -module.

Consider the $D(m|n)$ -module

$$F(\lambda) := x_k^a \mathbb{C}[x_1^{\pm 1}, \dots, x_m^{\pm 1}, x_{-1}, \dots, x_{-n}],$$

where $k = 1, a \notin \mathbb{Z}_{\geq 0}$ if $\lambda = a\omega_1$, and $a \in \mathbb{C}$ if $\lambda = -(1+a)\omega_{k-1} + a\omega_k$ for $2 \leq k \leq m$.

Let $f_k := x_1^{-1} \dots x_{k-1}^{-1} x_k^a$ and let Y_k denote the $D(m|n)_0$ -submodule in $F(\lambda)$ generated by f_k . Note that the weight of f_k equals $a\omega_1$ for $k = 1$, and equals $-(1+a)\omega_{k-1} + a\omega_k$ for $2 \leq k \leq m$. Then Y_1 is a simple $D(m|n)_0$ -module and its pullback along $\Upsilon_{m|n}^+$ is isomorphic to $L(a\omega_1)$, since by direct computation one can see that any vector annihilated by all $x_{i-1}\partial_i$ for $2 \leq i \leq m$ is proportional to f_1 . For $k > 1$, consider the $D(m|n)_0$ -submodule $Z_k \subset Y_k$ generated by $x_{i-1}\partial_i(f)$ for $2 \leq i \leq m$. Then $f \notin Z_k$ and hence $X_k := Y_k/Z_k \neq 0$. Furthermore, the pullback along $\Upsilon_{m|n}^+$ of X_k is isomorphic to $L(-(1+a)\omega_{k-1} + a\omega_k)$ (again because any vector annihilated by all $x_{i-1}\partial_i$ is proportional to f_k). The statement is proved. \square

In the rest of this section $\mathfrak{g} = \mathfrak{sl}(\infty|\infty)$ and $A = D(\infty|\infty)$. We fix an exhaustion $\mathfrak{g} = \varinjlim \mathfrak{g}_k$, where $\mathfrak{g}_k \simeq \mathfrak{sl}(k|k)$. By \mathfrak{g}^\pm we denote the ideals of \mathfrak{g}_0 with respective roots $\varepsilon_i - \varepsilon_j$ and $\delta_i - \delta_j$, and we write Υ^\pm instead of $\Upsilon_{\infty|\infty}^\pm$.

Lemma 5.3. *Let M be a simple bounded weight \mathfrak{g} -module not integrable over \mathfrak{g}^+ or \mathfrak{g}^- . Then $\text{Ann}_{U(\mathfrak{g})} M$ contains $\ker \Upsilon^+$, or respectively $\ker \Upsilon^-$, and therefore M is multiplicity free.*

Proof. Let $v \in M$ be a nonzero weight vector and let $M_k := U(\mathfrak{g}_k)v$. If $k > d(M)$ then Corollary 5.2 implies that $\text{Ann}_{U(\mathfrak{g}_k)} M_k$ contains $\ker \Upsilon_{k|k}^\pm$. Therefore $\text{Ann}_{U(\mathfrak{g})} M = \varinjlim \text{Ann}_{U(\mathfrak{g}_k)} M_k$ contains $\ker \Upsilon^\pm = \varinjlim \ker \Upsilon_k^\pm$. Since every simple weight A_0 -module is multiplicity free, the second assertion follows. \square

Remark 5.4. One may observe that Lemma 5.3 holds also for the Lie superalgebra $\mathfrak{sl}(\infty|n)$, $n \in \mathbb{Z}_{>0}$, where one replaces \mathfrak{g}^+ by the simple ideal $\mathfrak{sl}(\infty)$ of $\mathfrak{sl}(\infty|n)_0$ and Υ^+ by $\Upsilon_{\infty|n}^+$. \circ

The simple bounded integrable \mathfrak{g} -modules have been classified in [CP], Theorem 5.9. Therefore, in order to classify all simple bounded weight \mathfrak{g} -modules it suffices to prove the following.

Theorem 5.5. *Let M be a simple bounded non-integrable \mathfrak{g} -module. Then*

- (a) M is multiplicity free.
- (b) M is obtained from a simple weight A_0 -module by pullback via precisely one of the homomorphisms Υ^+ or Υ^- , and accordingly either \mathfrak{g}^- or \mathfrak{g}^+ acts integrably on M .
- (c) Pullback via Υ^\pm establishes a bijection between isomorphism classes of simple, non-integrable over \mathfrak{g}^\pm , bounded \mathfrak{g} -modules and isomorphism classes of simple non-integrable weight A_0 -modules.

Proof. (a) follows directly from Lemma 5.3.

(b) Let C^\pm denote the image of Υ^\pm . It is easy to see that $C^\pm \subsetneq A_0$. Lemma 5.3 implies that every simple non-integrable weight \mathfrak{g} -module is obtained by pullback from a simple C^+ - or a C^- -module. Therefore, to prove (2) we need to show that a weight \mathfrak{g} -module obtained by pullback from a weight C^\pm -module is in fact obtained by pullback from the restriction of a weight A_0 -module to C^\pm . It suffices to prove the statement for C^+ , since the other case follows by applying the obvious automorphism of \mathfrak{g} .

Recall the basis $\{u_i\}_{i \in \mathbb{Z}}$ of \mathfrak{h}_A introduced in Section 2.4. By a slight abuse of notation we denote by the same letter the preimage of u_i in the Cartan subalgebra of $\mathfrak{gl}(\infty|\infty)$. Then $\{w_i = u_i - u_{-1} \mid i \neq -1\}$ is a basis of the Cartan subalgebra of \mathfrak{g} . Let N be a simple weight \mathfrak{g} -module, $\mu \in \text{supp}N$ and $c \in \mathbb{C}$. Note that we can endow N with a $\mathfrak{gl}(\infty|\infty)$ -module structure by setting $u_{-1}v := (c + \nu(u_{-1}) - \mu(u_{-1}))v$ for every $v \in N^\nu$. We denote this $\mathfrak{gl}(\infty|\infty)$ -module by $N(\mu, c)$.

We claim that if M is the pullback of some simple weight C^+ -module, then we can find μ, c such that the $\mathfrak{gl}(\infty|\infty)$ -module $N(\mu, c)$ is the pullback of some weight A_0 -module. Clearly, we can assume that M is not trivial. We pick some $\kappa \in \text{supp}M$ such that $\kappa(w_i) \neq 0$ for some $i \leq -2$. One readily sees that the relation $w_i^3 = w_i$ implies $\kappa(w_i) = \pm 1, 0$ for $i \leq -2$. Next, we choose a negative i such that $\kappa(w_i) \neq 0$. It easily follows from the linearity of κ that $\kappa(w_j) = 0$ or $\kappa(w_j) = \kappa(w_i)$ for every negative j . Finally, we set $c = 0$ if $\kappa(w_i) = 1$ and $c = 1$ if $\kappa(w_i) = -1$. Then $\text{supp}M(\kappa, c) \subset \mathfrak{h}_A^\vee$ and, since the restriction of $M(\kappa, c)$ to \mathfrak{g} is the pullback of some weight C^+ -module, the $\mathfrak{gl}(\infty|\infty)$ -module $M(\kappa, c)$ is the pullback of a weight A_0 -module.

(c) Follows from Proposition 2.23(b). \square

Remark 5.6. It is likely that Theorem 5.5 holds also for $\mathfrak{sl}(\infty|n)$. \circ

Remark 5.7. Note that the definition of \mathfrak{h}_A^\vee implies that if M is the pullback of a weight A_0 -module via Υ^+ (respectively, Υ^-), then for $\sum_i a_i \varepsilon_i + \sum_j b_j \delta_j \in \text{supp}M$ we have $a_i \in \{0, 1\}$ (respectively, $b_j \in \{0, 1\}$). \circ

Proposition 5.8. *A simple bounded weight \mathfrak{g} -module M is determined, up to isomorphism and a possible parity change, by $\text{supp}M$.*

Proof. Here we consider the case of non-integrable modules, and leave as an exercise to the reader to check our claim for integrable modules using the classification result of [CP]. Let us observe that if M and N are not integrable, and one is obtained by pullback via Υ^+ while the other is obtained by pullback via Υ^- , then M and N cannot have the same support.

Now, without loss of generality we can assume that M and N are obtained by pullback from simple weight A_0 -modules X and Y , respectively. Suppose that $\text{supp}M = \text{supp}N$ but $\text{supp}X \neq \text{supp}Y$. Then $\text{supp}X = \text{supp}Y \pm \tau$ where $\tau = \sum_{i>0} (\varepsilon_i - \delta_i)$. Since the supports of X and Y are subsets of \mathfrak{h}_A^\vee , this is only possible if $\text{supp}X = \{0\}$, $\text{supp}Y = \{\pm\tau\}$ or vice versa. Then, both M and N are necessarily trivial and we have

a contradiction. Consequently, $\text{supp}M = \text{supp}N$ implies $\text{supp}X = \text{supp}Y$, and then the A_0 -modules X and Y are isomorphic up to parity change by Theorem 2.22(a). This completes the proof. \square

Let $M^\pm(\mu)$ denote the simple weight \mathfrak{g} -module obtained by pullback from the simple weight A_0 -module $Y(\mu)$ via Υ^\pm .

Proposition 5.9. *Every multiplicity free simple weight \mathfrak{g} -module M is isomorphic to the pullback of a simple weight A_0 -module via Υ^+ or Υ^- . If M is obtained by pullback via both Υ^+ and Υ^- , then M is isomorphic to $V, \Pi V, V_*, \Pi V_*, \mathbb{C}$ or $\Pi\mathbb{C}$.*

Proof. By Theorem 5.5 all non-integrable simple bounded \mathfrak{g} -modules are pullbacks of A_0 -modules via Υ^+ or Υ^- , and hence are multiplicity free. Therefore it suffices to check the statement for integrable multiplicity free modules.

Theorem 5.9 in [CP] implies that, in addition to the six modules $V, \Pi V, V_*, \Pi V_*, \mathbb{C}, \Pi\mathbb{C}$ there are four families of multiplicity free simple integrable \mathfrak{g} -modules $S_{\mathcal{A}}^\infty V, S_{\mathcal{A}}^\infty V_*, \Lambda_{\mathcal{A}}^\infty V, \Lambda_{\mathcal{A}}^\infty V_*$. If one observes that all other three families are obtained from $S_{\mathcal{A}}^\infty V$ by a twist from a proper automorphism of $\mathfrak{sl}(\infty|\infty)$, it remains to check that any simple module of the form $S_{\mathcal{A}}^\infty V$ is isomorphic to $M^-(\mu_{\mathcal{A}})$ or $\Pi M^-(\mu_{\mathcal{A}})$ for a weight $\mu_{\mathcal{A}} \in \mathfrak{h}_{\mathcal{A}}^\vee$.

Recall from [CP] that \mathcal{A} is a sequence of pairs (a_n, b_n) where $a_1 \leq a_2 \leq \dots$ is a sequence of positive integers and $b_n \in \{0, 1\}$ with the condition that $b_n = b_{n+1}$ if $a_n = a_{n+1}$. Moreover, $S_{\mathcal{A}}^\infty V$ is defined as the direct limit $\varinjlim \Pi^{b_n} S^{a_n} V_n$ where V_n is the natural $\mathfrak{sl}(n|n)$ -module. Let

$$\mu_{\mathcal{A}} := \sum_{i>0} (b_i \varepsilon_i + (a_i - a_{i-1} - b_i) \delta_i),$$

where we set $a_0 = 0$. Then a direct verification shows that $\text{supp}S_{\mathcal{A}}^\infty V = \text{supp}M^-(\mu_{\mathcal{A}})$. Since a simple multiplicity free weight \mathfrak{g} -module is determined by its support up to isomorphism and a possible application by Π , we conclude that $S_{\mathcal{A}}^\infty V \simeq M^-(\mu_{\mathcal{A}})$ if the weight space $(S_{\mathcal{A}}^\infty V)^{\mu_{\mathcal{A}}}$ has parity equal to $p(\mu_{\mathcal{A}})$, and $S_{\mathcal{A}}^\infty V \simeq \Pi M^-(\mu_{\mathcal{A}})$ otherwise. In fact, the parity of the weight space $(S_{\mathcal{A}}^\infty V)^{\mu_{\mathcal{A}}}$ depends only on b_1 : the weight space $(S_{\mathcal{A}}^\infty V)^{\mu_{\mathcal{A}}}$ is purely even for $b_1 = 0$ and purely odd for $b_1 = 1$.

Finally, the fact that each of the six modules $V, \Pi V, V_*, \Pi V_*, \mathbb{C}, \Pi\mathbb{C}$ is obtained by pullback via both Υ^+ and Υ^- is straightforward. \square

Proposition 5.10. *If M is a simple bounded weight \mathfrak{g} -module then M is semisimple as a \mathfrak{g}_0 -module.*

Proof. The statement is clear for integrable modules since every bounded integrable \mathfrak{g}_0 -module is semisimple by Theorem 3.7 in [PSer2]. Therefore, without loss of generality we can assume that M is isomorphic to $M^\pm(\mu)$. Consider the lattice $Q_{(A_0)_0}$

with generators $\varepsilon_i - \varepsilon_j, \delta_i - \delta_j$. Set

$$Y(\mu)^n := \bigoplus_{\nu \in \mu + n(\varepsilon_1 - \delta_1) + Q_{(A_0)_{\bar{0}}}} Y(\mu)^\nu.$$

Then $Y(\mu)^n$ is a simple $(A_0)_{\bar{0}}$ -module and $Y(\mu) = \bigoplus_{n \in \mathbb{Z}} Y(\mu)^n$. Obviously, the semisimplicity of $Y(\nu)$ over $(A_0)_{\bar{0}}$ implies semisimplicity of M over $\mathfrak{g}_{\bar{0}}$. \square

Theorem 5.11. (a) Let $\mathfrak{g} = \mathfrak{sl}(\infty)$. The following ideals are all bounded primitive ideals of $U(\mathfrak{g})$:

$$\text{Ann}_{U(\mathfrak{g})} \mathbb{S}_\lambda V, \text{Ann}_{U(\mathfrak{g})} \mathbb{S}_\lambda V_*, \ker \Upsilon^+, \ker \Upsilon^-.$$

(b) Let $\mathfrak{g} = \mathfrak{osp}(2a+1|2b)$ (respectively, $\mathfrak{g} = \mathfrak{osp}(2a|2b)$) with at least one of a and b equal infinity. Then $U(\mathfrak{g})$ has exactly three bounded primitive ideals: the augmentation ideal, $\text{Ann}_{U(\mathfrak{g})} V$, and $\ker \Theta_{a|b}$ (respectively, $\ker \Psi_{a|b}$).

Proof. (a) follows from Corollary 2.25, Theorem 5.5, Proposition 5.8, and the classification of simple bounded integrable \mathfrak{g} -modules in [CP]. (b) follows from Corollary 3.6. \square

Lemma 5.12. Let \mathcal{B} denote the category of bounded weight \mathfrak{g} -modules, and let $M, N \in \mathcal{B}$. Denote by $Q_{\mathfrak{g}}$ the root lattice of \mathfrak{g} .

- (a) $\text{Ext}_{\mathcal{B}}^1(M, N) = \text{Ext}_{\mathcal{B}}^1(N, M)$.
- (b) If M is simple and $\text{Ext}_{\mathcal{B}}^1(M, N) \neq 0$, then $\text{supp} M \subset \text{supp} N + Q_{\mathfrak{g}}$.
- (c) $\text{Ext}_{\mathcal{B}}^1(M^\pm(\mu), \Pi M^\pm(\nu)) = 0$ for all μ, ν .
- (d) If $\text{Ext}_{\mathcal{B}}^1(M^+(\mu), M^+(\nu)) \neq 0$, then $\mu - \nu \in Q_{A_0}$ or at least one of $M^+(\mu)$ and $M^-(\nu)$ is trivial.
- (e) If $\text{Ext}_{\mathcal{B}}^1(M^+(\mu), M^-(\nu)) \neq 0$, then at least one of $M^+(\mu)$ and $M^-(\nu)$ is isomorphic to V, V_*, \mathbb{C} .
- (f) If M and N are simple and $d(M) > 1$ then $\text{Ext}_{\mathcal{B}}^1(M, N) = 0$.

Proof. (a) We consider the (contravariant) functor of contragredient duality \cdot^\vee on the category \mathcal{B} . Then $M^\vee \simeq M, N^\vee \simeq N$ and

$$\text{Ext}_{\mathcal{B}}^1(M, N) = \text{Ext}_{\mathcal{B}}^1(N^\vee, M^\vee) = \text{Ext}_{\mathcal{B}}^1(N, M).$$

(b) Let $0 \rightarrow M \rightarrow R \rightarrow N \rightarrow 0$ represent a nonzero element of $\text{Ext}_{\mathcal{B}}^1(M, N)$. Then, for some weight vector $v \in N$, the image of M in R is a submodule of $U(\mathfrak{g})v'$ where v' is a preimage of v in R of weight κ . Then $\text{supp} M \subset \kappa + Q_{\mathfrak{g}} \subset \text{supp} N + Q_{\mathfrak{g}}$.

(c) follows from comparing the parity of weight spaces of the modules $M^\pm(\mu)$ with the parity of the weight spaces of the modules $\Pi M^\pm(\nu)$.

(d) follows from (b).

(e) For a \mathfrak{g} -module M and a Lie subsuperalgebra \mathfrak{k} of \mathfrak{g} we denote by $\Gamma_{\mathfrak{k}} M$ the set of locally finite \mathfrak{k} -vectors, i.e.,

$$\Gamma_{\mathfrak{k}} M := \{m \in M \mid \dim \text{span}\{m, km, k^2 m, \dots\} < \infty \forall k \in \mathfrak{k}\}.$$

The superspace $\Gamma_{\mathfrak{k}}M$ is a \mathfrak{g} -submodule of M . This is established for Lie algebras in particular in Theorem 8.2 in [PH], and the proof for Lie superalgebras is the same.

By the semisimplicity result in [CP], at least one of $M^+(\mu)$ and $M^-(\nu)$ can be assumed non-integrable. Moreover, by (a), the statement is symmetric with respect to $M^+(\mu)$ and $M^-(\nu)$. Without loss of generality, assume that $M^-(\nu)$ is not integrable. Consider a non-split exact sequence

$$0 \rightarrow M^-(\nu) \rightarrow N \rightarrow M^+(\mu) \rightarrow 0.$$

Since $\Gamma_{\mathfrak{g}^+}M^-(\nu) = M^-(\nu)$, there exists a root α of \mathfrak{g}^- such that \mathfrak{g}_α acts freely on $M^-(\nu)$. If $\Gamma_{\mathfrak{g}_\alpha}N \neq 0$ then $\Gamma_{\mathfrak{g}_\alpha}N$ is a submodule of N which does not coincide with $M^-(\nu)$, i.e., the sequence splits. Consequently, $\Gamma_{\mathfrak{g}_\alpha}N = 0$. Hence, for any $\theta \in \text{supp}M^+(\mu)$ we have $\theta + n\alpha \in \text{supp}M^-(\nu)$ for $n \geq 2$. Thus, we get that $\theta_i \in \{0, 1\}$ for all $i > 0$, and therefore for all i by Remark 5.7. This is possible if and only if $M^-(\nu)$ is isomorphic to V , V_* or \mathbb{C} .

(f) If $d(M) > 1$ then using Theorem 5.9 in [CP] and Proposition 1.2 one can verify that M is isomorphic to $\mathbb{S}_\lambda V$, $\text{II}\mathbb{S}_\lambda V$, $\mathbb{S}_\lambda V_*$, or $\text{II}\mathbb{S}_\lambda V_*$ for some Young diagram λ with more than one row or more than one column. Assume $M = \mathbb{S}_\lambda V$ and $\text{Ext}^1(M, N) \neq 0$. Then the semisimplicity result (Theorem 6.1) in [CP] implies that N is not integrable. Consider a non-split exact sequence

$$0 \rightarrow N \rightarrow R \rightarrow M \rightarrow 0.$$

Suppose $N \simeq M^-(\nu)$ for some ν . The argument in the proof of (e) can be easily modified to show that $(\theta, \alpha^\vee) \in \{\pm 1, 0\}$ for any weight θ of M and any root α of \mathfrak{g}^+ . This implies that λ consists of a single column, and hence $d(M) = 1$. Similarly, if $N \simeq M^+(\nu)$ one proves that λ consists of a single row and $d(M) = 1$. \square

REFERENCES

- BBL. G. Benkart, D. Britten, F. Lemire, Modules with bounded weight multiplicities for simple Lie algebras, *Math. Z.* **225** (1997), 333–353.
- BKLM. D. Britten, O. Khomenko, F. Lemire, V. Mazorchuk, Complete reducibility of torsion free C_n -modules of finite degree, *J. Algebra* **276** (2004), 129–142.
- C. L. Calixto, On the structure of simple bounded weight modules of $\mathfrak{sl}(\infty)$, $\mathfrak{o}(\infty)$, $\mathfrak{sp}(\infty)$, *Comm. Alg.* **48** (2020), 4256–4280.
- CP. L. Calixto, I. Penkov, Integrable bounded weight modules of classical Lie superalgebras at infinity, *J. Pure Appl. Algebra*, DOI: 10.1016/j.jpaa.2021.106847
- DPS. E. Dan-Cohen, I. Penkov, V. Serganova, A Koszul category of representations of finitary Lie algebras, *Adv. Math.* **289** (2016), 250–278.
- DP. I. Dimitrov, I. Penkov, Weight modules of direct limit Lie algebras, *IMRN* **5** (1999), 223–249.
- Fer. S. Fernando, Lie algebra modules with finite-dimensional weight spaces I, *Trans. Amer. Math. Soc.* **322** (1990), 757–781.
- FG. R. Fiorese, F. Gavarini, Chevalley Supergroups, *Memoirs of the American Mathematical Society* **215** (2012), no. 1014.
- Fuks. D. B. Fuks, Cohomology of infinite-dimensional Lie algebras, *Contemporary Soviet Mathematics*, Consultants Bureau, New York, 1986.

- FGM. V. Futorny, D. Grantcharov, V. Mazorchuk, Weight modules over infinite dimensional Weyl algebras, *Proc. Amer. Math. Soc.* **142** (2014), 3049–3057.
- GP. D. Grantcharov, I. Penkov. Simple bounded weight modules of \mathfrak{sl}_∞ , \mathfrak{o}_∞ , \mathfrak{sp}_∞ , *Transform. Groups.* **25** (2020), 1125–1160.
- GS. D. Grantcharov, V. Serganova, Category of $\mathfrak{sp}(2n)$ -modules with bounded weight multiplicities, *Mosc. Math. J.* **6** (2006), 119–134.
- HS. J. Hartwig, V. Serganova, Clifford and Weyl superalgebras and spinor representations, *Transform. Groups.* **25** (2020), 1185–1207.
- Mar. E. Marczewski, Sur l’extension de l’ordre partiel, *Fund. Math. (in French)* **16** (1930), 386–389.
- Mat. O. Mathieu, Classification of irreducible weight modules, *Ann. Inst. Fourier* **50** (2000), 537–592.
- M. I. Musson, Lie superalgebras and enveloping algebras, Graduate Studies in Mathematics **131**, American Mathematical Society, 2012.
- PH. I. Penkov, C. Hoyt, Classical Lie algebras at infinity, Springer Monographs in Mathematics, Springer Verlag 2022.
- PSer1. I. Penkov, V. Serganova, Tensor representations of Mackey Lie algebras and their dense subalgebras, In: Developments and Retrospectives in Lie Theory: Algebraic Methods, Developments in Mathematics, **38**, Springer Verlag 2014, 291–330.
- PSer2. I. Penkov, V. Serganova, Categories of integrable $sl(\infty)$ -, $o(\infty)$ -, $sp(\infty)$ -modules. In: Representation Theory and Mathematical Physics, *Cont. Math.* **557** (2011), 335–357.
- PSer3. I. Penkov, V. Serganova, On bounded generalized Harish-Chandra modules, *Annales de l’Institut Fourier* **62** (2012), 477–496.
- PStyr. I. Penkov, K. Styrkas, Tensor representations of infinite-dimensional root-reductive Lie algebras, in Developments and Trends in Infinite-Dimensional Lie Theory, Progress in Mathematics 288, Birkhäuser, 2011, 127–150.
- SS. S. Sam, A. Snowden, Stability patterns in representation theory, *Forum Math. Sigma* **3** (2015), e11,108 pp.
- S. V. Serganova, Classical Lie superalgebras at infinity. In: Advances in Lie Superalgebras. Springer INdAM Series 7, Springer Verlag, 2014, 181–201.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT ARLINGTON, ARLINGTON, TX
76019-0408, USA

E-mail address: grandim@uta.edu

JACOBS UNIVERSITY BREMEN, CAMPUS RING 1, 28759 BREMEN, GERMANY

E-mail address: i.penkov@jacobs-university.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT BERKELEY, BERKELEY,
CA 94720, USA

E-mail address: serganov@math.berkeley.edu