

Oberwolfach Preprints



OWP 2014 - 05

NHAN-PHU CHUNG AND GUOHUA ZHANG

Weak Expansiveness for Actions of Sofic Groups

Mathematisches Forschungsinstitut Oberwolfach gGmbH
Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

Starting in 2007, the MFO publishes a preprint series which mainly contains research results related to a longer stay in Oberwolfach. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 30 free hard copies (DIN A4, black and white copy) by surface mail.

Of course, the full copy right is left to the authors. The MFO only needs the right to publish it on its website *www.mfo.de* as a documentation of the research work done at the MFO, which you are accepting by sending us your file.

In case of interest, please send a **pdf file** of your preprint by email to *rip@mfo.de* or *owlf@mfo.de*, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany

Tel +49 7834 979 50
Fax +49 7834 979 55
Email admin@mfo.de
URL www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.
Copyright of the content is held by the authors.

WEAK EXPANSIVENESS FOR ACTIONS OF SOFIC GROUPS

NHAN-PHU CHUNG AND GUOHUA ZHANG

ABSTRACT. In this paper, we shall introduce h -expansiveness and asymptotical h -expansiveness for actions of sofic groups. By the definitions, each h -expansive action of sofic groups is asymptotically h -expansive. We show that each expansive action of sofic groups is h -expansive, and, for any given asymptotically h -expansive action of sofic groups, the entropy function (with respect to measures) is upper semi-continuous and hence the system admits a measure with maximal entropy.

Observe that asymptotically h -expansive property was firstly introduced and studied by Misiurewicz for \mathbb{Z} -actions using the language of tail entropy. And thus in the remaining part of the paper, we shall compare our definitions of weak expansiveness for actions of sofic groups with the definitions given in the same spirit of Misiurewicz's ideas when the group is amenable. It turns out that these two definitions are equivalent in this setting.

1. INTRODUCTION

Dynamical system theory is the study of qualitative properties of group actions on spaces with certain structures. In this paper, by a *topological dynamical system* we mean a continuous action of a countable discrete sofic group on a compact metric space. Sofic groups were defined implicitly by Gromov in [Gro99a] and explicitly by Weiss in [Wei00], which include all amenable groups and residually finite groups.

Recently, Lewis Bowen introduced a notion of entropy for measure-preserving actions of a countable discrete sofic group admitting a generating measurable partition with finite entropy [Bow10, Bow12]. The main idea here is to replace the important Følner sequence of a countable discrete amenable group with a sofic approximation for a countable discrete sofic group. Very soon after [Bow10], in the spirit of L. Bowen's sofic measure-theoretic entropy, Kerr and Li developed an operator-algebraic approach to sofic entropy [KL11, KL13b] which applies not only to continuous actions of countable discrete sofic groups on compact metric spaces but also to all measure-preserving actions of countable discrete sofic groups on standard probability measure spaces. From then on, there are many other papers,

Date: March 16, 2014.

2010 Mathematics Subject Classification. Primary 37B05; Secondary 37C85, 54H15.

Key words and phrases. actions of sofic groups, expansive, h -expansive, asymptotically h -expansive, measures with maximal entropy, actions of amenable groups.

presenting different but equivalent definitions of sofic entropy [Ker13, Zha12], extending sofic entropy to sofic pressure [Chu13] and to sofic mean dimension [Li13], and discussing combinatorial independence for actions of sofic groups [KL13a].

Let X be a compact metric space. Any homeomorphism $T : X \rightarrow X$ generates naturally a topological dynamical system by considering the group $\{T^n : n \in \mathbb{Z}\}$. Even the given map $T : X \rightarrow X$ is just a continuous map (may be non-invertible), we still call it a topological dynamical system by considering the semi-group $\{T^n : n \in \mathbb{Z}_+\}$. A self-homeomorphism of a compact metric space is *expansive* if, for each pair of distinct points, some iterate of the homeomorphism separates them by a definite amount. Expansiveness is in fact a multifaceted dynamical condition which plays a very important role in the exploitation of hyperbolicity in smooth dynamical system [Mañ87]. In the setting of considering a continuous mapping over a compact metric space, two classes of weak expansiveness, the h -expansiveness and asymptotical h -expansiveness, were introduced by Rufus Bowen [Bow72] and Misiurewicz [Mis76], respectively. By the definitions, each h -expansive system is asymptotically h -expansive. Both of h -expansiveness and asymptotical h -expansiveness turn out to be important in the study of smooth dynamical system [Bur11, DFPV12, DM09, DN05, LVY].

It is direct to define expansiveness for actions of groups. That is, let G be a discrete group acting on a compact metric space X (with the metric ρ), then we say that (X, G) is *expansive* if there is $\delta > 0$ such that for any two different points x_1 and x_2 in X there exists $g \in G$ with $\rho(gx_1, gx_2) > \delta$. Such a δ is called an *expansive constant*. Symbolic systems are standard examples for expansive actions. This obvious extension of the notion of expansiveness has been investigated extensively in algebraic actions for \mathbb{Z}^d [ES02, LS99, Sch90, Sch95] and for more general groups [Bow11, CL, DS07]; and a general framework of dynamics of $d \in \mathbb{N}$ commuting homeomorphisms over a compact metric space, in terms of expansive behavior along lower dimensional subspaces of \mathbb{R}^d , was firstly proposed by Boyle and Lind [BL97].

Now, a natural question rises: how to define weak expansiveness using entropy techniques when considering actions of countable sofic groups?

The problem is addressed in this paper. When considering a continuous mapping over a compact metric space, R. Bowen introduced h -expansiveness using separated and spanning subsets by considering topological entropy of special subsets [Bow72], and then Misiurewicz introduced asymptotical h -expansiveness, weaker than h -expansiveness, using open covers by introducing tail entropy [Mis76]. Observe that the notion of tail entropy was first introduced by Misiurewicz in [Mis76] where he called it topological conditional entropy, and then Buzzi called this quantity local entropy in [Buz97]. Here, we follow Downarowicz and Serafin [DS02] and Downarowicz [Dow11]. It was Li who used firstly open covers for actions of

sofic groups to consider sofic mean dimension [Li13], and then this idea was used in [Zha12] to consider equivalently the entropy for actions of sofic groups. To fix the problem of defining weak expansiveness naturally for actions of sofic groups, we shall use open covers again to introduce the properties of h -expansiveness and asymptotical h -expansiveness in the spirit of Misiurewicz [Mis76]. The idea turns out to be successful. From the definitions, each h -expansive action of sofic groups is asymptotically h -expansive; we shall prove that each expansive action of sofic groups is h -expansive (Theorem 3.1), and hence, h -expansiveness and asymptotical h -expansiveness are indeed two classes of weak expansiveness. Additionally, similar to the setting of considering a continuous mapping over a compact metric space, for any given asymptotically h -expansive action of sofic groups, the entropy function (with respect to measures) is upper semi-continuous (Theorem 3.5) and hence the system admits really a measure with maximal entropy.

Observe that the asymptotically h -expansive property was first introduced and studied by Misiurewicz for \mathbb{Z} -actions using the language of tail entropy. We can define tail entropy for actions of amenable groups in the same spirit, and so it is quite natural to ask if we could define asymptotical h -expansiveness for actions of amenable groups along the line of tail entropy. The answer turns out to be true, that is, our definitions of weak expansiveness for actions of sofic groups are equivalent to the definitions given in the same spirit of Misiurewicz's ideas of using tail entropy when the group is amenable (Theorem 6.1). In [Mis76] Misiurewicz provided a typical example of an asymptotically h -expansive system, that is, any continuous endomorphism of a compact metric group with finite entropy is asymptotically h -expansive. We shall show that in fact this holds in a more general setting with the help of Theorem 6.1, precisely, any action of a countable discrete amenable group acting on a compact metric group by continuous automorphisms is asymptotically h -expansive if the action has finite entropy (Theorem 7.1).

The paper is organized as follows. In section 2 we prove that each expansive action of sofic groups admits a measure with maximal entropy based on the sofic measure-theoretic entropy introduced in [Bow10]. In section 3 we introduce h -expansive and asymptotically h -expansive actions of sofic groups in the spirit of Misiurewicz [Mis76]. Each h -expansive action of sofic groups is asymptotically h -expansive by the definitions. We show that each expansive action of sofic groups is h -expansive, and each asymptotically h -expansive action of sofic groups admits a measure with maximal entropy (in fact, its entropy function is upper semi-continuous with respect to measures). In section 4 we present our first interesting non-trivial example of an h -expansive action of sofic groups which is in fact the profinite action of a countable group. In order to understand further our introduced weak expansiveness for actions of sofic groups in the setting of amenable groups, in section 5 we define tail entropy

for actions of amenable groups in the same spirit of Misiurewicz. And then in section 6 we compare our definitions of weak expansiveness for actions of sofic groups with the definitions given in section 5 when the group is amenable. It turns out that these two definitions are equivalent in this setting. And then in section 7 we show that any action of a countable discrete amenable group acting on a compact metric group by continuous automorphisms is asymptotically h -expansive if the action has finite entropy.

2. EXPANSIVE ACTIONS OF SOFIC GROUPS

Let G be a countable discrete group. For each $d \in \mathbb{N}$, denote by $\text{Sym}(d)$ the permutation group of $\{1, \dots, d\}$; and then we say that G is *sofic* if there is a sequence $\Sigma = \{\sigma_i : G \rightarrow \text{Sym}(d_i), g \mapsto \sigma_{i,g}, d_i \in \mathbb{N}\}_{i \in \mathbb{N}}$ such that

$$\lim_{i \rightarrow \infty} \frac{1}{d_i} |\{a \in \{1, \dots, d_i\} : \sigma_{i,s} \sigma_{i,t}(a) = \sigma_{i,st}(a)\}| = 1 \text{ for all } s, t \in G$$

and

$$\lim_{i \rightarrow \infty} \frac{1}{d_i} |\{a \in \{1, \dots, d_i\} : \sigma_{i,s}(a) \neq \sigma_{i,t}(a)\}| = 1 \text{ for all distinct } s, t \in G.$$

Here, by $|\bullet|$ we mean the cardinality of a set \bullet . Such a sequence Σ with $\lim_{i \rightarrow \infty} d_i = \infty$ is referred as a *sofic approximation of G* . Observe that the condition $\lim_{i \rightarrow \infty} d_i = \infty$ is essential for the variational principle concerning entropy of actions of sofic groups (see [KL11] and [Zha12] for the global and local variational principles, respectively), and it is automatic if G is infinite.

Throughout the paper, G will be a countable discrete sofic group, with a fixed sofic approximation Σ as above and act on a compact metric space (X, ρ) .

In this section, based on sofic measure-theoretic entropy introduced in [Bow10], we mainly prove that, for an expansive action of sofic groups, the entropy function is upper semi-continuous with respect to measures, and hence the action admits a measure with maximal entropy. Additionally, we show that in general the entropy function of a finite open cover is also upper semi-continuous with respect to measures.

Denote by $M(X)$ the set of all Borel probability measures on X , which is a compact metric space if endowed with the well-known weak star topology; and by $M(X, G)$ the set of all G -invariant elements μ in $M(X)$, i.e., $\mu(A) = \mu(g^{-1}A)$ for each $g \in G$ and all $A \in \mathcal{B}_X$, where \mathcal{B}_X is the Borel σ -algebra of X . Note that if $M(X, G) \neq \emptyset$ then it is a compact metric space.

For a set Y , we denote by \mathfrak{F}_Y the set of all non-empty finite subsets of Y . By a *cover* of X we mean a family of subsets of X with the whole space as its union. If elements of a cover are pairwise disjoint, then it is called a *partition*. Denote by $\mathcal{C}_X, \mathcal{C}_X^o, \mathcal{C}_X^c$ and \mathcal{P}_X the set of all finite Borel covers, finite open covers, finite closed

covers and finite Borel partitions of X , respectively. For $\mathcal{V} \in \mathcal{C}_X$ and $\emptyset \neq K \subset X$ we set $\overline{\mathcal{V}} = \{\overline{V} : V \in \mathcal{V}\}$, and set $N(\mathcal{V}, K)$ to be the minimal cardinality of sub-families of \mathcal{V} covering K (with $N(\mathcal{V}, \emptyset) = 0$ by convention).

Throughout the whole paper, we shall fix the convention $\log 0 = -\infty$.

Now we recall the sofic measure-theoretic entropy introduced in [Bow10, §2].

Let $\alpha = \{A_1, \dots, A_k\} \in \mathcal{P}_X$, $k \in \mathbb{N}$ and $\sigma : G \rightarrow \text{Sym}(d)$, $d \in \mathbb{N}$, and let ζ be the uniform probability measure on $\{1, \dots, d\}$ and $\beta = \{B_1, \dots, B_k\}$ a partition of $\{1, \dots, d\}$. Assume $\mu \in M(X, G)$. For $F \in \mathfrak{F}_G$, we denote by $\text{Map}(F, k)$ the set of all functions $\phi : F \rightarrow \mathbb{N}$ such that $\phi(f) \leq k$ for all $f \in F$, and we set

$$d_F(\alpha, \beta) = \sum_{\phi \in \text{Map}(F, k)} |\mu(A_\phi) - \zeta(B_\phi)|,$$

where

$$A_\phi = \bigcap_{f \in F} f^{-1}A_{\phi(f)} \text{ and } B_\phi = \bigcap_{f \in F} \sigma(f)^{-1}B_{\phi(f)} \text{ for each } \phi \in \text{Map}(F, k).$$

Now for each $\varepsilon > 0$, let $AP_\mu(\sigma, \alpha : F, \varepsilon)$ (or just $AP(\sigma, \alpha : F, \varepsilon)$ if there is no any ambiguity) be the set of all partitions $\beta = \{B_1, \dots, B_k\}$ of $\{1, \dots, d\}$ with $d_F(\alpha, \beta) \leq \varepsilon$. In particular, $|AP(\sigma, \alpha : F, \varepsilon)| \leq k^d$. We define

$$\begin{aligned} H_{\mu, \Sigma}(\alpha : F, \varepsilon) &= \limsup_{i \rightarrow \infty} \frac{1}{d_i} \log |AP(\sigma_i, \alpha : F, \varepsilon)| \leq \log |\alpha|, \\ H_{\mu, \Sigma}(\alpha : F) &= \lim_{\varepsilon \rightarrow 0} H_{\mu, \Sigma}(\alpha : F, \varepsilon) = \inf_{\varepsilon > 0} H_{\mu, \Sigma}(\alpha : F, \varepsilon) \leq \log |\alpha|, \\ h_{\mu, \Sigma}(\alpha) &= \inf_{F \in \mathfrak{F}_G} H_{\mu, \Sigma}(\alpha : F) \leq \log |\alpha|. \end{aligned}$$

Observe that $AP(\sigma_i, \alpha : F, \varepsilon)$ may be empty, and in the case that $AP(\sigma_i, \alpha : F, \varepsilon) = \emptyset$ for all large enough $i \in \mathbb{N}$ we have $H_{\mu, \Sigma}(\alpha : F, \varepsilon) = -\infty$ by the convention $\log 0 = -\infty$. Hence, $h_{\mu, \Sigma}(\alpha)$ may take a value in $[0, \log |\alpha|] \cup \{-\infty\}$.

The main result of [Bow10] tells us that, if there exists an $\alpha \in \mathcal{P}_X$ generating the σ -algebra \mathcal{B}_X (in the sense of μ) then the quantity $h_{\mu, \Sigma}(\alpha)$ is independent of the selection of such a partition, and this quantity, denoted by $h_{\mu, \Sigma}(X, G)$, is called the *measure-theoretic μ -entropy of (X, G)* . Indeed, L. Bowen defined the measure-theoretic entropy in a more general case when the action admits a generating partition β (not necessary finite) with finite Shannon entropy [Bow10]. We say that the partition $\beta \subset \mathcal{B}_X$ *generates the σ -algebra \mathcal{B}_X (in the sense of μ)* if for each $B \in \mathcal{B}_X$ there exists $A \in \mathcal{A}$ such that $\mu(A \Delta B) = 0$, where \mathcal{A} is the smallest G -invariant sub- σ -algebra of \mathcal{B}_X containing β .

Observing that, for an expansive action (X, G) of sofic groups with an expansive constant $\delta > 0$, if $\xi \in \mathcal{P}_X$ satisfies $\text{diam}(\xi) < \delta$, where $\text{diam}(\xi)$ denotes the maximal diameter of subsets in ξ , then, for each $\mu \in M(X, G)$ (if $M(X, G) \neq \emptyset$), ξ generates \mathcal{B}_X [Wal82, Theorem 5.25], and so the quantity $h_{\mu, \Sigma}(X, G)$ is well defined.

For technical reasons for $r_1, r_2 \in [-\infty, \infty]$ we set $r_1 + r_2 = -\infty$ by convention in the case that either $r_1 = -\infty$ or $r_2 = -\infty$, and for $r_1, r_2 \in (-\infty, \infty]$ we set $r_1 + r_2 = \infty$ by convention in the case that either $r_1 = \infty$ or $r_2 = \infty$.

We say that a function $f : Y \rightarrow [-\infty, \infty)$ defined over a compact metric space Y is *upper semi-continuous* if $\limsup_{y' \rightarrow y} f(y') \leq f(y)$ for each $y \in Y$. The following result shows that each expansive action of sofic groups admits a measure with maximal entropy.

Theorem 2.1. *Let (X, G) be an expansive action of sofic groups with $M(X, G) \neq \emptyset$. Then $h_{\bullet, \Sigma}(X, G) : M(X, G) \rightarrow [0, \infty) \cup \{-\infty\}$ is an upper semi-continuous function.*

Proof. The proof is inspired by [Wal82, Theorem 8.2].

Let $\delta > 0$ be an expansive constant for (X, G) and $\xi \in \mathcal{P}_X$ with $\text{diam}(\xi) < \delta$. Then ξ generates \mathcal{B}_X and so $h_{\mu, \Sigma}(X, G) \in [0, \log |\xi|] \cup \{-\infty\}$ for each $\mu \in M(X, G)$.

Now fix $\eta > 0$ and $\mu \in M(X, G)$. It suffices to find an open set $U \subset M(X, G)$ containing μ such that $h_{\nu, \Sigma}(X, G) \leq h_{\mu, \Sigma}(X, G) + \eta$ for each $\nu \in U$.

We choose $F \in \mathfrak{F}_G$ and $\varepsilon > 0$ such that $H_{\mu, \Sigma}(\xi : F, 2\varepsilon) \leq h_{\mu, \Sigma}(X, G) + \eta$. Say $\xi = \{A_1, \dots, A_k\}$ and let $0 < \varepsilon_1 < \frac{\varepsilon}{2M^2}$ with $M = |\text{Map}(F, k)| = k^{|F|}$. Let $\phi \in \text{Map}(F, k)$. Since μ is regular, there exists a compact set $K_\phi \subset A_\phi$ with $\mu(A_\phi \setminus K_\phi) < \varepsilon_1$, and then for each $i = 1, \dots, k$ we define

$$L_i = \bigcup_{f \in F} \{fK_\phi : \phi(f) = i\} \subset A_i.$$

Then L_1, \dots, L_k are pairwise disjoint compact subsets of X , and so there exists $\xi' = \{A'_1, \dots, A'_k\} \in \mathcal{P}_X$ such that $\text{diam}(\xi') < \delta$ and, for each $j = 1, \dots, k$, $L_j \subset \text{int}(A'_j)$ where $\text{int}(A'_j)$ denotes the interior of A'_j . Observe that

$$K_\phi \subset \text{int} \left(\bigcap_{f \in F} f^{-1} A'_{\phi(f)} \right) = \text{int}(A'_\phi) \left(\text{here } A'_\phi = \bigcap_{f \in F} f^{-1} A'_{\phi(f)} \right)$$

by the construction of ξ' , and so using Urysohn's Lemma we can choose $u_\phi \in C(X)$, where $C(X)$ denotes the set of all real-valued continuous functions over X , with $0 \leq u_\phi \leq 1$ which equals 1 on K_ϕ and vanishes on $X \setminus \text{int}(A'_\phi)$. Set

$$U = \{\nu \in M(X, G) : |\nu(u_\phi) - \mu(u_\phi)| < \varepsilon_1 \text{ for all } \phi \in \text{Map}(F, k)\}$$

which is an open set of $M(X, G)$ containing μ . Let $\nu \in U$. Then $\nu(A'_\phi) \geq \nu(u_\phi) > \mu(u_\phi) - \varepsilon_1 \geq \mu(K_\phi) - \varepsilon_1$ and hence $\mu(A_\phi) - \nu(A'_\phi) < 2\varepsilon_1$ for each $\phi \in \text{Map}(F, k)$. Observe $\{A_\phi : \phi \in \text{Map}(F, k)\} \in \mathcal{P}_X$ and $\{A'_\phi : \phi \in \text{Map}(F, k)\} \in \mathcal{P}_X$. Note that if $p_1, \dots, p_m, q_1, \dots, q_m, c$ are nonnegative real numbers with $m \in \mathbb{N}$ such that $\sum_{i=1}^m p_i = \sum_{i=1}^m q_i = 1$ and $p_j - q_j < c$ for each $j = 1, \dots, m$ then

$$q_i - p_i = \sum_{j \neq i} (p_j - q_j) < mc$$

and hence $|p_i - q_i| < mc$ for any $i = 1, \dots, m$. This implies $|\nu(A'_\phi) - \mu(A_\phi)| < 2\varepsilon_1 M$ for each $\phi \in \text{Map}(F, k)$, and so

$$\sum_{\phi \in \text{Map}(F, k)} |\nu(A'_\phi) - \mu(A_\phi)| \leq 2\varepsilon_1 M^2 \leq \varepsilon.$$

Thus $AP_\nu(\sigma_i, \xi' : F, \varepsilon) \subset AP_\mu(\sigma_i, \xi : F, 2\varepsilon)$ for each $i \in \mathbb{N}$, and then

$$H_{\nu, \Sigma}(\xi' : F) \leq H_{\nu, \Sigma}(\xi' : F, \varepsilon) \leq H_{\mu, \Sigma}(\xi : F, 2\varepsilon) \leq h_{\mu, \Sigma}(X, G) + \eta.$$

As $\text{diam}(\xi') < \delta$, $\xi' \in \mathcal{P}_X$ generates \mathcal{B}_X by the construction of δ , and so we get $h_{\nu, \Sigma}(X, G) \leq h_{\mu, \Sigma}(X, G) + \eta$ for each $\nu \in U$ as desired. This finishes the proof. \square

In the spirit of L. Bowen's entropy as above, Kerr and Li introduced alternatively the sofic measure-theoretic entropy [KL11, KL13b] as follows.

Let (Y, ρ) be a metric space and $\varepsilon > 0$. A set $\emptyset \neq A \subset Y$ is said to be (ρ, ε) -separated if $\rho(x, y) \geq \varepsilon$ for all distinct $x, y \in A$. We write $N_\varepsilon(Y, \rho)$ for the maximal cardinality of finite non-empty (ρ, ε) -separated subsets of Y (and set $N_\varepsilon(\emptyset, \rho) = 0$ by convention). A basic fact is that if $\emptyset \neq A \subset Y$ is a maximal finite (ρ, ε) -separated subset of Y then for each $y \in Y$ there exists $x \in A$ such that $\rho(x, y) < \varepsilon$.

For each $d \in \mathbb{N}$ and $(x_1, \dots, x_d), (x'_1, \dots, x'_d) \in X^d$, we set

$$(2.1) \quad \rho_d((x_1, \dots, x_d), (x'_1, \dots, x'_d)) = \max_{i=1}^d \rho(x_i, x'_i).$$

For $F \in \mathfrak{F}_G, \delta > 0$ and $\sigma : G \rightarrow \text{Sym}(d), g \mapsto \sigma_g$ with $d \in \mathbb{N}$, put

$$X_{F, \delta, \sigma}^d = \left\{ (x_1, \dots, x_d) \in X^d : \max_{s \in F} \sqrt{\sum_{i=1}^d \frac{1}{d} \rho^2(sx_i, x_{\sigma_s(i)})} < \delta \right\};$$

and then for $\mu \in M(X)$ and $L \in \mathfrak{F}_{C(X)}$, we set

$$X_{F, \delta, \sigma, \mu, L}^d = \left\{ (x_1, \dots, x_d) \in X_{F, \delta, \sigma}^d : \max_{f \in L} \left| \frac{1}{d} \sum_{i=1}^d f(x_i) - \mu(f) \right| < \delta \right\}.$$

By [KL13b, Proposition 3.4] the *measure-theoretic μ -entropy of (X, G)* can be defined as (recalling the convention $\log 0 = -\infty$)

$$h_\mu(G, X) = \sup_{\varepsilon > 0} \inf_{L \in \mathfrak{F}_{C(X)}} \inf_{F \in \mathfrak{F}_G} \inf_{\delta > 0} \limsup_{i \rightarrow \infty} \frac{1}{d_i} \log N_\varepsilon(X_{F, \delta, \sigma_i, \mu, L}^{d_i}, \rho_{d_i}).$$

The sofic measure-theoretic entropy can be defined equivalently using finite open covers as follows [Zha12, §2]. We remark that it was Li who used firstly open covers for actions of sofic groups to consider sofic mean dimension in [Li13].

Let $\mathcal{U} \in \mathcal{C}_X$, we set (recalling the convention $\log 0 = -\infty$)

$$h_{F, \delta, \mu, L}(G, \mathcal{U}) = \limsup_{i \rightarrow \infty} \frac{1}{d_i} \log N(\mathcal{U}^{d_i}, X_{F, \delta, \sigma_i, \mu, L}^{d_i}).$$

In particular, $h_{F,\delta,\mu,L}(G, \mathcal{U})$ takes the value $-\infty$ once $X_{F,\delta,\sigma_i,\mu,L}^{d_i} = \emptyset$ for all $i \in \mathbb{N}$ large enough. Now we define the *measure-theoretic μ -entropy of \mathcal{U}* as

$$h_\mu(G, \mathcal{U}) = \inf_{L \in \mathfrak{F}_{C(X)}} \inf_{F \in \mathfrak{F}_G} \inf_{\delta > 0} h_{F,\delta,\mu,L}(G, \mathcal{U}) \leq \log N(\mathcal{U}, X).$$

It is not hard to check that

$$h_\mu(G, X) = \sup_{\mathcal{U} \in \mathcal{C}_X^\circ} h_\mu(G, \mathcal{U}).$$

Moreover, by the proof of [KL11, Theorem 6.1], it was proved implicitly $h_\mu(G, X) = -\infty$ (and hence $h_\mu(G, \mathcal{U}) = -\infty$ for all $\mathcal{U} \in \mathcal{C}_X^\circ$) for each $\mu \in M(X) \setminus M(X, G)$.

Observe that both of $h_\mu(G, \mathcal{U})$ and $h_\mu(G, X)$ may take the value of $-\infty$, and by [KL11, KL13b] if $\mu \in M(X, G)$ and \mathcal{B}_X admits a generating partition (in the sense of μ) with finite Shannon entropy then $h_\mu(G, X)$ is just the quantity $h_{\mu,\Sigma}(X, G)$ introduced before.

The following result is easy to obtain:

Proposition 2.2. *Let $\mathcal{U} \in \mathcal{C}_X^\circ$. Then $h_\bullet(G, \mathcal{U}) : M(X) \rightarrow [0, \log N(\mathcal{U}, X)] \cup \{-\infty\}$ is an upper semi-continuous function.*

Proof. Let $\mu \in M(X)$. For any $\varepsilon > 0$ we may choose $L \in \mathfrak{F}_{C(X)}$, $F \in \mathfrak{F}_G$ and $\delta > 0$ such that $h_{F,2\delta,\mu,L}(G, \mathcal{U}) \leq h_\mu(G, \mathcal{U}) + \varepsilon$. Now we consider the non-empty open set

$$\mu \in V = \{\nu \in M(X) : |\nu(f) - \mu(f)| < \delta \text{ for all } f \in L\}.$$

Then for each $\nu \in V$ we have $X_{F,\delta,\sigma,\nu,L}^d \subset X_{F,2\delta,\sigma,\mu,L}^d$ for each $\sigma : G \rightarrow \text{Sym}(d)$, $d \in \mathbb{N}$, which implies $h_\nu(G, \mathcal{U}) \leq h_{F,\delta,\nu,L}(G, \mathcal{U}) \leq h_{F,2\delta,\mu,L}(G, \mathcal{U}) \leq h_\mu(G, \mathcal{U}) + \varepsilon$. This implies that the considered function is upper semi-continuous. \square

3. WEAK EXPANSIVENESS FOR ACTIONS OF SOFIC GROUPS

Note that, when considering a continuous mapping over a compact metric space, since the introduction of h -expansiveness and asymptotical h -expansiveness, both of them turn out to be very important classes in the research area of dynamical systems. It is shown by R. Bowen [Bow72] that positively expansive systems, expansive homeomorphisms, endomorphisms of a compact Lie group and Axiom A diffeomorphisms are all h -expansive, by Misiurewicz [Mis76] that every continuous endomorphism of a compact metric group is asymptotically h -expansive if it has finite entropy, and by Buzzi [Buz97] that any C^∞ diffeomorphism on a compact manifold is asymptotically h -expansive. Moreover, there are more nice characterizations of asymptotical h -expansiveness obtained recently for this setting. For example, a topological dynamical system is asymptotically h -expansive if and only if it admits a principal extension to a symbolic system by Boyle and Downarowicz [BD04], i.e., a symbolic extension which preserves entropy for each invariant measure; if and

only if it is hereditarily uniformly lowerable by Huang, Ye and the second author of present paper [HYZ10] (for a detailed definition of the hereditarily uniformly lowerable property and its more story see [HYZ10]).

In this section we explore similar weak expansiveness for actions of sofic groups.

By [KL13b, Proposition 2.4] the *topological entropy of* (X, G) can be defined as

$$h(G, X) = \sup_{\varepsilon > 0} \inf_{F \in \mathfrak{F}_G} \inf_{\delta > 0} \limsup_{i \rightarrow \infty} \frac{1}{d_i} \log N_\varepsilon (X_{F, \delta, \sigma_i}^{d_i}, \rho_{d_i}),$$

which is introduced and discussed in [KL11, KL13b]. Before proceeding, we need to recall the topological entropy for actions of sofic groups introduced in [Zha12, §2] using finite open covers. Let $\mathcal{U} \in \mathcal{C}_X$. For $F \in \mathfrak{F}_G$ and $\delta > 0$ we set

$$h_{F, \delta}(G, \mathcal{U}) = \limsup_{i \rightarrow \infty} \frac{1}{d_i} \log N(\mathcal{U}^{d_i}, X_{F, \delta, \sigma_i}^{d_i}).$$

Observe again that $h_{F, \delta}(G, \mathcal{U})$ takes the value of $-\infty$ whenever $X_{F, \delta, \sigma_i}^{d_i} = \emptyset$ for all $i \in \mathbb{N}$ large enough. Now we define the *topological entropy of* \mathcal{U} as

$$h(G, \mathcal{U}) = \inf_{F \in \mathfrak{F}_G} \inf_{\delta > 0} h_{F, \delta}(G, \mathcal{U}) \leq \log N(\mathcal{U}, X).$$

It is not hard to check that

$$h(G, X) = \sup_{\mathcal{U} \in \mathcal{C}_X^\circ} h(G, \mathcal{U}).$$

Observe that both of $h(G, \mathcal{U})$ and $h(G, X)$ may take the value of $-\infty$.

The sofic topological entropy and sofic measure-theoretic entropy are related to each other [KL11, Theorem 6.1] and [Zha12, Theorem 4.1]: for $\mathcal{U} \in \mathcal{C}_X^\circ$,

$$(3.1) \quad h(G, X) = \sup_{\mu \in M(X, G)} h_\mu(G, X) \text{ and } h(G, \mathcal{U}) = \max_{\mu \in M(X, G)} h_\mu(G, \mathcal{U}),$$

where in the right-hand sides as above we set it as $-\infty$ by convention if $M(X, G) = \emptyset$.

In the spirit of Misiurewicz [Mis76], the above idea can be used to introduce weak expansiveness of h -expansiveness and asymptotical h -expansiveness for actions of sofic groups. Let $\mathcal{U}_1, \mathcal{U}_2 \in \mathcal{C}_X$. For $F \in \mathfrak{F}_G$ and $\delta > 0$ we set

$$h_{F, \delta}(G, \mathcal{U}_1 | \mathcal{U}_2) = \limsup_{i \rightarrow \infty} \frac{1}{d_i} \log \max_{V \in \mathcal{U}_2^{d_i}} N(\mathcal{U}_1^{d_i}, X_{F, \delta, \sigma_i}^{d_i} \cap V) \leq h_{F, \delta}(G, \mathcal{U}_1),$$

$$h(G, \mathcal{U}_1 | \mathcal{U}_2) = \inf_{F \in \mathfrak{F}_G} \inf_{\delta > 0} h_{F, \delta}(G, \mathcal{U}_1 | \mathcal{U}_2) \leq h(G, \mathcal{U}_1),$$

$$h(G, X | \mathcal{U}_2) = \sup_{\mathcal{U}_1 \in \mathcal{C}_X^\circ} h(G, \mathcal{U}_1 | \mathcal{U}_2) \leq h(G, X),$$

$$h^*(G, X) = \inf_{\mathcal{U}_2 \in \mathcal{C}_X^\circ} h(G, X | \mathcal{U}_2) \leq h(G, X).$$

Then $h(G, X | \{X\}) = h(G, X)$ by the definitions. We say that (X, G) is *h-expansive* if $h(G, X | \mathcal{U}) \leq 0$ for some $\mathcal{U} \in \mathcal{C}_X^\circ$, and *asymptotically h-expansive* if $h^*(G, X) \leq 0$.

Each h -expansive action of sofic groups is asymptotically h -expansive by the definitions. The next result shows that each expansive action of sofic groups is h -expansive, and thus these two kinds of expansiveness are indeed weak expansiveness.

Theorem 3.1. *Let (X, G) be an expansive action of sofic groups with $\kappa > 0$ an expansive constant and $\mathcal{U} \in \mathcal{C}_X$. Assume that $\text{diam}(\mathcal{U}) \leq c\kappa$ for some $c < 1$. Then $h(G, X|\mathcal{U}) \leq 0$.*

Proof. Let $\mathcal{V} \in \mathcal{C}_X^o$ and $\varepsilon > 0$. It suffices to prove that $h(G, \mathcal{V}|\mathcal{U}) \leq \varepsilon$.

Let $\tau > 0$ be a Lebesgue number of \mathcal{V} . As (X, G) is an expansive action of sofic groups with $\kappa > 0$ an expansive constant, it is not hard to choose $F \in \mathfrak{F}_G$ such that $\max_{s \in F} \rho(sx, sx') < \kappa$ implies $\rho(x, x') < \frac{\tau}{2}$ (for example see [Wal82, Chapter 5, §5.6]). Now let $\delta > 0$ be small enough such that

$$(3.2) \quad |\Theta_d| \cdot |\mathcal{V}|^{\frac{8\delta^2|F|}{(1-c)^2\kappa^2} \cdot d} < e^{\varepsilon d}$$

for all $d \in \mathbb{N}$ large enough, where Θ_d is the set of all subsets θ of $\{1, \dots, d\}$ with

$$|\theta| < \frac{8\delta^2|F|}{(1-c)^2\kappa^2} \cdot d.$$

For any map $\sigma : G \rightarrow \text{Sym}(d), g \mapsto \sigma_g$ with $d \in \mathbb{N}$, recall

$$X_{F,\delta,\sigma}^d = \left\{ (x_1, \dots, x_d) \in X^d : \max_{s \in F} \sqrt{\sum_{i=1}^d \frac{1}{d} \rho^2(sx_i, x_{\sigma_s(i)})} < \delta \right\},$$

in particular, once $(x_1^*, \dots, x_d^*) \in X_{F,\delta,\sigma}^d$ then, by direct calculations,

$$(3.3) \quad \left| \left\{ i \in \{1, \dots, d\} : \rho(sx_i^*, x_{\sigma_s(i)}^*) \geq \frac{(1-c)\kappa}{2} \text{ for some } s \in F \right\} \right| < \frac{4\delta^2|F| \cdot d}{(1-c)^2\kappa^2}.$$

Now let $V \in \mathcal{U}^d$ and say $(x_1, \dots, x_d) \in X_{F,\delta,\sigma}^d \cap V$ (if it is not empty). For any $(x'_1, \dots, x'_d) \in X_{F,\delta,\sigma}^d \cap V$, applying (3.3) to (x_1, \dots, x_d) and (x'_1, \dots, x'_d) and observing $\rho(sx_i, sx'_i) \leq \rho(sx_i, x_{\sigma_s(i)}) + \rho(x_{\sigma_s(i)}, x'_{\sigma_s(i)}) + \rho(sx'_i, x'_{\sigma_s(i)})$, it is easy to see

$$|\{i \in \{1, \dots, d\} : \rho(sx_i, sx'_i) \geq \kappa \text{ for some } s \in F\}| < \frac{8\delta^2|F|}{(1-c)^2\kappa^2} \cdot d.$$

In other words, if we associate each $\theta \in \Theta_d$ with $X_{F,\delta,\sigma,V,\theta}^d$, the set of all (x'_1, \dots, x'_d) in $X_{F,\delta,\sigma}^d \cap V$ with $\theta = |\{i \in \{1, \dots, d\} : \rho(sx_i, sx'_i) \geq \kappa \text{ for some } s \in F\}|$, then

$$(3.4) \quad \bigcup_{\theta \in \Theta_d} X_{F,\delta,\sigma,V,\theta}^d = X_{F,\delta,\sigma}^d \cap V.$$

For each $\theta \in \Theta_d$, if $(x_1'', \dots, x_d''), (x_1^\#, \dots, x_d^\#) \in X_{F,\delta,\sigma,V,\theta}^d$ then, by the selection of F , $\rho(x_i'', x_i) < \frac{\tau}{2}, \rho(x_i^\#, x_i) < \frac{\tau}{2}$ and then $\rho(x_i'', x_i^\#) < \tau$ for all $i \in \{1, \dots, d\} \setminus \theta$,

thus $X_{F,\delta,\sigma,V,\theta}^d$ can be covered by at most $|\mathcal{V}|^{|\theta|}$ many elements of \mathcal{V}^d . This implies

$$\begin{aligned} N(\mathcal{V}^d, X_{F,\delta,\sigma}^d \cap V) &\leq \sum_{\theta \in \Theta_d} N(\mathcal{V}^d, X_{F,\delta,\sigma,V,\theta}^d) \text{ (using (3.4))} \\ &\leq |\Theta_d| \cdot |\mathcal{V}|^{\frac{8\delta^2|F|}{(1-c)^2\kappa^2} \cdot d} < e^{\varepsilon d} \text{ (using (3.2)),} \end{aligned}$$

and hence $h(G, \mathcal{V}|\mathcal{U}) \leq \varepsilon$ by the definition, finishing the proof. \square

Observing that, for $\mathcal{V}_1, \mathcal{V}_2 \in \mathcal{C}_X$ and $K \subset X$,

$$(3.5) \quad N(\mathcal{V}_1, K) \leq N(\mathcal{V}_2, K) \cdot \max_{V \in \mathcal{V}_2} N(\mathcal{V}_1, K \cap V),$$

it is direct to obtain the following easy while useful observation.

Lemma 3.2. *Let $\mathcal{U}_1, \mathcal{U}_2 \in \mathcal{C}_X$ and $\mu \in M(X)$. Then*

$$h_\mu(G, \mathcal{U}_1) \leq h_\mu(G, \mathcal{U}_2) + h(G, \mathcal{U}_1|\mathcal{U}_2) \text{ and } h_\mu(G, X) \leq h_\mu(G, \mathcal{U}_2) + h(G, X|\mathcal{U}_2),$$

$$h(G, \mathcal{U}_1) \leq h(G, \mathcal{U}_2) + h(G, \mathcal{U}_1|\mathcal{U}_2) \text{ and } h(G, X) \leq h(G, \mathcal{U}_2) + h(G, X|\mathcal{U}_2).$$

Proof. Let $\{F_n : n \in \mathbb{N}\} \subset \mathfrak{F}_G$ increase to the whole group G and $\{\delta_n > 0 : n \in \mathbb{N}\}$ decrease to 0. By the definitions it is direct to see

$$h_\mu(G, \mathcal{U}) = \inf_{L \in \mathfrak{F}_{C(X)}} \lim_{n \rightarrow \infty} h_{F_n, \delta_n, \mu, L}(G, \mathcal{U})$$

for each $\mathcal{U} \in \mathcal{C}_X$ and

$$h(G, \mathcal{U}_1|\mathcal{U}_2) = \lim_{n \rightarrow \infty} h_{F_n, \delta_n}(G, \mathcal{U}_1|\mathcal{U}_2).$$

Now for each $n \in \mathbb{N}$ and $L \in \mathfrak{F}_{C(X)}$, using (3.5) it is easy to obtain

$$h_{F_n, \delta_n, \mu, L}(G, \mathcal{U}_1) \leq h_{F_n, \delta_n, \mu, L}(G, \mathcal{U}_2) + h_{F_n, \delta_n}(G, \mathcal{U}_1|\mathcal{U}_2),$$

and then $h_\mu(G, \mathcal{U}_1) \leq h_\mu(G, \mathcal{U}_2) + h(G, \mathcal{U}_1|\mathcal{U}_2)$ by taking limits on both sides, in particular, $h_\mu(G, \mathcal{U}_1) = -\infty$ when $h_\mu(G, \mathcal{U}_2) = -\infty$ (and hence $h_\mu(G, X) = -\infty$ if and only if $h_\mu(G, \mathcal{U}_2) = -\infty$), which implies $h_\mu(G, X) \leq h_\mu(G, \mathcal{U}_2) + h(G, X|\mathcal{U}_2)$ even if $h(G, X|\mathcal{U}_2) = \infty$ (recalling our technical convention of $-\infty + \infty = -\infty$). The remaining items can be proved similarly. \square

As direct corollaries, we have:

Corollary 3.3. *$h^*(G, X) < \infty$ if and only if $h(G, X) < \infty$, and $h^*(G, X) = -\infty$ if and only if $h(G, X) = -\infty$.*

Corollary 3.4. *Assume that (X, G) is h -expansive. Then there exists $\mathcal{U} \in \mathcal{C}_X^o$ with*

$$h(G, X) = h(G, \mathcal{U}) \text{ and } h_\mu(G, X) = h_\mu(G, \mathcal{U}) \text{ for each } \mu \in M(X).$$

Moreover, we can prove the following result.

Theorem 3.5. *Let $\mu \in M(X)$. Then*

$$(3.6) \quad \limsup_{\nu \rightarrow \mu} h_\nu(G, X) \leq h_\mu(G, X) + h^*(G, X).$$

In particular, if (X, G) is asymptotically h -expansive then the function $h_\bullet(G, X) : M(X) \rightarrow [0, \infty) \cup \{-\infty\}$ is upper semi-continuous.

Proof. Let $\mathcal{U} \in \mathcal{C}_X^o$. By Proposition 2.2 and Lemma 3.2 we have

$$\begin{aligned} \limsup_{\nu \rightarrow \mu} h_\nu(G, X) &\leq \limsup_{\nu \rightarrow \mu} h_\nu(G, \mathcal{U}) + h(G, X|\mathcal{U}) \\ &\leq h_\mu(G, \mathcal{U}) + h(G, X|\mathcal{U}) \leq h_\mu(G, X) + h(G, X|\mathcal{U}). \end{aligned}$$

Then (3.6) follows directly by taking the infimum over all $\mathcal{U} \in \mathcal{C}_X^o$.

Now we assume that (X, G) is asymptotically h -expansive, that is, $h^*(G, X) \leq 0$. Then $h(G, X) < \infty$ by Corollary 3.3, and hence $h_\eta(G, X) \in [0, \infty) \cup \{-\infty\}$ for each $\eta \in M(X)$ by (3.1) (recalling the fact that $h_\eta(G, X) = -\infty$ for each $\eta \in M(X) \setminus M(X, G)$). Moreover, using (3.6) we obtain

$$\limsup_{\nu \rightarrow \mu} h_\nu(G, X) \leq h_\mu(G, X) + h^*(G, X) \leq h_\mu(G, X)$$

from our technical convention. This finishes the proof. \square

Remark 3.6. *As a direct corollary of Theorem 3.5, we have: if the action (X, G) is asymptotically h -expansive then both of its sofic topological mean dimension and its sofic metric mean dimension with respect to any compatible metric are at most zero [Li13, Proposition 4.3 and Theorem 6.1]. For the definition of sofic topological mean dimension and sofic metric mean dimension see [Li13, §2 and §4], respectively.*

Combining Theorem 3.5 with (3.1) one has:

Corollary 3.7. *Each asymptotically h -expansive action of sofic groups admits a measure with maximal entropy.*

In general, the converse does not hold, for example, any \mathbb{Z} -action with infinite entropy admits a measure with infinite entropy, whereas, it is not asymptotically h -expansive. Furthermore, [Mis76, Example 6.4] shows us a \mathbb{Z} -action with finite entropy such that it is not asymptotically h -expansive, while its each invariant measure has maximal entropy. Observe that, as we shall show later, the definitions of weak expansiveness given here for actions of sofic groups are equivalent to definitions given in the same spirit of Misiurewicz's ideas when the group is amenable.

Remark 3.8. *Theorem 2.1 is a consequence of Theorem 3.1 and Theorem 3.5.*

4. PROFINITE ACTIONS

In this section we provide our first interesting non-trivial h -expansive action of sofic groups using the language of the profinite action.

Recall that the action (X, G) is *distal* if $\inf_{g \in G} \rho(gx, gy) > 0$ for all distinct $x, y \in X$, and *equicontinuous* if for each $\delta > 0$ there exists $\varepsilon > 0$ such that $\rho(x, y) \leq \varepsilon$ implies $\rho(gx, gy) \leq \delta$ for all $g \in G$.

The following result should be known, we provide here a proof for completeness.

Lemma 4.1. *Assume that the action (X, G) is equicontinuous. Then it is distal. And if additionally X is infinite then it is not expansive.*

Proof. First we prove that (X, G) is distal. Else, there exist distinct points $x_1, x_2 \in X$ with $\inf_{g \in G} \rho(gx_1, gx_2) = 0$. As (X, G) is equicontinuous, there exists $\varepsilon > 0$ such that $\rho(x, y) < \varepsilon$ implies $\rho(gx, gy) \leq \frac{1}{2}\rho(x_1, x_2)$ for each $g \in G$. Say $g' \in G$ with $\rho(g'x_1, g'x_2) < \varepsilon$, and then $0 < \rho(x_1, x_2) = \rho((g')^{-1}g'x_1, (g')^{-1}g'x_2) \leq \frac{1}{2}\rho(x_1, x_2)$ by the selection of ε , a contradiction.

Now additionally we assume that X is infinite. If (X, G) is expansive, then there exists $\delta > 0$ with $\sup_{g \in G} \rho(gx, gy) > \delta$ for all distinct $x, y \in X$. Using again the equicontinuity of (X, G) , we could choose $\varepsilon' > 0$ such that $\rho(x, y) < \varepsilon'$ implies $\rho(gx, gy) \leq \delta$ for each $g \in G$. As X is infinite, by the compactness of X we could choose distinct points $y_1, y_2 \in X$ with $\rho(y_1, y_2) < \varepsilon'$, a contradiction to the selection of δ . That is, (X, G) is not expansive, finishing the proof. \square

See [Aus88] for a more detailed story of distal actions and equicontinuous actions.

Observe that, by the definitions each action of sofic groups is h -expansive if it has topological entropy at most zero, and Kerr and Li proved that each distal action of sofic groups has topological entropy at most zero [KL13a, Corollary 8.5]. Thus each distal action of sofic groups is h -expansive. With help of this observation, we can provide our first interesting h -expansive example.

Let G be a countable group. A *chain* of G is a sequence $G = G_0 \geq G_1 \geq \dots$ of subgroups with finite indices in G . For a chain (G_n) , we have a tree structure $T(G, (G_n))$ defined naturally as follows. The vertices are $\{gG_n : n \in \mathbb{N}, g \in G\}$ and (g_1G_n, g_2G_m) is an edge if $m = n + 1$ and $g_2G_m \subset g_1G_n$. The boundary $\partial T(G, (G_n))$ of $T(G, (G_n))$ consists of all sequences (x_0, x_1, \dots) of vertices with x_n adjacent to x_{n+1} for each $n \in \mathbb{Z}_+$. Then $\partial T(G, (G_n))$ is a compact metrizable space endowed with the topology generated by the open basis consisting of all subsets $O_x = \{(x_0, x_1, \dots) \in \partial T(G, (G_n)) : x_N = x\}$ with $x \in G/G_N$ and $N \in \mathbb{Z}_+$. The natural left actions of G on G/G_n induce the *profinite action* $(\partial T(G, (G_n)), G)$, an action of G on $\partial T(G, (G_n))$ by homeomorphisms. In this case the profinite action

$(\partial T(G, (G_n)), G)$ is equicontinuous, since for any $x = gG_n \in G/G_n$ and $h \in G$ we have $hO_x = O_y$ with $y = hgG_n \in G/G_n$.

Combining with the above discussions, we obtain:

Proposition 4.2. *Let G be a countable sofic group and (G_n) a chain of G . Then the profinite action of G on $\partial T(G, (G_n))$ is h -expansive. Furthermore if $\partial T(G, (G_n))$ of $T(G, (G_n))$ is infinite then the action is not expansive.*

Dynamical properties of profinite actions have been studied extensively for residually finite groups in [AE12, AN12]. They were also used to investigate orbit equivalence rigidity for Kazhdan property (T) groups [Ioa11, OP10].

5. TAIL ENTROPY FOR ACTIONS OF COUNTABLE DISCRETE AMENABLE GROUPS

In this section we shall introduce tail entropy for actions of countable discrete amenable groups in the same spirit of Misiurewicz.

Recall that G is a countable discrete group. Denote by e_G the unit of G . G is called *amenable*, if there exists a sequence $\{F_n : n \in \mathbb{N}\} \subset \mathfrak{F}_G$, called a *Følner sequence* of G , such that

$$\lim_{n \rightarrow \infty} \frac{|gF_n \Delta F_n|}{|F_n|} = 0, \forall g \in G.$$

In the class of countable discrete groups, amenable groups include all solvable groups and groups with subexponential growth. In the group $G = \mathbb{Z}$, the sequence $F_n = \{0, 1, \dots, n-1\}$ defines a Følner sequence, as, indeed, does $\{a_n, a_n+1, \dots, a_n+n-1\}$ for any sequence $\{a_n\}_{n \in \mathbb{N}} \subset \mathbb{Z}$; and in the finite group G , if $\{F_n : n \in \mathbb{N}\}$ is a Følner sequence of G , then $F_n = G$ once n is large enough.

Throughout this section and next section, additionally, we will assume that G is always a countable discrete amenable group.

The well-known Ornstein-Weiss Lemma plays a crucial role in the study of entropy theory for actions of amenable groups [OW87]. The following version of it is taken from [Gro99b, 1.3.1]. For a more detailed story see the section of Appendix.

Proposition 5.1. *Let $f : \mathfrak{F}_G \rightarrow \mathbb{R}$ be a nonnegative function such that $f(Eg) = f(E)$ and $f(E \cup F) \leq f(E) + f(F)$ for all $E, F \in \mathfrak{F}_G$ and $g \in G$. Then for any Følner sequence $\{F_n : n \in \mathbb{N}\}$ of G the sequence $\left\{ \frac{f(F_n)}{|F_n|} : n \in \mathbb{N} \right\}$ converges and the value of the limit is independent of the selection of the Følner sequence $\{F_n : n \in \mathbb{N}\}$.*

Let $\emptyset \neq T \subset G$. We say that T *tiles* G if there exists $\emptyset \neq G_T \subset G$ such that $\{Tc : c \in G_T\}$ forms a partition of G , that is, $Tc_1 \cap Tc_2 = \emptyset$ if c_1 and c_2 are different elements from G_T and $\bigcup_{c \in G_T} Tc = G$. Denote by \mathcal{T}_G the set of all non-empty finite subsets of G which tile G . Observe that $\mathcal{T}_G \neq \emptyset$, as $\mathcal{T}_G \supset \{\{g\} : g \in G\}$.

As shown by [Wei03, Theorem 3.3 and Proposition 3.6], tiling sets play a key role in establishing a counterpart of Rokhlin's Lemma for actions of countable discrete amenable groups. The class of countable discrete amenable groups admitting a *tiling Følner sequence* (i.e., a Følner sequence consisting of tiling subsets of the group) is large, including all countable residually finite amenable groups [Wei01].

The following result is proved as [DZ, Proposition 2.5].

Proposition 5.2. *Let $f : \mathfrak{F}_G \rightarrow \mathbb{R}$ be a function and G admit a tiling Følner sequence. Assume that $f(Eg) = f(E)$ and $f(E \cup F) \leq f(E) + f(F)$ whenever $g \in G$ and $E, F \in \mathfrak{F}_G$ satisfy $E \cap F = \emptyset$. Then for any tiling Følner sequence $\{F_n : n \in \mathbb{N}\}$ of G , the sequence $\left\{ \frac{f(F_n)}{|F_n|} : n \in \mathbb{N} \right\}$ converges and the value of the limit is independent of the choice of the tiling Følner sequence $\{F_n : n \in \mathbb{N}\}$. Furthermore,*

$$\lim_{n \rightarrow \infty} \frac{f(F_n)}{|F_n|} = \inf_{F \in \mathfrak{F}_G} \frac{f(F)}{|F|} \quad \left(\text{and so } = \inf_{n \in \mathbb{N}} \frac{f(F_n)}{|F_n|} \right).$$

Let $\mathcal{W}_1, \mathcal{W}_2 \in \mathcal{C}_X$. If each element of \mathcal{W}_1 is contained in some element of \mathcal{W}_2 then we say that \mathcal{W}_1 is *finer than* \mathcal{W}_2 (denote by $\mathcal{W}_1 \succeq \mathcal{W}_2$ or $\mathcal{W}_2 \preceq \mathcal{W}_1$). The join $\mathcal{W}_1 \vee \mathcal{W}_2$ is given by $\mathcal{W}_1 \vee \mathcal{W}_2 = \{W_1 \cap W_2 : W_1 \in \mathcal{W}_1, W_2 \in \mathcal{W}_2\} \in \mathcal{C}_X$, which extends naturally to a finite collection of covers. Let $F \in \mathfrak{F}_G$, we set $(\mathcal{W}_1)_F = \bigvee_{g \in F} g^{-1} \mathcal{W}_1$, and then we consider a nonnegative function $m_{\mathcal{W}_1, \mathcal{W}_2} : \mathfrak{F}_G \rightarrow \mathbb{R}$ given by

$$m_{\mathcal{W}_1, \mathcal{W}_2}(F) = \max_{K \in (\mathcal{W}_2)_F} \log N((\mathcal{W}_1)_F, K) \text{ for each } F \in \mathfrak{F}_G.$$

It is easy to obtain the following useful observation.

Lemma 5.3. $m_{\mathcal{W}_1, \mathcal{W}_2}(E \cup F) \leq m_{\mathcal{W}_1, \mathcal{W}_2}(E) + m_{\mathcal{W}_1, \mathcal{W}_2}(F)$ for all $E, F \in \mathfrak{F}_G$.

Proof. Let $E, F \in \mathfrak{F}_G$. From the definition we choose $K \in (\mathcal{W}_2)_{E \cup F}$ with $m_{\mathcal{W}_1, \mathcal{W}_2}(E \cup F) = \log N((\mathcal{W}_1)_{E \cup F}, K)$. Say $K_1 \in (\mathcal{W}_2)_E$ and $K_2 \in (\mathcal{W}_2)_F$ with $K = K_1 \cap K_2$ (no matter if E and F are disjoint), such K_1 and K_2 must exist. Now let $\mathcal{V}_1 \subset (\mathcal{W}_1)_E$ cover K_1 with $|\mathcal{V}_1| = N((\mathcal{W}_1)_E, K_1)$ and let $\mathcal{V}_2 \subset (\mathcal{W}_1)_F$ cover K_2 with $|\mathcal{V}_2| = N((\mathcal{W}_1)_F, K_2)$. Obviously we can cover $K_1 \cap K_2$ (i.e. K) using the family $\mathcal{V}_1 \vee \mathcal{V}_2$. Observing that $|\mathcal{V}_1 \vee \mathcal{V}_2| \leq |\mathcal{V}_1| \cdot |\mathcal{V}_2|$ and each element of $\mathcal{V}_1 \vee \mathcal{V}_2$ is contained in some element of $(\mathcal{W}_1)_{E \cup F}$, we have that $N((\mathcal{W}_1)_{E \cup F}, K) \leq |\mathcal{V}_1 \vee \mathcal{V}_2| \leq N((\mathcal{W}_1)_E, K_1) \cdot N((\mathcal{W}_1)_F, K_2)$, which implies the conclusion directly. \square

It is easy to check G -invariance of the nonnegative function $m_{\mathcal{W}_1, \mathcal{W}_2} : \mathfrak{F}_G \rightarrow \mathbb{R}$. Observing Lemma 5.3, we could apply Proposition 5.1 to define

$$h^a(G, \mathcal{W}_1 | \mathcal{W}_2) = \lim_{n \rightarrow \infty} \frac{1}{|F_n|} m_{\mathcal{W}_1, \mathcal{W}_2}(F_n) \geq 0,$$

which is independent of the selection of the Følner sequence $\{F_n : n \in \mathbb{N}\}$. In particular, if G admits a tiling Følner sequence then by Proposition 5.2 one has

$$h^a(G, \mathcal{W}_1 | \mathcal{W}_2) = \inf_{F \in \mathcal{J}_G} \frac{1}{|F|} m_{\mathcal{W}_1, \mathcal{W}_2}(F) \left(\text{and so } = \inf_{n \in \mathbb{N}} \frac{1}{|F_n|} m_{\mathcal{W}_1, \mathcal{W}_2}(F_n) \right) \geq 0.$$

Then we define the *tail entropy of (X, G) with respect to \mathcal{W}_2* by

$$h^a(G, X | \mathcal{W}_2) = \sup_{\mathcal{U} \in \mathcal{C}_X^\circ} h^a(G, \mathcal{U} | \mathcal{W}_2),$$

and define the *topological entropy of (X, G)* , $h^a(G, X)$, and *tail entropy of (X, G)* , $h^{a,*}(G, X)$, respectively, as: $h^a(G, X) = h^a(G, X | \{X\})$ and

$$h^{a,*}(G, X) = \inf_{\mathcal{V} \in \mathcal{C}_X^\circ} h^a(G, X | \mathcal{V}) \left(= \inf_{\mathcal{V} \in \mathcal{C}_X^\circ} h^a(G, X | \overline{\mathcal{V}}) \right) \geq 0.$$

Recalling that in the special case of $G = \mathbb{Z}$ acting on a compact metric space X , equivalently, giving a homeomorphism $T : X \rightarrow X$, the above definition recovers the definition given by Misiurewicz in [Mis76], which was then used to discuss weak expansiveness for \mathbb{Z} -actions. In fact, Misiurewicz [Mis76] introduced tail entropy in the setting of giving a compact Hausdorff space and a continuous transformation of X into itself.

6. COMPARISON BETWEEN SOFIC AND AMENABLE CASES

The following result is the main result of this section, which shows that our definitions of weak expansiveness for actions of sofic groups are equivalent to definitions given in the same spirit of Misiurewicz's ideas when the group is amenable.

Theorem 6.1. *(X, G) is asymptotically h -expansive if and only if $h^{a,*}(G, X) = 0$. Similarly, (X, G) is h -expansive if and only if $h^a(G, X | \mathcal{V}) = 0$ for some $\mathcal{V} \in \mathcal{C}_X^\circ$.*

Let Y be a finite set, $\{A_i : i \in I\} \subset \{\emptyset\} \cup \mathfrak{F}_Y$ and $\delta \geq 0$. We say that $\{A_i : i \in I\}$ δ -covers Y if $\left| \bigcup_{i \in I} A_i \right| \geq \delta |Y|$. $\{A_i : i \in I\}$ are ε -disjoint if there exist pairwise disjoint subsets $B_i \subset A_i$ with $|B_i| \geq (1 - \varepsilon)|A_i|$ for each $i \in I$.

The next result is the Rokhlin Lemma for sofic approximations of countable discrete groups [KL13b, Lemma 4.5].

Lemma 6.2. *Let Γ be a countable group with the unit e and $0 \leq \tau < 1, 0 < \eta < 1$. Then there are an $l \in \mathbb{N}$ and $\eta', \eta'' > 0$ such that, whenever $e \in E_1 \subset \cdots \subset E_l$ are finite subsets of Γ with $|E_{k-1}^{-1} E_k \setminus E_k| \leq \eta' |E_k|$ for $k = 2, \dots, l$, there exists $e \in E \in \mathfrak{F}_\Gamma$ such that for every good enough sofic approximation $\sigma : \Gamma \rightarrow \text{Sym}(d)$ for Γ with some $d \in \mathbb{N}$ (i.e., $\sigma : \Gamma \rightarrow \text{Sym}(d)$ is a map such that*

$$\sigma_{st}(a) = \sigma_s \sigma_t(a), \sigma_s(a) \neq \sigma_{s'}(a), \sigma_e(a) = a$$

for all $a \in B$ with $B \subset \{1, \dots, d\}$ satisfying $|B| \geq (1 - \eta'')d$ and $s, t, s' \in E$ with $s \neq s'$, and any set $V \subset \{1, \dots, d\}$ with $|V| \geq (1 - \tau)d$, there exist $C_1, \dots, C_l \subset V$ such that

- (1) the sets $\sigma(E_k)C_k, k \in \{1, \dots, l\}$ are pairwise disjoint;
- (2) $\{\sigma(E_k)C_k : k \in \{1, \dots, l\}\}$ $(1 - \tau - \eta)$ -covers $\{1, \dots, d\}$;
- (3) $\{\sigma(E_k)c : c \in C_k\}$ is η -disjoint for each $k \in \{1, \dots, l\}$; and
- (4) for every $k \in \{1, \dots, l\}$ and $c \in C_k$, $E_k \ni s \mapsto \sigma_s(c)$ is bijective.

Before proceeding, we also need the following easy observation.

Recall that ρ is a compatible metric on the compact metric space X .

Lemma 6.3. *Let $K \subset X$ be a closed subset and $F \in \mathfrak{F}_G, \mathcal{U} \in \mathcal{C}_X^o$. Then there exists $\delta > 0$ such that*

$$K_{F,\delta} = \left\{ (x_s)_{s \in F} \in X^F : \max_{s \in F} \rho(x_s, sx) < \delta \text{ for some } x \in K \right\}$$

can be covered by at most $N(\mathcal{U}_F, K)$ elements of \mathcal{U}^F .

Proof. Obviously, there exists $\mathcal{V} \subset \mathcal{U}^F$ such that $|\mathcal{V}| \leq N(\mathcal{U}_F, K)$ and

$$\cup \mathcal{V} \supset K_F \text{ where } K_F = \{(sx)_{s \in F} : x \in K\}.$$

For example, let $\mathcal{W} \subset \mathcal{U}_F$ such that $|\mathcal{W}| = N(\mathcal{U}_F, K)$ and $\cup \mathcal{W} \supset K$. Now for each $W \in \mathcal{W}$, as $W \in \mathcal{U}_F$, say $W = \bigcap_{s \in F} s^{-1}U(s)$ with $U(s) \in \mathcal{U}$ for each $s \in F$, we set

$$\widehat{W} = \prod_{s \in F} U(s) \in \mathcal{U}^F. \text{ Then we can take } \mathcal{V} \text{ to be } \{\widehat{W} : W \in \mathcal{W}\}.$$

Note that $\cup \mathcal{V}$ is an open subset of X^F and $K_F \subset X^F$ is a closed subset, there exists $\delta > 0$ such that $K_{F,\delta} \subset \cup \mathcal{V}$. This finishes the proof. \square

Then, following the ideas of [KL13b, Lemma 5.1] we have:

Proposition 6.4. *Let $\mathcal{U}, \mathcal{U}_2 \in \mathcal{C}_X^o$ and $\mathcal{U}_1 \in \mathcal{C}_X^c$ with $\mathcal{U}_1 \succeq \mathcal{U}_2$. Then*

$$h(G, \mathcal{U}|\mathcal{U}_1) \leq h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) \text{ and thus } h(G, X|\mathcal{U}_1) \leq h^a(G, X|\overline{\mathcal{U}_2}).$$

Proof. Let $\varepsilon > 0$. We choose $1 > \eta > 0$ small enough such that

$$(6.1) \quad \frac{h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + \varepsilon}{1 - \eta} + 2\eta \log |\mathcal{U}| \leq h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + 2\varepsilon$$

and $K \in \mathfrak{F}_G, \delta' > 0$ such that, once $F \in \mathfrak{F}_G$ satisfies $|KF\Delta F| \leq \delta'|F|$ then

$$(6.2) \quad \frac{1}{|F|} \max_{K \in (\overline{\mathcal{U}_2})_F} \log N(\mathcal{U}_F, K) \leq h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + \varepsilon.$$

Observing that $\mathcal{U}_2 \in \mathcal{C}_X^o$ and $\mathcal{U}_1 \in \mathcal{C}_X^c$ satisfy $\mathcal{U}_1 \succeq \mathcal{U}_2$, we can choose $\delta'' > 0$ such that, for each $U_1 \in \mathcal{U}_1$ there exists $U_2 \in \mathcal{U}_2$ containing the open δ'' -neighborhood of U_1 . Now let $l \in \mathbb{N}$ and $\eta' > 0$ be as given by Lemma 6.2 with respect to $\tau = \eta$

and η . In \mathfrak{F}_G we take $e_G \in F_1 \subset \cdots \subset F_l$ such that $|F_{k-1}^{-1}F_k \setminus F_k| \leq \eta'|F_k|$ for each $k = 2, \dots, l$ and $|KF_k \Delta F_k| \leq \delta'|F_k|$ for all $k = 1, \dots, l$. As the group G is amenable, such subsets F_1, \dots, F_l must exist. Thus, by (6.2),

$$(6.3) \quad \max_{k=1}^l \frac{1}{|F_k|} \max_{K \in (\overline{\mathcal{U}_2})_{F_k}} \log N(\mathcal{U}_{F_k}, K) \leq h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + \varepsilon.$$

For each $k = 1, \dots, l$ and any $K \in (\overline{\mathcal{U}_2})_{F_k}$, let $\delta(k, K) > 0$ be as given by Lemma 6.3 with respect to K, F_k and \mathcal{U} , and then set $\delta_k = \min\{\delta(k, K) : K \in (\overline{\mathcal{U}_2})_{F_k}\}$. We take $\delta > 0$ such that $\delta \leq \min\{(\delta'')^2, \delta_1^2, \dots, \delta_l^2, \frac{\eta}{|F_l|}\}$ and

$$(6.4) \quad \sum_{j=0}^{\lfloor |F_l| \delta d \rfloor} \binom{d}{j} < (1 + \varepsilon)^d \text{ for all large enough } d \in \mathbb{N}.$$

Now let $\sigma : G \rightarrow \text{Sym}(d)$ be a good enough sofic approximation for G with some $d \in \mathbb{N}$ (and hence $d \in \mathbb{N}$ is large enough). If $(x_1, \dots, x_d) \in X_{F_l, \delta, \sigma}^d$ then

$$\max_{s \in F_l} \sqrt{\sum_{i=1}^d \frac{1}{d} \rho^2(sx_i, x_{\sigma_s(i)})} < \delta,$$

which implies that $|J(x_1, \dots, x_d, F_l)| \geq (1 - |F_l| \delta)d$, where

$$J(x_1, \dots, x_d, F_l) = \left\{ i \in \{1, \dots, d\} : \max_{s \in F_l} \rho(sx_i, x_{\sigma_s(i)}) < \sqrt{\delta} \right\}.$$

Denote by Θ the set of all subsets of $\{1, \dots, d\}$ with at least $(1 - |F_l| \delta)d$ many elements, and for each $\theta \in \Theta$ denote by $X_{F_l, \delta, \sigma, \theta}^d$ the set of all $(x_1, \dots, x_d) \in X_{F_l, \delta, \sigma}^d$ with $J(x_1, \dots, x_d, F_l) = \theta$. Then

$$(6.5) \quad |\Theta| = \sum_{j=0}^{\lfloor |F_l| \delta d \rfloor} \binom{d}{j} < (1 + \varepsilon)^d \text{ (using (6.4)) and } \bigcup_{\theta \in \Theta} X_{F_l, \delta, \sigma, \theta}^d = X_{F_l, \delta, \sigma}^d.$$

Let $\theta \in \Theta$. As σ is good enough, by Lemma 6.2 there exist $C_1, \dots, C_l \subset \theta$ with

- (1) the sets $\sigma(F_k)C_k, k \in \{1, \dots, l\}$ are pairwise disjoint;
- (2) $\{\sigma(F_k)c : c \in C_k\}$ is η -disjoint for each $k = 1, \dots, l$;
- (3) $\{\sigma(F_k)C_k : k \in \{1, \dots, l\}\}$ $(1 - 2\eta)$ -covers $\{1, \dots, d\}$; and
- (4) for every $k \in \{1, \dots, l\}$ and $c \in C_k$, $F_k \ni s \mapsto \sigma_s(c)$ is bijective.

Set $J_\theta = \{1, \dots, d\} \setminus \bigcup \{\sigma(F_k)C_k : k \in \{1, \dots, l\}\}$. Then

$$(6.6) \quad |J_\theta| \leq 2\eta d \text{ and } \sum_{k=1}^l |F_k| \cdot |C_k| \leq \frac{1}{1 - \eta} \sum_{k=1}^l |\sigma(F_k)C_k| \leq \frac{d}{1 - \eta}.$$

Now let $W \in (\mathcal{U}_1)^d$, say $W = \prod_{i=1}^d U_1^{(i)}$ and for each $i = 1, \dots, d$ the open δ'' -neighborhood of $U_1^{(i)}$ is contained in $U_2^{(i)} \in \mathcal{U}_2$. For each $k = 1, \dots, l$ and any

$c_k \in C_k$, as $C_k \subset \theta$ and $F_k \subset F_l$, if $(x_1, \dots, x_d) \in X_{F_l, \delta, \sigma, \theta}^d \cap W$, then

$$\max_{s \in F_k} \rho(x_{\sigma_s(c_k)}, sx_{c_k}) < \sqrt{\delta} \leq \delta'',$$

and so $sx_{c_k} \in U_2^{(\sigma_s(c_k))}$ (as $x_{\sigma_s(c_k)} \in U_1^{(\sigma_s(c_k))}$) for all $s \in F_k$ by the selection of δ'' , thus

$$x_{c_k} \in \bigcap_{s \in F_k} s^{-1} \overline{U_2^{(\sigma_s(c_k))}} \text{ (denoted by } Q) \in (\overline{\mathcal{U}_2})_{F_k},$$

which implies by applying Lemma 6.3 and (6.3) that we can cover

$$\begin{aligned} & \{(x_i)_{i \in \sigma(F_k)c_k} : (x_1, \dots, x_d) \in X_{F_l, \delta, \sigma, \theta}^d \cap W\} \\ & \subset \left\{ (x_i)_{i \in \sigma(F_k)c_k} : \max_{s \in F_k} \rho(x_{\sigma_s(c_k)}, sx) < \delta_k \text{ for some } x \in Q \right\} \end{aligned}$$

by at most (observing the selection of δ_k)

$$N(\mathcal{U}_{F_k}, Q) \leq e^{|F_k| \cdot [h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + \varepsilon]}$$

elements of $\mathcal{U}^{\sigma(F_k)c_k}$, and so it is not hard to cover

$$\{(x_i)_{i \in \sigma(F_k)c_k} : (x_1, \dots, x_d) \in X_{F_l, \delta, \sigma, \theta}^d \cap W\}$$

using at most

$$e^{|C_k| \cdot |F_k| \cdot [h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + \varepsilon]}$$

elements of $\mathcal{U}^{\sigma(F_k)c_k}$. Thus

$$\begin{aligned} \log N(\mathcal{U}^d, X_{F_l, \delta, \sigma, \theta}^d \cap W) & \leq \sum_{k=1}^l |C_k| \cdot |F_k| \cdot [h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + \varepsilon] + |J_\theta| \log |\mathcal{U}| \\ & \leq d \left(\frac{h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + \varepsilon}{1 - \eta} + 2\eta \log |\mathcal{U}| \right) \text{ (using (6.6))} \\ (6.7) \quad & \leq d(h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + 2\varepsilon) \text{ (using (6.1)).} \end{aligned}$$

Combining (6.5) with (6.7) we obtain

$$\log N(\mathcal{U}^d, X_{F_l, \delta, \sigma}^d \cap W) \leq d(h^a(G, \mathcal{U}|\overline{\mathcal{U}_2}) + 2\varepsilon + \log(1 + \varepsilon)).$$

By the arbitrariness of $\varepsilon > 0$ and $W \in (\mathcal{U}_1)^d$ we obtain the conclusion. \square

We also have [KL13b, Lemma 4.6], an improved version of Lemma 6.2 for an amenable group. Recall that the group G is amenable throughout the whole section.

Lemma 6.5. *Let $0 \leq \tau < 1, 0 < \eta < 1$ and $K \in \mathfrak{F}_G, \delta > 0$. Then there are an $l \in \mathbb{N}$ and $F_1, \dots, F_l \in \mathfrak{F}_G$ with $|KF_k \setminus F_k| < \delta|F_k|$ and $|F_k K \setminus F_k| < \delta|F_k|$ for all $k = 1, \dots, l$, such that for every good enough sofic approximation $\sigma : G \rightarrow \text{Sym}(d)$ for G with some $d \in \mathbb{N}$ and any set $V \subset \{1, \dots, d\}$ with $|V| \geq (1 - \tau)d$, there exist $C_1, \dots, C_l \subset V$ such that*

- (1) *the sets $\sigma(F_k)C_k, k \in \{1, \dots, l\}$ are pairwise disjoint;*

- (2) $\{\sigma(F_k)C_k : k \in \{1, \dots, l\}\}$ $(1 - \tau - \eta)$ -covers $\{1, \dots, d\}$; and
(3) for every $k \in \{1, \dots, l\}$, the map $F_k \times C_k \ni (s, c) \mapsto \sigma_s(c)$ is bijective.

Let $\mathcal{U} \in \mathcal{C}_X$, $\varepsilon > 0$ and $F \in \mathfrak{F}_G$, $\delta > 0$. We set

$$h_{F,\delta}(G, \varepsilon|\mathcal{U}) = \limsup_{i \rightarrow \infty} \frac{1}{d_i} \log \max_{V \in \mathcal{U}^{d_i}} N_\varepsilon(X_{F,\delta,\sigma_i}^{d_i} \cap V, \rho_{d_i}) \quad (\text{recalling (2.1)}),$$

$$h(G, \varepsilon|\mathcal{U}) = \inf_{F \in \mathfrak{F}_G} \inf_{\delta > 0} h_{F,\delta}(G, \varepsilon|\mathcal{U}).$$

Now let $\mathcal{V} \in \mathcal{C}_X$ and $\varepsilon_1, \varepsilon_2 > 0$. Assume that $\text{diam}(\mathcal{V}) < \varepsilon_1$ and any open ball with radius ε_2 is contained in some element of \mathcal{V} . It is easy to obtain

$$h_{F,\delta}(G, \varepsilon_1|\mathcal{U}) \leq h_{F,\delta}(G, \mathcal{V}|\mathcal{U}) \leq h_{F,\delta}(G, \varepsilon_2|\mathcal{U}),$$

$$h(G, \varepsilon_1|\mathcal{U}) \leq h(G, \mathcal{V}|\mathcal{U}) \leq h(G, \varepsilon_2|\mathcal{U}).$$

Thus we have

$$(6.8) \quad h(G, X|\mathcal{U}) = \lim_{\varepsilon \rightarrow 0} h(G, \varepsilon|\mathcal{U}) = \sup_{\varepsilon > 0} h(G, \varepsilon|\mathcal{U}).$$

Following the ideas of [KL13b, Lemma 5.2] we have:

Proposition 6.6. *Let $\mathcal{U} \in \mathcal{C}_X$. Then $h(G, X|\mathcal{U}) \geq h^a(G, X|\mathcal{U})$.*

Proof. Let $\mathcal{U}_1 \in \mathcal{C}_X^o$. We only need to prove $h(G, X|\mathcal{U}) \geq h^a(G, \mathcal{U}_1|\mathcal{U})$.

We choose $\varepsilon > 0$ such that any open ball with ρ -radius ε is contained in some element of \mathcal{U}_1 , and let $\theta > 0$, $F \in \mathfrak{F}_G$, $\delta > 0$. We are to finish the proof by showing

$$(6.9) \quad \frac{1}{d} \log \max_{V \in \mathcal{U}^d} N_\varepsilon(X_{F,\delta,\sigma}^d \cap V, \rho_d) \geq h^a(G, \mathcal{U}_1|\mathcal{U}) - 3\theta$$

once $\sigma : G \rightarrow \text{Sym}(d)$ is a good enough sofic approximation for G with some $d \in \mathbb{N}$.

Let $M > 0$ be large enough and $\delta' > 0$ small enough such that the diameter of the space X is at most M and

$$(6.10) \quad \sqrt{\delta'}M < \frac{\delta}{2}, \delta' \log |\mathcal{U}| < \theta \text{ and } (1 - \delta')h^a(G, \mathcal{U}_1|\mathcal{U}) \geq h^a(G, \mathcal{U}_1|\mathcal{U}) - \theta.$$

Applying Lemma 6.5, there are an $l \in \mathbb{N}$ and $F_1, \dots, F_l \in \mathfrak{F}_G$ so that

$$(6.11) \quad \min_{k=1}^l \min_{s \in F} \frac{|s^{-1}F_k \cap F_k|}{|F_k|} \geq 1 - \delta'$$

and

$$(6.12) \quad \min_{k=1}^l \frac{1}{|F_k|} \max_{K \in \mathcal{U}_{F_k}} \log N((\mathcal{U}_1)_{F_k}, K) \geq \max\{0, h^a(G, \mathcal{U}_1|\mathcal{U}) - \theta\},$$

such that once $\sigma : G \rightarrow \text{Sym}(d)$ is a good enough sofic approximation for G with some $d \in \mathbb{N}$ then there exist $C_1, \dots, C_l \subset \{1, \dots, d\}$ satisfying

- (1) the sets $\sigma(F_k)C_k, k \in \{1, \dots, l\}$ are pairwise disjoint;
(2) $\{\sigma(F_k)C_k : k \in \{1, \dots, l\}\}$ $(1 - \delta')$ -covers $\{1, \dots, d\}$;

- (3) for every $k \in \{1, \dots, l\}$, the map $F_k \times C_k \ni (s, c) \mapsto \sigma_s(c)$ is bijective; and
(4) for all $k \in \{1, \dots, l\}$ and $s \in F$, $s_k \in F_k$, $c_k \in C_k$, $\sigma_{ss_k}(c_k) = \sigma_s \sigma_{s_k}(c_k)$.

Remark again that the group G is amenable, such subsets F_1, \dots, F_l must exist.

Now assume that $\sigma : G \rightarrow \text{Sym}(d)$ is a good enough sofic approximation for G with some $d \in \mathbb{N}$ and let $C_1, \dots, C_l \subset \{1, \dots, d\}$ be constructed as above.

For each $k \in \{1, \dots, l\}$ and any $K \in \mathcal{U}_{F_k}$, we take a maximal $(\rho_{F_k}, \varepsilon)$ -separated subset $E_{k,K}$ of K which is obviously finite, where

$$\rho_{F_k}(x, x') = \max_{g \in F_k} \rho(gx, gx') \text{ for } x, x' \in X.$$

Then for each $y \in K$ there exists an $x \in E_{k,K}$ such that $\rho_{F_k}(x, y) < \varepsilon$. Observe that any open ball with ρ -radius ε is contained in some element of \mathcal{U}_1 , and hence any open ball with ρ_{F_k} -radius ε is contained in some element of $(\mathcal{U}_1)_{F_k}$, thus

$$(6.13) \quad |E_{k,K}| \geq N((\mathcal{U}_1)_{F_k}, K).$$

Now let (y_1, \dots, y_l) be any l -tuple with

$$y_k \in \prod_{c \in C_k} E_{k,K_c}, \text{ where } k \in \{1, \dots, l\} \text{ and } K_c \in \mathcal{U}_{F_k} \text{ for each } c \in C_k.$$

From the construction of C_1, \dots, C_l , it is not hard to see that there exists at least one point $(x_1, \dots, x_d) \in X^d$ such that once $i \in \sigma(F_k)C_k$ for some $k \in \{1, \dots, l\}$, say $i = \sigma_{s_k}(c_k)$ with $s_k \in F_k$ and $c_k \in C_k$, then $x_i = s_k y_k(c_k)$. Let $(x_1, \dots, x_d) \in X^d$ be such a point (corresponding to the l -tuple (y_1, \dots, y_l)). Let $s \in F$ and $i \in \{1, \dots, d\}$. Once $i = \sigma_{s_k}(c_k)$ for some $s_k \in F_k$ and $c_k \in C_k$, $k \in \{1, \dots, l\}$, if $ss_k \in F_k$, then $sx_i = ss_k y_k(c_k) = x_{\sigma_{ss_k}(c_k)} = x_{\sigma_s \sigma_{s_k}(c_k)} = x_{\sigma_s(i)}$. Which implies that

$$(6.14) \quad \frac{1}{d} \sum_{i=1}^d \rho^2(sx_i, x_{\sigma_s(i)}) = \frac{1}{d} \sum_{i \in \{1, \dots, d\} \setminus E} \rho^2(sx_i, x_{\sigma_s(i)}) \leq \frac{M^2}{d} |\{1, \dots, d\} \setminus E|,$$

where

$$E = \bigcup_{k=1}^l \sigma(s^{-1}F_k \cap F_k)C_k.$$

Using the construction of C_1, \dots, C_l again, by (6.11) one has

$$(6.15) \quad |E| = \sum_{k=1}^l |s^{-1}F_k \cap F_k| \cdot |C_k| \geq (1 - \delta') \sum_{k=1}^l |F_k| \cdot |C_k| \geq d(1 - \delta')^2 \geq d(1 - 2\delta').$$

Combining (6.14) with (6.15), we obtain

$$\frac{1}{d} \sum_{i=1}^d \rho^2(sx_i, x_{\sigma_s(i)}) \leq 2\delta' M^2.$$

In particular, $(x_1, \dots, x_d) \in X_{F, \delta, \sigma}^d$ follows from the selection of M and δ' .

On one hand, from the above constructions, it is easy to check that, given $K_c \in \mathcal{U}_{F_k}$ for each $k \in \{1, \dots, l\}$ and any $c \in C_k$, each point $(x_1, \dots, x_d) \in X^d$ corresponding to an l -tuple from $\prod_{k=1}^l \prod_{c \in C_k} E_{k, K_c}$ is contained in a fixed element of

$$\mathcal{W} \doteq \prod_{k=1}^l \mathcal{U}^{\sigma(F_k)C_k} \times \{X\}^{\{1, \dots, d\} \setminus \bigcup_{k=1}^l \sigma(F_k)C_k}.$$

On the other hand, assume that $(x_1, \dots, x_d) \in X^d$ and $(x'_1, \dots, x'_d) \in X^d$ correspond to distinct l -tuples (y_1, \dots, y_l) and (y'_1, \dots, y'_l) from $\prod_{k=1}^l \prod_{c \in C_k} E_{k, K_c}$, where $K_c \in \mathcal{U}_{F_k}$ for each $k \in \{1, \dots, l\}$ and any $c \in C_k$, respectively. Thus, for some $k \in \{1, \dots, l\}$ and $c \in C_k$, $y_k(c)$ and $y'_k(c)$ are distinct elements of E_{k, K_c} , in particular, $\rho_{F_k}(y_k(c), y'_k(c)) \geq \varepsilon$, and then

$$\begin{aligned} \rho_d((x_1, \dots, x_d), (x'_1, \dots, x'_d)) &\geq \max_{i \in \sigma(F_k)c} \rho(x_i, x'_i) \\ &= \max_{s \in F_k} \rho(sy_k(c), sy'_k(c)) = \rho_{F_k}(y_k(c), y'_k(c)) \geq \varepsilon. \end{aligned}$$

This implies that

$$\begin{aligned} \max_{W \in \mathcal{W}} N_\varepsilon(X_{F, \delta, \sigma}^d \cap W, \rho_d) &\geq \left| \prod_{k=1}^l \prod_{c \in C_k} E_{k, K_c} \right| = \prod_{k=1}^l \prod_{c \in C_k} |E_{k, K_c}| \\ &\geq \prod_{k=1}^l \prod_{c \in C_k} N((\mathcal{U}_1)_{F_k}, K_c) \text{ (using (6.13)).} \end{aligned}$$

Combining the above estimation with the fact that $\{\sigma(F_k)C_k : k \in \{1, \dots, l\}\}$ $(1 - \delta')$ -covers $\{1, \dots, d\}$, we obtain that

$$\begin{aligned} &\log \left[|\mathcal{U}|^{\delta' d} \max_{V \in \mathcal{U}^d} N_\varepsilon(X_{F, \delta, \sigma}^d \cap V, \rho_d) \right] \\ &\geq \log \max_{W \in \mathcal{W}} N_\varepsilon(X_{F, \delta, \sigma}^d \cap W, \rho_d) \\ &\geq \sum_{k=1}^l |C_k| \max_{K \in \mathcal{U}_{F_k}} \log N((\mathcal{U}_1)_{F_k}, K) \\ &\geq \sum_{k=1}^l |C_k| \cdot |F_k| \cdot \max\{0, h^a(G, \mathcal{U}_1 | \mathcal{U}) - \theta\} \text{ (using (6.12))} \\ &\geq d(1 - \delta') \max\{0, h^a(G, \mathcal{U}_1 | \mathcal{U}) - \theta\} \\ (6.16) \quad &\geq d[h^a(G, \mathcal{U}_1 | \mathcal{U}) - 2\theta] \text{ (using (6.10)).} \end{aligned}$$

Then (6.9) follows from (6.10) and (6.16). \square

Now let us prove our main result of this section.

Proof of Theorem 6.1. Assume that (X, G) is asymptotically h -expansive. Let $\varepsilon > 0$. Then there exists $\mathcal{U}_1 \in \mathcal{C}_X^o$ such that $h(G, X|\mathcal{U}_1) < \varepsilon$. Thus $h^{a,*}(G, X) \leq h^a(G, X|\mathcal{U}_1) < \varepsilon$ by Proposition 6.6, and finally $h^{a,*}(G, X) = 0$.

Now assume that $h^{a,*}(G, X) = 0$ and let $\varepsilon > 0$. By the definition, there exists $\mathcal{U}_3 \in \mathcal{C}_X^o$ with $h^a(G, X|\overline{\mathcal{U}_3}) < \varepsilon$. As X is a compact metric space, we can take $\mathcal{U}_4 \in \mathcal{C}_X^o$ with $\overline{\mathcal{U}_4} \supseteq \overline{\mathcal{U}_3}$, then $h(G, X|\overline{\mathcal{U}_4}) < \varepsilon$ by Proposition 6.4, and so $h^*(G, X) \leq h(G, X|\overline{\mathcal{U}_4}) < \varepsilon$, thus $h^*(G, X) \leq 0$. That is, (X, G) is asymptotically h -expansive.

We could prove similarly the remaining part of the theorem. \square

Setting $\mathcal{U}_1 = \mathcal{U}_2 = \{X\}$ in Proposition 6.4 and $\mathcal{U} = \{X\}$ in Proposition 6.6, we obtain directly the following observation [KL13b, Theorem 5.3].

Corollary 6.7. $h(G, X) = h^a(G, X)$.

7. ACTIONS OVER COMPACT METRIC GROUPS

In this section we shall provide more interesting asymptotically h -expansive examples when we consider actions of countable discrete amenable groups.

Recall that any C^∞ diffeomorphism on a compact manifold is asymptotically h -expansive by Buzzi [Buz97]. Moreover, if we consider a differentiable action (X, G) in the sense that the homeomorphism of X given by each $g \in G$ is a $C^{(1)}$ map, where X is a compact smooth manifold (here we allow a smooth manifold to have different dimensions for different connected components, even including zero dimension) and G is a countable discrete amenable group containing \mathbb{Z} as a subgroup of infinite index, then the action (X, G) has zero topological entropy by Li and Thom [LT14, Lemma 5.7], and so it is h -expansive.

Moreover, inspired by Misiurewicz's work [Mis76, §7], in the following we prove:

Theorem 7.1. *Let G be a countable discrete amenable group acting on a compact metric group X by continuous automorphisms. Then the action (X, G) is asymptotically h -expansive if and only if $h^a(G, X) < \infty$.*

Remark 7.2. *Let G be a countable discrete amenable group acting on a compact metrizable group X by continuous automorphisms. If (X, G) has finite topological entropy then, by Corollary 3.7, Corollary 6.7 and Theorem 7.1, it admits an invariant measure with maximal entropy. If the amenable group G is infinite and (X, G) has infinite topological entropy, then, by the affinity property of measure-theoretic entropy (for actions of a countable discrete infinite amenable group), it will also admit an invariant measure with maximal entropy. Indeed, if let μ be the normalized Haar measure of the compact metric group X (and hence μ is automatically G -invariant), then the measure-theoretic μ -entropy coincides with the topological entropy of the*

system [Den06, Theorem 2.2]. Remark that X was assumed to be abelian in [Den06, Theorem 2.2], which is in fact not needed.

Remark 7.3. *The conclusion of Theorem 7.1 does not hold if we remove the group structure from X even for the \mathbb{Z} -actions. There are many such actions. The first one may be the Gurevic's example (for the detailed construction see for example [Wal82, Page 192]), which is a \mathbb{Z} -action having finite entropy and without any invariant measure attaining maximal entropy. Another example is [Mis76, Example 6.4], which is a \mathbb{Z} -action with finite entropy such that it is not asymptotically h -expansive, while its each invariant measure has maximal entropy.*

The “only if” part of Theorem 7.1 comes from Corollary 3.3. The proof of its “if” part relies on the concept of homogeneous measures. And so first we recall the definition of homogeneous measures following [Mis76, §7].

Let G be a group acting on a compact metric space X , and denote by α the action. Let $\mu \in M(X)$. For each $\mathcal{U} \in \mathcal{C}_X^o$, we set

$$P(\mathcal{U}) = \bigcup_{x \in X} \left\{ U \in \mathcal{U} : x \in U \text{ and } \mu(U) = \max_{V \in \mathcal{U}, x \in V} \mu(V) \right\} \in \mathcal{C}_X^o.$$

The measure μ is called α -homogeneous if there exist mappings $D : \mathcal{C}_X^o \rightarrow \mathcal{C}_X^o$ and $c : \mathcal{C}_X^o \rightarrow (0, \infty)$ such that for any $\mathcal{U} \in \mathcal{C}_X^o$ and each $F \in \mathcal{F}_G$ we have

$$\mu(V) \leq c(\mathcal{U})\mu(U) \text{ for all } U \in P(\mathcal{U}_F) \text{ and } V \in D(\mathcal{U})_F.$$

In general, it is not easy to check if a measure is homogeneous. While for $G = \mathbb{Z}$ Misiurewicz gave a sufficient condition for the existence of such a measure [Mis76, Theorem 7.2], which can be generalized to a general group G as follows.

Following the spirit of Misiurewicz [Mis76], let H be a group acting on a compact metric space (X, ρ) with the action Φ , that is, Φ is a homomorphism of H into the group of all homeomorphisms of X . Recall that Φ is *transitive* if for any $x, y \in X$ there exists $g \in H$ with $gx = y$, and *equicontinuous* if for each $\varepsilon > 0$ there exists $\delta > 0$ such that $\rho(x_1, x_2) < \delta$ implies $\rho(gx_1, gx_2) < \varepsilon$ for all $x_1, x_2 \in X$ and $g \in H$. Recall that $\mu \in M(X)$ is *invariant with respect to Φ* if $g\mu = \mu$ for each $g \in H$. While the transitivity here is different from the usual one in topological dynamics, the definitions of equicontinuity and invariance of a measure are just as usual.

Lemma 7.4. *Let G be a group acting on a compact metric space X , and denote by α the action. Let H be a group with T , an action of G on H by homomorphisms $g \mapsto T_g$ for each $g \in G$, equivalently, each $T_g : H \rightarrow H$ is a homomorphism. Now assume that Φ is a transitive equicontinuous action of H on X by homeomorphisms such that $\mu \in M(X)$ is invariant with respect to Φ and $g(hx) = (T_g h)(gx)$ for all $x \in X, g \in G$ and $h \in H$. Then μ is α -homogeneous.*

Proof. Let $\mathcal{U} \in \mathcal{C}_X^\circ$. We aim to construct some $D(\mathcal{U}) \in \mathcal{C}_X^\circ$ with $c(\mathcal{U}) = 1$ such that both of mappings D and c satisfy the condition of μ being α -homogeneous.

Let $\varepsilon > 0$ be a Lebesgue number of \mathcal{U} . By the equicontinuity of Φ there exists $\delta > 0$ such that $\rho(x_1, x_2) \leq \delta$ implies $\rho(hx_1, hx_2) < \frac{\varepsilon}{3}$ for all $x_1, x_2 \in X$ and $h \in H$. Now let W be a non-empty open set with diameter at most δ . For each $x \in X$, denote by W_x the set of all $y \in X$ such that both x and y are contained in hW for some $h \in H$, then $\rho(x, y) < \frac{\varepsilon}{3}$ for each $y \in W_x$, and so $W_x \subset U_x$ for some $U_x \in \mathcal{U}$.

Observe that the family of non-empty open subsets $\{hW : h \in H\}$ covers X by the transitivity of the action Φ , and so we could choose $D(\mathcal{U}) \in \mathcal{C}_X^\circ$ with $D(\mathcal{U}) \subset \{hW : h \in H\}$. Now we show that $D(\mathcal{U})$ and $c(\mathcal{U}) = 1$ satisfy the required properties.

Let $F \in \mathcal{F}_G$. Set $W_{F,x} = \bigcap_{\gamma \in F} \gamma^{-1}W_{\gamma x} \subset \bigcap_{\gamma \in F} \gamma^{-1}U_{\gamma x} \in \mathcal{U}_F$ for each $x \in X$. Let $U \in D(\mathcal{U})_F$, and say $U = \bigcap_{g \in F} g^{-1}(h_g W)$ with $h_g \in H$ for any $g \in F$. Then $gx \in h_g W$ and hence $h_g W \subset W_{gx}$ for all $x \in U$ and $g \in F$. Thus, for any $x \in U$,

$$U = \bigcap_{g \in F} g^{-1}(h_g W) \subset \bigcap_{g \in F} g^{-1}W_{gx} = W_{F,x} \ni x$$

and hence (observing $W_{F,x} \subset \bigcap_{\gamma \in F} \gamma^{-1}U_{\gamma x} \in \mathcal{U}_F$)

$$(7.1) \quad \mu(U) \leq \mu(W_{F,x}) \leq \max\{\mu(V) : x \in V \text{ and } V \in \mathcal{U}_F\}.$$

Let $x \in X$ and $k \in H$. For $y \in X$, $y \in kW_x$, equivalently $k^{-1}y \in W_x$, if and only if both x and $k^{-1}y$ are contained in hW for some $h \in H$, equivalently both kx and y are contained in khW for some $h \in H$, if and only if $y \in W_{kx}$. Thus $kW_x = W_{kx}$. Now let $h \in H, Y \subset X$ and $g \in G$. By assumptions, $g(h(g^{-1}Y)) = (T_g h)(g(g^{-1}Y)) = (T_g h)Y$, equivalently, $h(g^{-1}Y) = g^{-1}((T_g h)Y)$. And hence

$$h(g^{-1}W_{gx}) = g^{-1}((T_g h)W_{gx}) = g^{-1}(W_{(T_g h)(gx)}) = g^{-1}(W_{g(hx)}).$$

Which implies that, for all $x \in X$ and any $h \in H, F \in \mathcal{F}_G$,

$$hW_{F,x} = h \left(\bigcap_{\gamma \in F} \gamma^{-1}W_{\gamma x} \right) = \bigcap_{\gamma \in F} h(\gamma^{-1}W_{\gamma x}) = \bigcap_{\gamma \in F} \gamma^{-1}W_{\gamma(hx)} = W_{F,hx}.$$

As the action Φ is transitive and μ is invariant with respect to Φ , we obtain

$$(7.2) \quad \mu(W_{F,x}) = \mu(W_{F,y}) \text{ for all } x, y \in X \text{ and any } F \in \mathcal{F}_G.$$

Summing up, for each $F \in \mathcal{F}_G$ and any $U \in D(\mathcal{U})_F$ and $Q \in P(\mathcal{U}_F)$, let $x \in U$, and by the construction of $P(\mathcal{U}_F)$ we could choose $y \in Q$ with $\mu(Q) = \max\{\mu(V) : y \in V \text{ and } V \in \mathcal{U}_F\}$, and then

$$\begin{aligned} \mu(U) &\leq \mu(W_{F,x}) \text{ (using (7.1))} = \mu(W_{F,y}) \text{ (using (7.2))} \\ &\leq \max\{\mu(V) : y \in V \text{ and } V \in \mathcal{U}_F\} \text{ (using (7.1) again)} = \mu(Q). \end{aligned}$$

Recalling $c(\mathcal{U}) = 1$, we see that the measure μ is α -homogeneous. \square

As a direct corollary, we have:

Proposition 7.5. *Let G be a group acting on a compact metric group X by continuous automorphisms, and denote by α the action. Let $\mu \in M(X)$ be the normalized Haar measure of the group X . Then μ is α -homogeneous.*

Proof. Set $H = X$ and let T be the action α of G on H by automorphisms $T_g : H \rightarrow H, h \mapsto gh$. Now let Φ be the action of H on X by homeomorphisms $h : X \rightarrow X, x \mapsto hx$ for all $h \in X$. Obviously, the action Φ is transitive and the measure μ is invariant with respect to the action Φ and

$$g(hx) = (gh)(gx) = (T_g h)(gx) \text{ for all } x \in X, g \in G \text{ and } h \in H,$$

recalling that G acts on $H = X$ by automorphisms. Observe that by the well-known Birkhoff-Kakutani Theorem the compact metric group X admits a left-invariant compatible metric (for example see [EW11, Page 430, Lemma C.2]), which implies that the action Φ is equicontinuous. Thus the measure μ is α -homogeneous by Lemma 7.4. \square

For a \mathbb{Z} -action acting on a compact metric space X , if the action admits a homogeneous measure and has finite topological entropy, then it is asymptotically h -expansive [Mis76, Theorem 7.1]. In the following, we shall see that the proof there also works for a general countable discrete infinite amenable group G .

Let (X, G) be an action of a countable group G on a compact metric space X with $\mu \in M(X), F \in \mathcal{F}_G, \mathcal{U} \in \mathcal{C}_X^\circ$. We put

$$M_F(\mathcal{U}) = \max_{U \in \mathcal{U}_F} \mu(U) > 0 \text{ and } m_F(\mathcal{U}) = \min_{U \in P(\mathcal{U}_F)} \mu(U).$$

For $\mathcal{U}, \mathcal{V} \in \mathcal{C}_X^\circ$, $M_F(\mathcal{V}) \leq M_F(\mathcal{U})$ whenever $\mathcal{V} \succeq \mathcal{U}$, and if μ is α -homogeneous then

$$(7.3) \quad M_F(\mathcal{V}) \leq M_F(D(\mathcal{U})) \leq c(\mathcal{U})m_F(\mathcal{U}) \text{ and hence } m_F(\mathcal{U}) \geq \frac{M_F(\mathcal{V})}{c(\mathcal{U})} > 0$$

whenever $\mathcal{V} \succeq D(\mathcal{U})$. Let $\mathcal{U} \in \mathcal{C}_X^\circ$ and $Y \subset X$. The *star of Y with respect to \mathcal{U}* and the *star of \mathcal{U}* is defined, respectively, as

$$st(Y, \mathcal{U}) = \cup \{U \in \mathcal{U} : U \cap Y \neq \emptyset\} \text{ and } St\mathcal{U} = \{st(U, \mathcal{U}) : U \in \mathcal{U}\} \in \mathcal{C}_X^\circ.$$

Following [Mis76, Lemma 7.1], it is easy to obtain:

Lemma 7.6. *Let (X, G) be an action of a countable group G on a compact metric space X with $\mu \in M(X)$ and $\mathcal{U}, \mathcal{V} \in \mathcal{C}_X^\circ$ with $\mathcal{V} \succeq \mathcal{U}$. Then*

$$\max_{K \in \mathcal{U}_F} N((St\mathcal{V})_F, K) \cdot m_F(\mathcal{V}) \leq M_F(St\mathcal{U}).$$

Proof. Observe $St(\mathcal{V}_F) \succeq (St\mathcal{V})_F$ as done in [Mis76, Lemma 3.3]. Let $U \in \mathcal{U}_F$ with

$$(7.4) \quad \max_{K \in \mathcal{U}_F} N((St\mathcal{V})_F, K) \leq \max_{K \in \mathcal{U}_F} N(St(\mathcal{V}_F), K) = N(St(\mathcal{V}_F), U) \text{ (denoted by } p).$$

Using [Mis76, Lemma 3.1], we could select a disjoint family $\mathcal{C} \subset P(\mathcal{V}_F)$ such that $C \cap U \neq \emptyset$ for each $C \in \mathcal{C}$ and $U \subset \cup\{st(C, P(\mathcal{V}_F)) : C \in \mathcal{C}\}$. Observing $P(\mathcal{V}_F) \subset \mathcal{V}_F$ by the definition, $\mathcal{C} \subset \mathcal{V}_F$ (and hence $st(C, \mathcal{V}_F) \in St(\mathcal{V}_F)$ for each $C \in \mathcal{C}$) and $U \subset \cup\{st(C, \mathcal{V}_F) : C \in \mathcal{C}\}$, which implies $p \leq |\mathcal{C}|$. Now say $U = \bigcap_{\gamma \in F} \gamma^{-1}U_\gamma$ with $U_\gamma \in \mathcal{U}$ for each $\gamma \in F$, then

$$\begin{aligned}
st(U, \mathcal{V}_F) &= \cup \left\{ W \in \mathcal{V}_F : W \cap \bigcap_{\gamma \in F} \gamma^{-1}U_\gamma \neq \emptyset \right\} \\
&\subset \bigcap_{\gamma \in F} st(\gamma^{-1}U_\gamma, \gamma^{-1}\mathcal{V}) = \bigcap_{\gamma \in F} \gamma^{-1}st(U_\gamma, \mathcal{V}) \\
(7.5) \quad &\subset \bigcap_{\gamma \in F} \gamma^{-1}st(U_\gamma, \mathcal{U}) \text{ (as } \mathcal{V} \succeq \mathcal{U}) \in (St\mathcal{U})_F \text{ (as } U_\gamma \in \mathcal{U} \text{ for all } \gamma \in F).
\end{aligned}$$

As $\mathcal{C} \subset \mathcal{V}_F$ and $C \cap U \neq \emptyset$ for each $C \in \mathcal{C}$, $\cup\mathcal{C} \subset st(U, \mathcal{V}_F)$ by the definition. Now recalling that $\mathcal{C} \subset P(\mathcal{V}_F)$ is a disjoint family and $p \leq |\mathcal{C}|$, we have

$$p \cdot m_F(\mathcal{V}) \leq \mu(\cup\mathcal{C}) \leq \mu(st(U, \mathcal{V}_F)) \leq M_F(St\mathcal{U})$$

by (7.5). Then the conclusion follows from (7.4) and the above estimation. \square

The following result generalizes [Mis76, Theorem 7.1] to the case of G being a general countable discrete infinite amenable group.

Proposition 7.7. *Let α be the action (X, G) with G a countable discrete infinite amenable group and $\mu \in M(X, G)$. Assume that μ is α -homogeneous and $h^a(G, X) < \infty$. Then the action (X, G) is asymptotically h -expansive.*

Proof. As μ is α -homogeneous, let $D : \mathcal{C}_X^\circ \rightarrow \mathcal{C}_X^\circ$ and $c : \mathcal{C}_X^\circ \rightarrow (0, \infty)$ correspond to μ . Setting $\mathcal{U} = \{X\}$ (and hence $St\mathcal{U} = \{X\}$) in Lemma 7.6 we obtain

$$(7.6) \quad N((St\mathcal{V})_F, X) \cdot m_F(\mathcal{V}) \leq 1, \text{ equivalently, } m_F(\mathcal{V}) \leq \frac{1}{N((St\mathcal{V})_F, X)}$$

for each $\mathcal{V} \in \mathcal{C}_X^\circ$. And then once $\mathcal{V}, \mathcal{W} \in \mathcal{C}_X^\circ$ satisfy $\mathcal{W} \succeq D(\mathcal{V})$, $N(\mathcal{W}_F, X)M_F(\mathcal{W}) \geq 1$ and hence combining (7.3) with (7.6) we obtain

$$(7.7) \quad m_F(\mathcal{V}) \geq \frac{1}{N(\mathcal{W}_F, X)c(\mathcal{V})} \text{ and } M_F(\mathcal{W}) \leq \frac{c(\mathcal{V})}{N((St\mathcal{V})_F, X)}.$$

Now fix any $\mathcal{C}, \mathcal{D} \in \mathcal{C}_X^\circ$. We choose $\mathcal{V} \in \mathcal{C}_X^\circ$ with $St\mathcal{V} \succeq \mathcal{C}, \mathcal{A} \in \mathcal{C}_X^\circ$ with $St\mathcal{A} \succeq D(\mathcal{V})$ and $\mathcal{E} \in \mathcal{C}_X^\circ$ with $St\mathcal{E} \succeq \mathcal{D}$ by [Mis76, Proposition 3.5]; and then find $\mathcal{G} \in \mathcal{C}_X^\circ$ with $\bar{\mathcal{G}} \succeq \mathcal{A}, \mathcal{B} \in \mathcal{C}_X^\circ$ with $\mathcal{B} \succeq \mathcal{A}$ and $\mathcal{B} \succeq \mathcal{E}$ (and hence $St\mathcal{B} \succeq \mathcal{D}$) and $\mathcal{W} \in \mathcal{C}_X^\circ$ with $\mathcal{W} \succeq D(\mathcal{B})$. Observing $m_F(\mathcal{B}) > 0$ by (7.3) for any $F \in \mathcal{F}_G$, we have

$$\max_{K \in (\bar{\mathcal{G}})_F} N((St\mathcal{B})_F, K) \leq \max_{K \in \mathcal{A}_F} N((St\mathcal{B})_F, K) \text{ (as } \bar{\mathcal{G}} \succeq \mathcal{A})$$

$$\begin{aligned} &\leq \frac{M_F(St\mathcal{A})}{m_F(\mathcal{B})} \text{ (using Lemma 7.6, as } \mathcal{B} \succeq \mathcal{A}) \\ &\leq c(\mathcal{V}) \cdot c(\mathcal{B}) \cdot \frac{N(\mathcal{W}_F, X)}{N((St\mathcal{V})_F, X)} \text{ (using (7.7)),} \end{aligned}$$

as $St\mathcal{A} \succeq D(\mathcal{V})$ and $\mathcal{W} \succeq D(\mathcal{B})$; and then (recalling that G is a countable discrete infinite amenable group and observing $St\mathcal{V} \succeq \mathcal{C}$ and $St\mathcal{B} \succeq \mathcal{D}$)

$$h^a(G, \mathcal{D}|\overline{\mathcal{G}}) \leq h^a(G, St\mathcal{B}|\overline{\mathcal{G}}) \leq h^a(G, \mathcal{W}) - h^a(G, St\mathcal{V}) \leq h^a(G, X) - h^a(G, \mathcal{C}).$$

By the arbitrariness of $\mathcal{D} \in \mathcal{C}_X^o$, one has

$$h^{a,*}(G, X) \leq h^a(G, X|\overline{\mathcal{G}}) \leq h^a(G, X) - h^a(G, \mathcal{C}),$$

and then by the arbitrariness of $\mathcal{C} \in \mathcal{C}_X^o$ and observing $h^a(G, X) < \infty$, we obtain $h^{a,*}(G, X) = 0$. That is, (X, G) is asymptotically h -expansive by Theorem 6.1. \square

Now we are ready to prove Theorem 7.1.

Proof of Theorem 7.1. The “only if” part comes from Corollary 3.3.

Now we prove the “if” part. If G is finite, it is trivial to see from the definition that $h^a(G, X) < \infty$ implies the finiteness of the group X , then the action (X, G) is expansive and hence asymptotically h -expansive. Now assume that G is infinite. Let $\mu \in M(X)$ be the normalized Haar measure of the compact metric group X . By Proposition 7.5 and Proposition 7.7, we only need to show $\mu \in M(X, G)$, which is obvious by the assumption, as it is well known that each continuous surjective homomorphism of the compact metric group X preserves μ . \square

It is natural to ask the following question which we haven’t solved currently.

Question 7.8. Does Theorem 7.1 hold when the acting group is sofic?

At the end of this section, we present further discussions about actions of countable groups over a compact metrizable abelian group by continuous automorphisms.

Let α be an action of a countable group G acting on a compact metrizable abelian group X by continuous automorphisms. On one hand, by [CL, Theorem 3.1], the action (X, G) is expansive if and only if there exist some $k \in \mathbb{N}$, some left $\mathbb{Z}G$ -submodule J of $(\mathbb{Z}G)^k$, and some $A \in M_k(\mathbb{Z}G)$ being invertible in $M_k(\ell^1(G))$ such that the left $\mathbb{Z}G$ -module \widehat{X} is isomorphic to $(\mathbb{Z}G)^k/J$ and the rows of A are contained in J . On the other hand, suppose that G is amenable and \widehat{X} is a finitely presented left $\mathbb{Z}G$ -module, and write \widehat{X} as $(\mathbb{Z}G)^k/(\mathbb{Z}G)^n A$ for some $k, n \in \mathbb{N}$ and $A \in M_{n \times k}(\mathbb{Z}G)$, then $h^a(G, X) < \infty$ if and only if the additive map $(\mathbb{Z}G)^k \rightarrow (\mathbb{Z}G)^n$ sending a to aA^* is injective [CL, Theorem 4.11], which implies that (X, G) is asymptotically h -expansive by Theorem 7.1. As a consequence, if G is amenable, then for any non-zero divisor f in $\mathbb{Z}G$ which is not invertible in $\ell^1(G)$, the principal algebraic action

α_f , the canonical action of G over $\widehat{\mathbb{Z}G/\mathbb{Z}G}f$, is asymptotically h -expansive but not expansive. See [CL, §3 and §4] for more details.

8. APPENDIX

In this appendix, we will give some comments about Proposition 5.1. We assume again that G is always a countable discrete amenable group.

There are several versions for the well-known Ornstein-Weiss Lemma for amenable groups. Among them, the following version of it (cf. [LW00, Theorem 6.1], see also [HYZ11, Lemma 2.4]) may be used mostly in the literature.

Proposition 8.1. *Let $f : \mathfrak{F}_G \rightarrow \mathbb{R}$ be a function such that $0 \leq f(Eg) = f(E) \leq f(E \cup F) \leq f(E) + f(F)$ for all $E, F \in \mathfrak{F}_G$ and $g \in G$. Then for any Følner sequence $\{F_n : n \in \mathbb{N}\}$ of G the sequence $\left\{ \frac{f(F_n)}{|F_n|} : n \in \mathbb{N} \right\}$ converges and the value of the limit is independent of the selection of the Følner sequence $\{F_n : n \in \mathbb{N}\}$.*

While, as shown by the following remark, it is not clear for us how to define the tail entropy $h^a(G, \mathcal{W}_1 | \mathcal{W}_2)$ with $\mathcal{W}_1, \mathcal{W}_2 \in \mathcal{C}_X$ for actions (X, G) of countable discrete amenable groups using Proposition 8.1.

Remark 8.2. *Let $\mathcal{U}, \mathcal{W} \in \mathcal{C}_X$. It is easy to check that the function $m_{\mathcal{U}, \mathcal{W}} : \mathfrak{F}_G \rightarrow \mathbb{R}$ satisfies $0 \leq m_{\mathcal{U}, \mathcal{W}}(Eg) = m_{\mathcal{U}, \mathcal{W}}(E)$ and $m_{\mathcal{U}, \mathcal{W}}(E \cup F) \leq m_{\mathcal{U}, \mathcal{W}}(E) + m_{\mathcal{U}, \mathcal{W}}(F)$ for all $E, F \in \mathfrak{F}_G$ and $g \in G$; whereas, it may happen that $m_{\mathcal{U}, \mathcal{W}}(E) > m_{\mathcal{U}, \mathcal{W}}(E \cup F)$ for some $E, F \in \mathfrak{F}_G$ even in the case of $\mathcal{U} \in \mathcal{C}_X^o$ and $\mathcal{W} \in \mathcal{C}_X^c$. Thus we could not obtain directly from Proposition 8.1 the convergence of the sequence $\left\{ \frac{1}{|F_n|} m_{\mathcal{U}, \mathcal{W}}(F_n) : n \in \mathbb{N} \right\}$ for a given Følner sequence $\{F_n : n \in \mathbb{N}\}$ of G .*

In the following we shall present such an example. Let $G = \mathbb{Z}_2 \times \mathbb{Z}$. Then G is a countable discrete amenable group with unit $(0, 0)$. We consider the compact metric space $X = \{a, b\}^G$, where G acts naturally on it. Set

$$\mathcal{U} = \{[a]_{(0,0)}, [b]_{(0,0)}\} \in \mathcal{C}_X^o \text{ and } \mathcal{W} = (1, 0)^{-1} \mathcal{U} \in \mathcal{C}_X^c$$

with $[i]_{(0,0)} = \{(x_g)_{g \in G} : x_{(0,0)} = i\}$, $i \in \{a, b\}$. Let $S \in \mathfrak{F}_{\mathbb{Z}}$ and set

$$E = \{(0, s) : s \in S\} \in \mathfrak{F}_G \text{ and } F = \{(1, s) : s \in S\} = (1, 0) \cdot E \in \mathfrak{F}_G.$$

It is straightforward to check that $\mathcal{U}_F = \mathcal{U}_{(1,0) \cdot E} = \mathcal{W}_E$ and $\mathcal{U}_E = \mathcal{W}_F$, and then

$$m_{\mathcal{U}, \mathcal{W}}(E) = m_{\mathcal{U}, \mathcal{W}}(F) = |S| \log 2 > 0 = m_{\mathcal{U}, \mathcal{W}}(E \cup F).$$

In fact, given a Følner sequence $\{F_n : n \in \mathbb{N}\}$ of G , in a previous version of the paper the authors proved convergence of the sequence $\left\{ \frac{1}{|F_n|} m_{\mathcal{W}_1, \mathcal{W}_2}(F_n) : n \in \mathbb{N} \right\}$ only in the case of $\mathcal{W}_1 \in \mathcal{C}_X^o$ and $\mathcal{W}_2 \in \mathcal{C}_X^c$ using another way different from Proposition 5.1 (at that time the authors were unaware of Proposition 5.1), where the value of the limit is independent of the selection of a Følner sequence of the group G . Hanfeng Li

told us that the sequence $\left\{\frac{1}{|F_n|}m_{\mathcal{W}_1, \mathcal{W}_2}(F_n) : n \in \mathbb{N}\right\}$ converges for all $\mathcal{W}_1, \mathcal{W}_2 \in \mathcal{C}_X$, which helped us to find finally Proposition 5.1 in [Gro99b]. Following Gromov’s idea in [Gro99b], it is not hard to prove Proposition 5.1 using the terminology and results by Ornstein and Weiss in [OW87] (see also [RW00, WZ92]), which is small modifications of the proof of Proposition 8.1 presented in [HYZ11].

ACKNOWLEDGEMENTS

N.-P. Chung would like to thank Lewis Bowen for asking and discussing the question “When the actions of sofic groups are expansive, do they admit measures of maximal entropy?” which started this project. The joint work began when N.-P. Chung visited School of Mathematical Sciences, Fudan University on July, 2012. Part of the work was carried out while N.-P. Chung attended Arbeitsgemeinschaft “Limits of Structures” at Oberwolfach on April, 2013.

As indicated in the section of Appendix, both authors thank Hanfeng Li for many helpful suggestions which resulted in substantial improvements to this paper. The authors are also grateful to Tomasz Downarowicz, Ben Hayes and Wen Huang for helpful comments.

This research was supported through the programme “Research in Pairs” by the Mathematisches Forschungsinstitut Oberwolfach in 2013. We are grateful to MFO for a warm hospitality. N.-P. Chung is also supported by Max Planck Society, and G. H. Zhang is also supported by FANEDD (201018) and NSFC (11271078).

REFERENCES

- [AE12] Miklós Abért and Gábor Elek, *Dynamical properties of profinite actions*, Ergodic Theory Dynam. Systems **32** (2012), no. 6, 1805–1835. MR 2995875
- [AN12] Miklós Abért and Nikolay Nikolov, *Rank gradient, cost of groups and the rank versus Heegaard genus problem*, J. Eur. Math. Soc. (JEMS) **14** (2012), no. 5, 1657–1677. MR 2966663
- [Aus88] Joseph Auslander, *Minimal flows and their extensions*, North-Holland Mathematics Studies, vol. 153, North-Holland Publishing Co., Amsterdam, 1988, Notas de Matemática [Mathematical Notes], 122. MR 956049 (89m:54050)
- [BD04] Mike Boyle and Tomasz Downarowicz, *The entropy theory of symbolic extensions*, Invent. Math. **156** (2004), no. 1, 119–161. MR 2047659 (2005d:37015)
- [BL97] Mike Boyle and Douglas Lind, *Expansive subdynamics*, Trans. Amer. Math. Soc. **349** (1997), no. 1, 55–102. MR 1355295 (97d:58115)
- [Bow72] Rufus Bowen, *Entropy-expansive maps*, Trans. Amer. Math. Soc. **164** (1972), 323–331. MR 0285689 (44 #2907)

- [Bow10] Lewis Bowen, *Measure conjugacy invariants for actions of countable sofic groups*, J. Amer. Math. Soc. **23** (2010), no. 1, 217–245. MR 2552252 (2011b:37010)
- [Bow11] ———, *Entropy for expansive algebraic actions of residually finite groups*, Ergodic Theory Dynam. Systems **31** (2011), no. 3, 703–718. MR 2794944 (2012j:37009)
- [Bow12] ———, *Sofic entropy and amenable groups*, Ergodic Theory Dynam. Systems **32** (2012), no. 2, 427–466. MR 2901354
- [Bur11] David Burguet, *C^2 surface diffeomorphisms have symbolic extensions*, Invent. Math. **186** (2011), no. 1, 191–236. MR 2836054 (2012k:37049)
- [Buz97] Jérôme Buzzi, *Intrinsic ergodicity of smooth interval maps*, Israel J. Math. **100** (1997), 125–161. MR 1469107 (99g:58071)
- [Chu13] Nhan-Phu Chung, *Topological pressure and the variational principle for actions of sofic groups*, Ergodic Theory Dynam. Systems **33** (2013), 1363–1390.
- [CL] Nhan-Phu Chung and Hanfeng Li, *Homoclinic group, IE group and expansive algebraic actions*, arXiv:1103.1567, preprint.
- [Den06] Christopher Deninger, *Fuglede-Kadison determinants and entropy for actions of discrete amenable groups*, J. Amer. Math. Soc. **19** (2006), no. 3, 737–758 (electronic). MR 2220105 (2007b:37010)
- [DFPV12] Lorenzo J. Díaz, Todd Fisher, Maria José Pacifico, and José L. Vieitez, *Entropy-expansiveness for partially hyperbolic diffeomorphisms*, Discrete Contin. Dyn. Syst. **32** (2012), no. 12, 4195–4207. MR 2966742
- [DM09] Tomasz Downarowicz and Alejandro Maass, *Smooth interval maps have symbolic extensions: the antarctic theorem*, Invent. Math. **176** (2009), no. 3, 617–636. MR 2501298 (2010d:37038)
- [DN05] Tomasz Downarowicz and Sheldon Newhouse, *Symbolic extensions and smooth dynamical systems*, Invent. Math. **160** (2005), no. 3, 453–499. MR 2178700 (2006j:37021)
- [Dow11] Tomasz Downarowicz, *Entropy in dynamical systems*, New Mathematical Monographs, vol. 18, Cambridge University Press, Cambridge, 2011.
- [DS02] Tomasz Downarowicz and Jacek Serafin, *Fiber entropy and conditional variational principles in compact non-metrizable spaces*, Fund. Math. **172** (2002), no. 3, 217–247. MR 1898686 (2003b:37027)
- [DS07] Christopher Deninger and Klaus Schmidt, *Expansive algebraic actions of discrete residually finite amenable groups and their entropy*, Ergodic Theory Dynam. Systems **27** (2007), no. 3, 769–786. MR 2322178 (2008d:37009)
- [DZ] Anthony H. Dooley and Guohua Zhang, *Local entropy theory of a random dynamical system*, Mem. Amer. Math. Soc., to appear.

- [ES02] Manfred Einsiedler and Klaus Schmidt, *The adjoint action of an expansive algebraic \mathbb{Z}^d -action*, *Monatsh. Math.* **135** (2002), no. 3, 203–220. MR 1897576 (2003b:37003)
- [EW11] Manfred Einsiedler and Thomas Ward, *Ergodic theory with a view toward-
s number theory*, *Graduate Texts in Mathematics*, vol. 259, Springer-Verlag London Ltd., London, 2011. MR 2723325 (2012d:37016)
- [Gro99a] Mikhael Gromov, *Endomorphisms of symbolic algebraic varieties*, *J. Eur. Math. Soc. (JEMS)* **1** (1999), no. 2, 109–197. MR 1694588 (2000f:14003)
- [Gro99b] ———, *Topological invariants of dynamical systems and spaces of holomorphic maps. I*, *Math. Phys. Anal. Geom.* **2** (1999), no. 4, 323–415. MR 1742309 (2001j:37037)
- [HYZ10] Wen Huang, Xiangdong Ye, and Guohua Zhang, *Lowering topological entropy over subsets*, *Ergodic Theory Dynam. Systems* **30** (2010), no. 1, 181–209. MR 2586351 (2011f:37027)
- [HYZ11] ———, *Local entropy theory for a countable discrete amenable group action*, *J. Funct. Anal.* **261** (2011), no. 4, 1028–1082. MR 2803841
- [Ioa11] Adrian Ioana, *Cocycle superrigidity for profinite actions of property (T) groups*, *Duke Math. J.* **157** (2011), no. 2, 337–367. MR 2783933 (2012c:37006)
- [Ker13] David Kerr, *Sofic measure entropy via finite partitions*, *Groups Geom. Dyn.* **7** (2013), no. 3, 617–632. MR 3095712
- [KL11] David Kerr and Hanfeng Li, *Entropy and the variational principle for actions of sofic groups*, *Invent. Math.* **186** (2011), no. 3, 501–558. MR 2854085
- [KL13a] ———, *Combinatorial independence and sofic entropy*, *Comm. Math. Stat.* **1** (2013), no. 2, 213–257.
- [KL13b] ———, *Soficity, amenability, and dynamical entropy*, *Amer. J. Math.* **135** (2013), no. 3, 721–761. MR 3068400
- [Li13] Hanfeng Li, *Sofic mean dimension*, *Adv. Math.* **244** (2013), 570–604. MR 3077882
- [LS99] Douglas Lind and Klaus Schmidt, *Homoclinic points of algebraic \mathbb{Z}^d -actions*, *J. Amer. Math. Soc.* **12** (1999), no. 4, 953–980. MR 1678035 (2000d:37002)
- [LT14] Hanfeng Li and Andreas Thom, *Entropy, determinants, and L^2 -torsion*, *J. Amer. Math. Soc.* **27** (2014), no. 1, 239–292.
- [LVY] Gang Liao, Marcelo Viana, and Jiagang Yang, *Entropy conjecture for diffeomorphisms away from tangencies*, *J. Eur. Math. Soc. (JEMS)*, to appear.
- [LW00] Elon Lindenstrauss and Benjamin Weiss, *Mean topological dimension*, *Israel J. Math.* **115** (2000), 1–24. MR 1749670 (2000m:37018)
- [Mañ87] Ricardo Mañé, *Ergodic theory and differentiable dynamics*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*, vol. 8, Springer-Verlag, Berlin, 1987, Translated from the Portuguese by Silvio Levy. MR 889254 (88c:58040)

- [Mis76] Michał Misiurewicz, *Topological conditional entropy*, *Studia Math.* **55** (1976), no. 2, 175–200. MR 0415587 (54 #3672)
- [OP10] Narutaka Ozawa and Sorin Popa, *On a class of II_1 factors with at most one Cartan subalgebra*, *Ann. of Math. (2)* **172** (2010), no. 1, 713–749. MR 2680430 (2011j:46101)
- [OW87] Donald S. Ornstein and Benjamin Weiss, *Entropy and isomorphism theorems for actions of amenable groups*, *J. Analyse Math.* **48** (1987), 1–141. MR 910005 (88j:28014)
- [RW00] Daniel J. Rudolph and Benjamin Weiss, *Entropy and mixing for amenable group actions*, *Ann. of Math. (2)* **151** (2000), no. 3, 1119–1150. MR 1779565 (2001g:37001)
- [Sch90] Klaus Schmidt, *Automorphisms of compact abelian groups and affine varieties*, *Proc. London Math. Soc. (3)* **61** (1990), no. 3, 480–496. MR 1069512 (91j:28015)
- [Sch95] ———, *Dynamical systems of algebraic origin*, *Progress in Mathematics*, vol. 128, Birkhäuser Verlag, Basel, 1995. MR 1345152 (97c:28041)
- [Wal82] Peter Walters, *An introduction to ergodic theory*, *Graduate Texts in Mathematics*, vol. 79, Springer-Verlag, New York, 1982. MR 648108 (84e:28017)
- [Wei00] Benjamin Weiss, *Sofic groups and dynamical systems*, *Sankhyā Ser. A* **62** (2000), no. 3, 350–359, *Ergodic theory and harmonic analysis (Mumbai, 1999)*. MR 1803462 (2001j:37022)
- [Wei01] ———, *Monotileable amenable groups*, *Topology, ergodic theory, real algebraic geometry*, *Amer. Math. Soc. Transl. Ser. 2*, vol. 202, Amer. Math. Soc., Providence, RI, 2001, pp. 257–262. MR 1819193 (2001m:22014)
- [Wei03] ———, *Actions of amenable groups*, *Topics in dynamics and ergodic theory*, *London Math. Soc. Lecture Note Ser.*, vol. 310, Cambridge Univ. Press, Cambridge, 2003, pp. 226–262. MR 2052281 (2005d:37008)
- [WZ92] Thomas Ward and Qing Zhang, *The Abramov-Rokhlin entropy addition formula for amenable group actions*, *Monatsh. Math.* **114** (1992), no. 3-4, 317–329. MR 1203977 (93m:28023)
- [Zha12] Guohua Zhang, *Local variational principle concerning entropy of a sofic group action*, *J. Funct. Anal.* **262** (2012), no. 4, 1954–1985. MR 2873866

NHAN-PHU CHUNG, MAX PLANCK INSTITUTE-MATHEMATICS IN THE SCIENCES, INSELSTRASSE 22, 04103 LEIPZIG, GERMANY

E-mail address: chung@mis.mpg.de

GUOHUA ZHANG, SCHOOL OF MATHEMATICAL SCIENCES AND LMNS, FUDAN UNIVERSITY, SHANGHAI 200433, CHINA

E-mail address: chiaths.zhang@gmail.com