

MATHEMATISCHES FORSCHUNGSINSTITUT OBERWOLFACH

Report No. 44/2022

DOI: 10.4171/OWR/2022/44

## Mini-Workshop: Zero-Range and Point-Like Singular Perturbations: For a Spillover to Analysis, PDE and Differential Geometry

Organized by  
Vladimir Georgiev, Pisa  
Alessandro Michelangeli, Bonn/Trieste

2 October – 8 October 2022

**ABSTRACT.** The field of contact interactions and perturbations of differential operators supported on subsets with non-trivial co-dimension is an increasingly active mainstream of mathematical physics (in particular, operator and spectral theory and quantum mechanics), with intimately related applications and mathematical challenges in partial differential equations, and neighbouring sectors of analysis, PDEs, and differential geometry. This Mini-Workshop fostered intense and prolific discussions on recent advances and trends in the field.

*Mathematics Subject Classification (2020):* 35B40, 35J75, 35Q55, 47N20, 53B21, 81Q35.

### Introduction by the Organizers

The Oberwolfach Mini-Workshop 2240a, *Zero-Range and Point-Like Singular Perturbations: For a Spillover to Analysis, PDE and Differential Geometry*, organised by Vladimir Georgiev (Pisa) and Alessandro Michelangeli (Bonn) took place on 2-8 October 2022 in hybrid on-site and online form, and was structured in seminar presentations and an amount of round table discussions. The 13 speakers, of which 11 on-site and 2 online, came from universities from France, Germany, Italy, Portugal, Serbia, and Spain.

This meeting was aimed at promoting and facilitating the transfer of techniques, tools, and open problems from the field of **contact interactions** and **perturbations of differential operators supported on subsets with non-trivial co-dimension**, an increasingly active mainstream of mathematical physics (in particular, operator and spectral theory and quantum mechanics), to intimately

related applications and mathematical challenges in the area of partial differential equations, and neighbouring sectors of analysis, PDEs, and differential geometry. The latter areas are naturally involved when one investigates the heat or Schrödinger evolution flow generated by singular-perturbed Hamiltonian of relevance, the associated standing waves, or when one transposes the problem in the language of stochastic processes generated by such operators, or again when point-like singular-perturbed PDE's are placed on relevant manifold (typically, almost-Riemannian structures).

Inter-particle interactions of virtually zero range ('contact interactions') became a natural tool when in the 1930's quantum mechanics began to be applied to the newly observed nuclear phenomena, as the decrease by a factor  $10^{-5}$  from the atomic to the nuclear scale made it plausible to model the interaction among nucleons as a delta-like interaction. In 1932 Wigner [77] calculated that the nuclear forces must be of very short range and very strong magnitude, leading first Bethe and Peierls [21, 22], then Thomas [76], Fermi, [42], and Breit [28] to describe the neutron-proton scattering, and then some 20 years later, in 1955, Ter-Martirosyan and Skornyakov [74] to describe the three-body problem with zero-range interaction, by means of the Schrödinger equation in the approximation of a two-body potential of very short range, virtually null. Whereas nuclear physics obviously developed into a much finer corpus of knowledge, the 'delta-like' idealisation remained for some decades an efficient tool for a formal first-order perturbation theory in application to atomic physics [40]. Modern advances in the manipulation of cold atoms have today conferred the zero-range modelling a renewed topicality, owing to the experimental possibility of tuning the interaction's effective scattering length to very large values and the interaction's effective range to zero, by means of a magnetically induced Feshbach resonance [71, Section 5.4.2]. Ultra-cold gases in the so-called 'unitary regime' (infinite scattering length and zero-range) are nowadays well preparable and intensively studied (see [30, 27, 69] and references therein).

An intense mathematical activity, at the crossroads of functional analysis, operator theory, and spectral theory paralleled the above physical developments, often being triggered by the latter, until the present days. This includes: the rigorous constructions of self-adjoint Hamiltonians for two-, three-, and  $N$ -body quantum systems with zero-range interactions [18, 65, 75, 39, 32, 61, 62, 63, 33, 60, 16, 58, 59, 67, 68, 15, 56], the study of their spectral properties [64, 8, 66, 62, 63, 79, 17, 56], the approximation by means of ordinary Schrödinger operators with potentials supported at very short scales [8, 9, 13, 50, 14], among other themes, with tools and techniques from spectral and scattering theory, self-adjoint extension schemes, and the like.

Notably, each of the above subjects has witnessed and is witnessing both fabulous, technically ingenious breakthroughs, and almost desperate, deep open problems. And most importantly, for the purposes of the Mini-Workshop, many mathematical questions that are central in the math-phys field sketched above naturally involves specific techniques from, and naturally give rise independent problems in

the realm of neighbouring subjects such as the theory of partial differential equations, differential geometry, and stochastic processes, where several studies have been already initiated.

Let us survey two main representative lines around which the workshop's discussions unfolded.

**Subject line I – Linear and non-linear Schrödinger and heat equations with point-like Laplace perturbation.** This is a natural and active continuation, in the field of PDE, of the quantum mechanical problem of the *time evolution* of particle systems subject to mutual zero-range interaction. The prototypical object of interest is the singular-perturbed Laplacian  $-\Delta_\alpha$  on  $L^2(\mathbb{R}^d)$ ,  $d \in \{1, 2, 3\}$ , which is a self-adjoint extension of  $-\Delta|_{C_0^\infty(\mathbb{R}^d \setminus \{0\})}$ , namely of the free Laplacian initially restricted to smooth functions that vanish in the vicinity of the origin. The parameter  $\alpha \in \mathbb{R} \cup \{\infty\}$  measures, through a suitable expression, the scattering length of the quantum particle that feels the non-trivial interaction supported at  $x = 0$  (or, re-interpreting the  $x$ -variable as a relative variable, the scattering length in the two-body interaction of zero range). For example, in  $d = 3$  dimensions,

$$(1) \quad \begin{aligned} \text{dom}(-\Delta_\alpha) &= \left\{ u \in L^2(\mathbb{R}^3) \mid u = f + \frac{f(0)}{4\pi\alpha + 1} \frac{e^{-|x|}}{|x|} \text{ with } f \in H^2(\mathbb{R}^3) \right\}, \\ (-\Delta_\alpha + 1)u &= (-\Delta + 1)f \end{aligned}$$

( $\text{dom} \equiv \text{domain}$ ). The realisation with  $\alpha = \infty$  is nothing but the self-adjoint Laplacian on the  $H^2$ -domain. This naturally leads to considering linear and non-linear Schrödinger equations

$$(2) \quad i\partial_t u = -\Delta_\alpha u,$$

$$(3) \quad i\partial_t u = -\Delta_\alpha u + g(u),$$

for non-linearities of physical relevance such as

$$(4) \quad g(u) = \lambda|u|^p u, \quad g(u) = (w * |u|^2)u$$

(for suitable  $\lambda, p > 0$  and  $w : \mathbb{R}^3 \rightarrow \mathbb{R}$ ). For the linear problem, in all dimensions, the explicit propagator is known [7, 36], and dispersive and Strichartz estimates [41, 51, 38], as well as the completeness of the wave operators [38, 31, 78] have been recently proved. In dimension  $d = 1$  the point-like nature of the singularity has the explicit structure  $-\Delta + \delta(x)$  (in the sense of sum of quadratic form, where  $\delta(x)$  is the Dirac distribution), thereby allowing for an accurate analysis of (2)-(3) in terms of local and global well-posedness, blow-up, scattering, asymptotic stability, solitons, standing waves, ground state [5, 37, 2, 4, 3, 10, 12, 53, 52, 34, 11, 54, 55, 35]. The same questions in the case of *dimension two or three* encounter a much more uncharted territory, with very recent characterisations [49, 29] of the adapted (i.e., singular-perturbed) fractional Sobolev spaces

$$(5) \quad H_\alpha^s(\mathbb{R}^3) := \text{dom}((-\Delta_\alpha + \lambda \mathbb{1})^{\frac{s}{2}}), \quad s > 0,$$

for  $\lambda > 0$  large enough, the proof of local and global well-posedness for certain three-dimensional [5, 29] and two-dimensional [1, 44] singular-perturbed NLS. This

leaves an ample room open for improvements and unanswered questions concerning solution theory, standing waves, scattering, stability and instability, etc., when  $d = 2, 3$ . Part of the workshop focussed on this, as well as on the counterpart issues for the linear and non-linear heat equation with point-like perturbation.

**Subject line II – Dispersive and scattering properties of the Schrödinger and heat flow on degenerate Riemannian manifold of Grushin type.** This is another natural and active spin-off, in the field of Riemannian geometry and analysis of PDEs on manifold, of the problem of a quantum particle subject to the zero-range interaction supported in a region of non-trivial co-dimension, which is in fact a region where the metric blows up. The study of a quantum particle on degenerate Riemannian manifolds, and the problem of the purely geometric confinement away from the singularity locus of the metric, as opposite to the dynamical transmission across the singularity, has recently attracted a considerable amount of attention in relation to Grushin structures and to the induced confining effective potentials on cylinder, cone, and plane (as in the works [70, 23, 26, 73, 25, 43, 46, 24, 72, 47, 20, 45, 48]), as well as, more generally, on two-step two-dimensional almost-Riemannian structures [23, 20, 19], or also generalisations to almost-Riemannian structures in any dimension and of any step, and even to sub-Riemannian geometries, provided that certain geometrical assumptions on the singular set are taken [43, 73]. On a related note, a satisfactory interpretation of the heat-confinement in the Grushin cylinder is known in terms of Brownian motions [24] and random walks [6]. The prototypical playground is the Grushin-type manifold  $M_\alpha \equiv (M, g_\alpha)$ , for given  $\alpha \in \mathbb{R}$ , with

$$(6) \quad \begin{aligned} M &= \mathbb{R}_x^- \times \mathbb{R}_y + \mathbb{R}_x^+ \times \mathbb{R}_y && \text{(Grushin plane),} \\ M &= \mathbb{R}_x^- \times \mathbb{S}_y^1 + \mathbb{R}_x^+ \times \mathbb{S}_y^1 && \text{(Grushin cylinder)} \end{aligned}$$

and with metric

$$(7) \quad g_\alpha = dx \otimes dx + |x|^{-2\alpha} dy \otimes dy,$$

thus with associated Laplace-Beltrami operator

$$(8) \quad \Delta_\alpha = \frac{\partial^2}{\partial x^2} + |x|^{2\alpha} \frac{\partial^2}{\partial y^2} - \frac{\alpha}{|x|} \frac{\partial}{\partial x}.$$

For such models, the *geometric quantum confinement* in each half-cylinder corresponds to the essential self-adjointness of  $-\Delta_\alpha$  in  $L^2(M, |x|^{-\alpha} dx \wedge dy)$  on its minimal domain of smooth functions supported away from the singularity; the *quantum transmission* between the two halves of  $M$  corresponds instead to the lack of essential self-adjointness, in which case the type of transmission is governed by the Schrödinger equation induced by a self-adjoint extension of  $-\Delta_\alpha$ . We underline the fundamental novelty of the quantum setting, where depending on the boundary conditions for  $\Delta_\alpha$  the particle may or may not trespass the singularity locus, whereas instead classically  $M_\alpha$  is always geodesically incomplete (the classical particle reaches the boundary in finite time). Next to the recently established identification of the  $\alpha$ -regimes of essential self-adjointness or lack-of,

the classification of the self-adjoint extensions of  $-\Delta_\alpha$ , and the study of their spectral and scattering properties [46, 72, 47, 45], the relevant problem arises of the dispersive and scattering properties of the Schrödinger and heat flow on  $M_\alpha$ . For concreteness in the case of the Grushin cylinder (compact  $y$ -variable), one can see [45, 48] that in each Fourier mode conjugate to  $y$  one is led to the study of PDEs of the form

$$(9) \quad \begin{cases} i\partial_t u = -\partial_x^2 u + \frac{\alpha(\alpha+2)}{4x^2} u, \\ u_0(t)^- = u_0(t)^+ \quad \text{with } u_0^\pm(t) := \lim_{x \rightarrow 0^\pm} |x|^{\frac{\alpha}{2}} u(t, x), \\ u_1(t)^- = -u_1(t)^+ \quad \text{with } u_1^\pm(t) := \lim_{x \rightarrow 0^\pm} |x|^{-(1+\frac{\alpha}{2})} (u(t, x) - |x|^{-\frac{\alpha}{2}} u_0^\pm(t)), \\ u(0, x) = \varphi(x) \quad \text{with } \varphi \in L^2(\mathbb{R}) \end{cases}$$

in the unknown  $u \equiv u(t, x)$ , with  $t \geq 0$  and  $x \in \mathbb{R} \setminus \{0\}$ : a free Schrödinger equation except for the non-trivial *boundary conditions of contact interaction* at  $x = 0$  (a signature of a non-trivial left $\leftrightarrow$ right transmission protocol). Similar considerations hold for the associated heat equation. Along a completely open programme recently proposed and advertised in [48], the next crucial challenges are the study of dispersive, smoothing, and Strichartz estimates for linear problems of the type (9), as well as the well-posedness, ground states, scattering, stability/instability of the associated non-linear equations.

The Mini-Workshop aimed at gathering a small number of mathematicians of various degrees of expertise, with an amount of direct or indirect scientific links already transversely present among them, and who are carriers of the mathematical-physical, differential-geometric, functional-analytic, and differential equation theoretic backgrounds, with the general goal of discussing the setup and the perspectives of ongoing and future investigations in each of the above fields, all stemming from, or intimately motivated by the grand picture of zero-range and contact interactions sketched in the previous Section. At the end of a week of mutual and profitable exchanges, this little group of already mildly connected researchers was meant to import into their own network of collaborations the corpus of general motivations, cross-disciplinary stimuli, technical tools, and bibliographic references discussed in the workshop. In this respect, the event set the basis of future larger meetings among neighbouring mathematical communities.

#### REFERENCES

- [1] R. ADAMI, F. BONI, R. CARLONE, AND L. TENTARELLI, *Ground states for the planar NLSE with a point defect as minimisers of the constrained energy*, arXiv:2109.09482 (2021).
- [2] R. ADAMI AND D. NOJA, *Existence of dynamics for a 1D NLS equation perturbed with a generalized point defect*, J. Phys. A, 42 (2009), pp. 495302, 19.
- [3] R. ADAMI AND D. NOJA, *Exactly solvable models and bifurcations: the case of the cubic NLS with a  $\delta$  or a  $\delta'$  interaction in dimension one*, Math. Model. Nat. Phenom., 9 (2014), pp. 1–16.
- [4] R. ADAMI, D. NOJA, AND N. VISCIGLIA, *Constrained energy minimisation and ground states for NLS with point defects*, Discrete Contin. Dyn. Syst. Ser. B, 18 (2013), pp. 1155–1188.
- [5] R. ADAMI AND A. SACCHETTI, *The transition from diffusion to blow-up for a nonlinear Schrödinger equation in dimension 1*, J. Phys. A, 38 (2005), pp. 8379–8392.

- [6] A. AGRACHEV, U. BOSCAIN, R. NEEL, AND L. RIZZI, *Intrinsic random walks in Riemannian and sub-Riemannian geometry via volume sampling*, ESAIM Control Optim. Calc. Var., 24 (2018), pp. 1075–1105.
- [7] S. ALBEVERIO, Z. BRZEŃNIAK, AND L. DABROWSKI, *Fundamental solution of the heat and Schrödinger equations with point interaction*, J. Funct. Anal., 130 (1995), pp. 220–254.
- [8] S. ALBEVERIO AND R. HØEGH-KROHN, *Point interactions as limits of short range interactions*, J. Operator Theory, 6 (1981), pp. 313–339.
- [9] S. ALBEVERIO, R. HØEGH-KROHN, AND T. T. WU, *A class of exactly solvable three-body quantum mechanical problems and the universal low energy behavior*, Physics Letters A, 83 (1981), pp. 105–109.
- [10] J. ANGULO PAVA AND L. C. F. FERREIRA, *On the Schrödinger equation with singular potentials*, Differential Integral Equations, 27 (2014), pp. 767–800.
- [11] J. ANGULO PAVA AND C. A. HERNÁNDEZ MELO, *On stability properties of the cubic-quintic Schrödinger equation with  $\delta$ -point interaction*, Commun. Pure Appl. Anal., 18 (2019), pp. 2093–2116.
- [12] V. BANICA AND N. VISCIGLIA, *Scattering for NLS with a delta potential*, J. Differential Equations, 260 (2016), pp. 4410–4439.
- [13] G. BASTI, C. CACCIAPUOTI, D. FINCO, AND A. TETA, *The three-body problem in dimension one: from short-range to contact interactions*, J. Math. Phys., 59 (2018), pp. 072104, 18.
- [14] ———, *Three-body Hamiltonian with regularized zero-range interactions in dimension three*, arXiv:2107.07188 (2022).
- [15] G. BASTI, R. FIGARI, AND A. TETA, *Regularized quadratic forms for a three boson system with zero-range interactions*, Rend. Mat. Appl. (7), 39 (2018), pp. 205–216.
- [16] G. BASTI AND A. TETA, *On the quantum mechanical three-body problem with zero-range interactions*, in Functional analysis and operator theory for quantum physics, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2017, pp. 71–93.
- [17] S. BECKER, A. MICHELANGELI, AND A. OTTOLINI, *Spectral Analysis of the 2+1 Fermionic Trimer with Contact Interactions*, Mathematical Physics, Analysis and Geometry, 21 (2018), p. 35.
- [18] F. BEREZIN AND L. FADDEEV, *A Remark on Schrodinger’s equation with a singular potential*, Sov.Math.Dokl., 2 (1961), pp. 372–375.
- [19] I. BESCHASTNYI, *Closure of the Laplace-Beltrami operator on 2D almost-Riemannian manifolds and semi-Fredholm properties of differential operators on Lie manifolds*, arXiv:2104.07745 (2021).
- [20] I. BESCHASTNYI, U. BOSCAIN, AND E. POZZOLI, *Quantum Confinement for the Curvature Laplacian  $-\Delta + cK$  on 2D-Almost-Riemannian Manifolds*, Potential Analysis, (2021).
- [21] H. BETHE AND R. PEIERLS, *Quantum Theory of the Dipion*, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences, 148 (1935), pp. 146–156.
- [22] H. A. BETHE AND R. PEIERLS, *The Scattering of Neutrons by Protons*, Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences, 149 (1935), pp. 176–183.
- [23] U. BOSCAIN AND C. LAURENT, *The Laplace-Beltrami operator in almost-Riemannian geometry*, Ann. Inst. Fourier (Grenoble), 63 (2013), pp. 1739–1770.
- [24] U. BOSCAIN AND R. W. NEEL, *Extensions of Brownian motion to a family of Grushin-type singularities*, Electron. Commun. Probab., 25 (2020), pp. Paper No. 29, 12.
- [25] U. BOSCAIN AND D. PRANDI, *Self-adjoint extensions and stochastic completeness of the Laplace-Beltrami operator on conic and anticonic surfaces*, J. Differential Equations, 260 (2016), pp. 3234–3269.
- [26] U. BOSCAIN, D. PRANDI, AND M. SERI, *Spectral analysis and the Aharonov-Bohm effect on certain almost-Riemannian manifolds*, Comm. Partial Differential Equations, 41 (2016), pp. 32–50.
- [27] E. BRAATEN AND H.-W. HAMMER, *Universality in few-body systems with large scattering length*, Physics Reports, 428 (2006), pp. 259–390.

- [28] G. BREIT, *The scattering of slow neutrons by bound protons. I. Methods of calculation*, Phys. Rev., 71 (1947), pp. 215–231.
- [29] C. CACCIAPUOTI, D. FINCO, AND D. NOJA, *Well posedness of the nonlinear Schrödinger equation with isolated singularities*, J. Differential Equations, 305 (2021), pp. 288–318.
- [30] Y. CASTIN AND F. WERNER, *The Unitary Gas and its Symmetry Properties*, in The BCS-BEC Crossover and the Unitary Fermi Gas, W. Zwerger, ed., vol. 836 of Lecture Notes in Physics, Springer Berlin Heidelberg, 2012, pp. 127–191.
- [31] H. D. CORNEAN, A. MICHELANGELI, AND K. YAJIMA, *Erratum: Two-dimensional Schrödinger operators with point interactions: threshold expansions, zero modes and  $L^p$ -boundedness of wave operators*, Rev. Math. Phys., 32 (2020), pp. 2092001, 5.
- [32] M. CORREGGI, G. DELL’ANTONIO, D. FINCO, A. MICHELANGELI, AND A. TETA, *Stability for a system of  $N$  fermions plus a different particle with zero-range interactions*, Rev. Math. Phys., 24 (2012), pp. 1250017, 32.
- [33] ———, *A Class of Hamiltonians for a Three-Particle Fermionic System at Unitarity*, Mathematical Physics, Analysis and Geometry, 18 (2015).
- [34] S. CUCCAGNA AND M. MAEDA, *On stability of small solitons of the 1-D NLS with a trapping delta potential*, SIAM J. Math. Anal., 51 (2019), pp. 4311–4331.
- [35] ———, *A survey on asymptotic stability of ground states of nonlinear Schrödinger equations II*, Discrete Contin. Dyn. Syst. Ser. S, 14 (2021), pp. 1693–1716.
- [36] P. D’ANCONA, V. PIERFELICE, AND A. TETA, *Dispersive estimate for the Schrödinger equation with point interactions*, Math. Methods Appl. Sci., 29 (2006), pp. 309–323.
- [37] F. F. G. DELLA CASA AND A. SACCHETTI, *Stationary states for non linear one-dimensional Schrödinger equations with singular potential*, Phys. D, 219 (2006), pp. 60–68.
- [38] G. DELL’ANTONIO, A. MICHELANGELI, R. SCANDONE, AND K. YAJIMA,  *$L^p$ -Boundedness of Wave Operators for the Three-Dimensional Multi-Centre Point Interaction*, Ann. Henri Poincaré, 19 (2018), pp. 283–322.
- [39] G. F. DELL’ANTONIO, R. FIGARI, AND A. TETA, *Hamiltonians for systems of  $N$  particles interacting through point interactions*, Ann. Inst. H. Poincaré Phys. Théor., 60 (1994), pp. 253–290.
- [40] Y. N. DEMKOV AND V. N. OSTROVSKII, *Zero-Range Potentials and Their Applications in Atomic Physics*, Physics of Atoms and Molecules, Springer US, 1988.
- [41] V. DUCHÈNE, J. L. MARZUOLA, AND M. I. WEINSTEIN, *Wave operator bounds for one-dimensional Schrödinger operators with singular potentials and applications*, J. Math. Phys., 52 (2011), pp. 013505, 17.
- [42] E. FERMI, *Sul moto dei neutroni nelle sostanze idrogenate*, Ricerca sci., 7 (1936), pp. 13–52.
- [43] V. FRANCESCHI, D. PRANDI, AND L. RIZZI, *On the essential self-adjointness of singular sub-Laplacians*, Potential Anal., 53 (2020), pp. 89–112.
- [44] N. FUKAYA, V. GEORGIEV, AND M. IKEDA, *On stability and instability of standing waves for 2d-nonlinear Schrödinger equations with point interaction*, arXiv:2109.04680 (2021).
- [45] M. GALLONE AND A. MICHELANGELI, *Quantum particle across Grushin singularity*, Journal of Physics A: Mathematical and Theoretical, (2021).
- [46] M. GALLONE, A. MICHELANGELI, AND E. POZZOLI, *On geometric quantum confinement in Grushin-type manifolds*, Z. Angew. Math. Phys., 70 (2019), pp. Art. 158, 17.
- [47] ———, *Geometric confinement and dynamical transmission of a quantum particle in Grushin cylinder*, arXiv:2003.07128 (2020).
- [48] ———, *Heat equation with inverse-square potential of bridging type across two half-lines*, arXiv.org:2112.01255 (2021).
- [49] V. GEORGIEV, A. MICHELANGELI, AND R. SCANDONE, *On fractional powers of singular perturbations of the Laplacian*, J. Funct. Anal., 275 (2018), pp. 1551–1602.
- [50] M. GRIESEMER, M. HOFACKER, AND U. LINDEN, *From Short-Range to Contact Interactions in the 1d Bose Gas*, Math. Phys. Anal. Geom., 23 (2020), pp. Paper No. 19, 28.
- [51] F. IANDOLI AND R. SCANDONE, *Dispersive estimates for Schrödinger operators with point interactions in  $\mathbb{R}^3$* , in Advances in Quantum Mechanics: Contemporary Trends and Open

- Problems, A. Michelangeli and G. Dell'Antonio, eds., Springer INdAM Series, vol. 18, Springer International Publishing, pp. 187–199.
- [52] I. IANNI, S. LE COZ, AND J. ROYER, *On the Cauchy problem and the black solitons of a singularly perturbed Gross-Pitaevskii equation*, SIAM J. Math. Anal., 49 (2017), pp. 1060–1099.
- [53] M. IKEDA AND T. INUI, *Global dynamics below the standing waves for the focusing semilinear Schrödinger equation with a repulsive Dirac delta potential*, Anal. PDE, 10 (2017), pp. 481–512.
- [54] S. MASAKI, J. MURPHY, AND J.-I. SEGATA, *Modified scattering for the one-dimensional cubic NLS with a repulsive delta potential*, Int. Math. Res. Not. IMRN, (2019), pp. 7577–7603.
- [55] ———, *Stability of small solitary waves for the one-dimensional NLS with an attractive delta potential*, Anal. PDE, 13 (2020), pp. 1099–1128.
- [56] A. MICHELANGELI, *Models of zero-range interaction for the bosonic trimer at unitarity*, Rev. Math. Phys., 33 (2021), pp. Paper No. 2150010, 101.
- [57] A. MICHELANGELI, A. OLGIATI, AND R. SCANDONE, *Singular Hartree equation in fractional perturbed Sobolev spaces*, J. Nonlinear Math. Phys., 25 (2018), pp. 558–588.
- [58] A. MICHELANGELI AND A. OTTOLINI, *On point interactions realised as Ter-Martirosyan-Skorniyakov Hamiltonians*, Rep. Math. Phys., 79 (2017), pp. 215–260.
- [59] ———, *Multiplicity of self-adjoint realisations of the  $(2+1)$ -fermionic model of Ter-Martirosyan—Skorniyakov type*, Rep. Math. Phys., 81 (2018), pp. 1–38.
- [60] A. MICHELANGELI AND P. PFEIFFER, *Stability of the  $(2+2)$ -fermionic system with zero-range interaction*, Journal of Physics A: Mathematical and Theoretical, 49 (2016), p. 105301.
- [61] A. MICHELANGELI AND C. SCHMIDBAUER, *Binding properties of the  $(2+1)$ -fermion system with zero-range interspecies interaction*, Phys. Rev. A, 87 (2013), p. 053601.
- [62] R. A. MINLOS, *A system of three quantum particles with point-like interactions*, Uspekhi Mat. Nauk, 69 (2014), pp. 145–172.
- [63] ———, *On point-like interaction of three particles: two fermions and another particle. II*, Mosc. Math. J., 14 (2014), pp. 617–637, 642–643.
- [64] R. A. MINLOS AND L. D. FADDEEV, *Comment on the problem of three particles with point interactions*, Soviet Physics JETP, 14 (1962), pp. 1315–1316.
- [65] ———, *On the point interaction for a three-particle system in quantum mechanics*, Soviet Physics JETP, 6 (1962), pp. 1072–1074.
- [66] A. MOGILNER AND M. SHERMATOV, *Binding of two fermions with a third different particle by a point interaction*, Physics Letters A, 149 (1990), pp. 398–400.
- [67] T. MOSER AND R. SEIRINGER, *Stability of a fermionic  $N + 1$  particle system with point interactions*, Comm. Math. Phys., 356 (2017), pp. 329–355.
- [68] ———, *Stability of the  $2 + 2$  fermionic system with point interactions*, Math. Phys. Anal. Geom., 21 (2018), pp. Paper No. 19, 13.
- [69] P. NAIDON AND S. ENDO, *Efimov physics: a review*, Reports on Progress in Physics, 80 (2017), p. 056001.
- [70] G. NENCIU AND I. NENCIU, *On confining potentials and essential self-adjointness for Schrödinger operators on bounded domains in  $\mathbb{R}^n$* , Ann. Henri Poincaré, 10 (2009), pp. 377–394.
- [71] C. J. PETHICK AND H. SMITH, *Bose–Einstein Condensation in Dilute Gases*, Cambridge University Press, second ed., 2008. Cambridge Books Online.
- [72] E. POZZOLI, *Quantum Confinement in  $\alpha$ -Grushin planes*, in Mathematical Challenges of Zero-Range Physics, A. Michelangeli, ed., Springer INdAM Series, Springer International Publishing, 2021, pp. 229–237.
- [73] D. PRANDI, L. RIZZI, AND M. SERI, *Quantum confinement on non-complete Riemannian manifolds*, J. Spectr. Theory, 8 (2018), pp. 1221–1280.
- [74] G. V. SKORNYAKOV AND K. A. TER-MARTIROSYAN, *Three Body Problem for Short Range Forces. I. Scattering of Low Energy Neutrons by Deuterons*, Sov. Phys. JETP, 4 (1956), pp. 648–661.

- [75] A. TETA, *Quadratic forms for singular perturbations of the Laplacian*, Publ. Res. Inst. Math. Sci., 26 (1990), pp. 803–817.
- [76] L. H. THOMAS, *The Interaction Between a Neutron and a Proton and the Structure of  $H^3$* , Phys. Rev., 47 (1935), pp. 903–909.
- [77] E. WIGNER, *On the Mass Defect of Helium*, Phys. Rev., 43 (1933), pp. 252–257.
- [78] K. YAJIMA,  *$L^p$ -boundedness of wave operators for 2D Schrödinger operators with point interactions*, Ann. Henri Poincaré, 22 (2021), pp. 2065–2101.
- [79] K. YOSHITOMI, *Finiteness of the discrete spectrum in a three-body system with point interaction*, Math. Slovaca, 67 (2017), pp. 1031–1042.



## Mini-Workshop: Zero-Range and Point-Like Singular Perturbations: For a Spillover to Analysis, PDE and Differential Geometry

### Table of Contents

Riccardo Adami	
<i>Two-dimensional systems with non-linear point interactions</i> . . . . .	2613
Ivan Beschastnyi	
<i>Self-adjoint extensions for the curvature Laplacian on Grushin manifolds</i>	2613
Filippo Boni	
<i>NLS ground states with singularities</i> . . . . .	2613
Ugo Boscain	
<i>Geometric confinement of the curvature Laplacian on almost Riemannian manifolds</i> . . . . .	2614
Luca Fanelli (joint with Luz Roncal, and Nico Michele Schiavone)	
<i>About Schrödinger and Dirac operators with scaling critical potentials</i> ..	2614
Valentina Franceschi	
<i>Pointed sub-Laplacians in three dimensions and Hardy inequalities</i> . . . . .	2616
Matteo Gallone (joint with Alessandro Michelangeli, Eugenio Pozzoli)	
<i>The Laplace-Beltrami operator on the Grushin Cylinder</i> . . . . .	2616
Marilena Ligabò (joint with Paolo Facchi, Fabio Deelan Cunden, Giancarlo Garnero)	
<i>Boundary conditions, product formulae and classical limit</i> . . . . .	2617
Diego Noja (joint with Claudio Cacciapuoti and Domenico Finco)	
<i>The NLS equation with a point interaction in two and three dimensions</i>	2618
Mario Rastrelli (joint with Vladimir S. Georgiev)	
<i>On the square of Laplacian with inverse square potential</i> . . . . .	2619
Raffaele Scandone (joint with Vladimir Georgiev, Alessandro Michelangeli)	
<i>Non-linear Schrödinger equations with point interactions: results and perspectives</i> . . . . .	2619
Ivana Vojnović (joint with Nevena Dugandžija, Alessandro Michelangeli)	
<i>Generalised solutions to non-linear Schrödinger equations with singularities</i> . . . . .	2620
Jens Wirth (joint with Jonas Brinker, Michael Ruzhansky)	
<i>Operators on Groups and Non-Commutative Fourier Transforms</i> . . . . .	2621



## Abstracts

### Two-dimensional systems with non-linear point interactions

RICCARDO ADAMI

Non-linear point interactions were introduced two decades ago in order to describe localised phenomena like trapping, beatings, concentration of wave packets, and so on. Seminal studies focussed on one and three-dimensional systems, leaving untouched the more technical two-dimensional case up to two years ago. We review the results and comment on the specific features exhibited by this case, together with open problems.

### REFERENCES

- [1] R. Adami, F. Boni, R. Carlone, L. Tentarelli, *Ground states for the planar NLSE with a point defect as minimisers of the constrained energy*, Calc. Var. **61** (2022) 195

### Self-adjoint extensions for the curvature Laplacian on Grushin manifolds

IVAN BESCHASTNYI

I discuss some results from an ongoing project with H. Quan (University of Washington), in which we study the curvature Laplacian on curved high-dimensional analogues of the  $\alpha$ -Grushin planes. We are able to prove that all self-adjoint properties of the Laplacian are encoded in a second order polynomial, which can be easily read off from the equation itself. In particular, we can determine whether the operator is essentially self-adjoint by computing its discriminant and construct all of its self-adjoint extensions from its roots.

### NLS ground states with singularities

FILIPPO BONI

We investigate the existence of ground states at fixed mass of the  $L^2$ -subcritical non-linear Schrödinger equation with a point interaction. First, the problem is considered in dimension two and three. We prove that ground states exist for every value of the mass and, up to a multiplication by a phase factor, they are positive, radially symmetric, decreasing along the radial direction and present a singularity where the interaction is placed. In order to obtain qualitative features of the ground states, we refine a classical result on rearrangements and move to equivalent variational formulations of the problem. Then, we present some future developments, among them the possibility of generalising similar models on hybrid structures.

## REFERENCES

- [1] R. Adami, F. Boni, R. Carlone, L. Tentarelli, *Existence, structure, and robustness of ground states of a NLSE in 3D with a point defect*, J. Math. Phys. **63** (2022) 071501

## Geometric confinement of the curvature Laplacian on almost Riemannian manifolds

UGO BOSCAIN

Two-dimensional almost-Riemannian structures of step 2 are natural generalisations of the Grushin plane. They are generalised Riemannian structures for which the vectors of a local orthonormal frame can become parallel. Under the 2-step assumption the singular set  $Z$ , where the structure is non-Riemannian, is a one-dimensional embedded submanifold. While approaching the singular set, all Riemannian quantities diverge. A remarkable property of these structures is that the geodesics can cross the singular set without singularities, but the heat and the solution to the Schrödinger equation (with the Laplace-Beltrami operator  $\Delta$  cannot. This is due to the fact that (under a natural compactness hypothesis), the Laplace-Beltrami operator is essentially self-adjoint on a connected component of the manifold without the singular set. In the literature such a counter-intuitive phenomenon is called geometric confinement.

For the heat equation an intuitive explanation of this fact can be given in terms of random walks. For the Schrödinger equation an intuitive explanation is more subtle, as the evolution of a quantum particle on a manifold can be done in several non-equivalent ways.

In this talk I describe the evolution (and the confinement) of a quantum particle described by the curvature Laplacian  $-\Delta + cK$  (here  $K$  is the Gaussian curvature and  $c > 0$  a constant) which originates in coordinate-free quantisation procedures (as, for instance, in path-integral or covariant Weyl quantisation).

## REFERENCES

- [1] I. Beschastnyi, U. Boscaín, E. Pozzoli, *Quantum Confinement for the Curvature Laplacian  $-\Delta + cK$  on 2D-Almost-Riemannian Manifolds*, Potential Anal. (2021)

## About Schrödinger and Dirac operators with scaling critical potentials

LUCA FANELLI

(joint work with Luz Roncal, and Nico Michele Schiavone)

Lower-order perturbations of the free Hamiltonians usually appear in Quantum Mechanics, as models describing the interaction of a free particle with an external field. In some cases, the perturbation lies at the same level as the free Hamiltonian, and the resulting conflict can generate interesting phenomena. We will introduce the Inverse Square and Coulomb potentials as toy models, and describe the main features of the complete Hamiltonians from the point of view of Fourier Analysis, Spectral Theory, and dispersive evolutions. In this talk, the potentials will usually

belong to  $\mathbb{C}$ , which is responsible of a lack of symmetry, and we hence are in the case of non-self-adjoint Hamiltonians. In the recent years, there is a growing interest devoted to this kind of operators, and it is quite surprising that it started around 25 years ago, after almost one century since the advent of Quantum Mechanics. Motivated by needs of Nuclear Physics, Scholtz-Geyer-Hahne [12] suggested an interesting model in which observables are represented by operators which are not necessarily self-adjoint, but merely quasi-self-adjoint” (roughly speaking, similar to self-adjoint operators). In this case, it is sufficient to change the inner product in the base Hilbert space, with the help of a metric operator related to the similarity, to end back in the usual Hermitian case. Later on, Bender-Boettcher [2] noticed that a large class of non-self-adjoint operators still possesses a real spectrum, and renewed the interest to the topic.

In the recent papers [4, 6, 7], we proved by purely real analytic methods some uniform resolvent estimates for the free Hamiltonian, inspired to the pioneer results by Kato and Yajima in [11]. Those are uniform version over the complex plane of some weighted variant of the so called *Hardy-Rellich* inequality, which has been recently investigated in [3] with a particular interest on the sharp constants. In this talk, we will present the recent results obtained in [8], concerned with the analogous questions in the sub-Riemannian setting of the Heisenberg Group. Some interesting open problems related with the spectrum of mass-less Dirac operators and the sharp constants of the uniform resolvent estimates will also be introduced.

#### REFERENCES

- [1] M. Muster, *Computing certain invariants of topological spaces of dimension three*, *Topology* **32** (1990), 100–120.
- [2] C. Bender, and S. Boettcher, *Real spectra in non-Hermitian Hamiltonians having PT-symmetry*, *Physical Review Letters*, **80(24)** (1998)
- [3] B. Cassano, L. Cossetti, and L. Fanelli, *Improved Hardy-Rellich Inequalities*, *Comm. Pure Appl. Anal.* **21** (2022), 867–889.
- [4] L. Cossetti, L. Fanelli and D. Krejčířík, *Absence of eigenvalues of Dirac and Pauli Hamiltonians via the method of multipliers*, *Comm. Math. Phys.* **379** (2020), 633–691.
- [5] L. D’Ambrosio, *Hardy-type inequalities related to degenerate elliptic differential operators*, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **5** (2005), 451–486.
- [6] L. Fanelli, D. Krejčířík, and L. Vega, *Spectral stability of Schrödinger operators with subordinated complex potentials*, *J. Spectr. Theory* **8** (2018), 575–604.
- [7] L. Fanelli, D. Krejčířík, and L. Vega, *Absence of eigenvalues of two-dimensional magnetic Schrödinger operators*, *J. Funct. Anal.* **275** (2018), no. 9, 2453–2472.
- [8] L. Fanelli, L. Roncal, and N. Schiavone, *Non self-adjoint perturbations of the Heisenberg Sublaplacian*, <https://arxiv.org/abs/2203.07296> (2022)
- [9] N. Garofalo and E. Lanconelli, *Frequency functions on the Heisenberg group, the uncertainty principle and unique continuation*, *Ann. Inst. Fourier (Grenoble)* **40** (1990), 313–356.
- [10] T. Ikebe, and Y. Saito, *Limiting absorption method and absolute continuity for the Schrödinger operator*, *J. Math. Kyoto Univ.* **12** (1972), 513–542.
- [11] T. Kato and K. Yajima, *Some examples of smooth operators and the associated smoothing effect*, *Rev. Math. Phys.* **1(4)** (1989), 481–496.
- [12] FG. Scholtz, HB Geyer, and FJW Hahne, *Quasi-Hermitian operators in quantum mechanics and the variational principle*, *Annals of Physics* **213(1)** (1992), 74–101

## Pointed sub-Laplacians in three dimensions and Hardy inequalities

VALENTINA FRANCESCHI

The aim of this seminar is to present recent results on the essential self-adjointness of pointed sub-Laplacians in three dimensions. We show that, unlike the Euclidean case, pointed sub-Laplacians (associated with smooth measures) are essentially self-adjoint in dimension 3. To this purpose, we focus on the case of the 3D Heisenberg sub-Laplacian, and we show its essential self-adjointness by exploiting non-commutative Fourier transform techniques. We then generalise the result to a class of three-dimensional sub-Riemannian manifolds. In connection with the main result, we present a discussion on Hardy inequalities in the Heisenberg group: contrary to the Euclidean case, a radial Hardy inequality, i.e., a Hardy inequality taking into account only the directional derivative with respect to the sub-Riemannian distance, does not hold in this context for any dimension. This underlines again a difference with respect to the Euclidean case, where essential self-adjointness of pointed Laplacians can be derived from Hardy inequalities.

### REFERENCES

- [1] U. Boscain, V. Franceschi, D. Prandi, R. Adami, *Point interactions for 3D sub-Laplacians*, Ann. Inst. H. Poincaré Anal. Non Linéaire **38** (2021), no. 4, pp. 1095–1113

## The Laplace-Beltrami operator on the Grushin Cylinder

MATTEO GALLONE

(joint work with Alessandro Michelangeli, Eugenio Pozzoli)

When a quantum particle is constrained on an orientable Riemannian manifold, one challenging problem that arises naturally is the question of the so-called ‘geometric quantum confinement’. This is the possibility that a particle whose initial wave-function is supported inside some portion of space may remain confined in such a region for all times when evolving according to the unitary group generated by the free Hamiltonian. This occurrence is related to the presence of singularities in the metric and to the (essential) self-adjointness of the Laplace-Beltrami operator. The prototypical example of space exhibiting this phenomenon is the ‘Grushin-like cylinder’, that is, roughly speaking, a cylinder with metric  $ds^2 = dx^2 + |x|^{-2\alpha} dy^2$ . In this talk I consider this manifold and present the classification of a physically interesting sub-family of self-adjoint realisations of the Laplace-Beltrami operator in the regime where it is not essentially self-adjoint. I discuss the advantages of the usage of the Kreĭn-Višik-Birman self-adjoint extension theory.

### REFERENCES

- [1] M. Gallone, A. Michelangeli, E. Pozzoli, *On Geometric Quantum Confinement in Grushin-type Manifolds*, Z. Angew. Math. Phys. **70** (2019) 158  
 [2] M. Gallone, A. Michelangeli, E. Pozzoli, *Quantum geometric confinement and dynamical transmission in Grushin cylinder*, Rev. Math. Phys. **34** (2022) 2250018

- [3] M. Gallone, A. Michelangeli, *Quantum particle across Grushin singularity* J. Phys. A: Math. Theor. **54** (2021) 215201
- [4] M. Gallone, A. Michelangeli, E. Pozzoli, *Heat equation with inverse-square potential of bridging type across two half-lines*, in Qualitative Properties of Dispersive PDEs, INdAM-Springer series, vol. 52, (2022) 141-164

## Boundary conditions, product formulae and classical limit

MARILENA LIGABÒ

(joint work with Paolo Facchi, Fabio Deelan Cunden, Giancarlo Garnero)

In this seminar some connection between (classical and quantum) boundary condition and product formulae will be illustrated. The starting point of this presentation is a result concerning a parametrisation of all the possible self-adjoint extensions of the Laplace operator on a bounded domain in terms of unitary operators at the boundary [1]. Then, using the Trotter product formula, a composition law for quantum boundary conditions will be defined [2]. Furthermore, the link between the occurrence of boundary conditions and frequent measurements on a quantum system will be discussed. This phenomenon is due to the existence of a one-parameter unitary group obtained as the limit of the Zeno product formula (defined by intertwining the time evolution group with an orthogonal projection) [3, 4]. The limiting dynamics is called Zeno dynamics and the corresponding generator is called Zeno Hamiltonian. Finally, it will be shown that the classical limit of a Zeno dynamics obtained in a cavity quantum electrodynamics experiment [5, 6] induces a change of topology in the classical phase space [7].

## REFERENCES

- [1] P. Facchi, G. Garnero, Marilena Ligabò., *Self-adjoint extensions and unitary operators on the boundary*, Letters in Mathematical Physics **108** (2018), 195–212.
- [2] P. Facchi, G. Garnero, Marilena Ligabò, *Quantum cavities with alternating boundary conditions*, Journal of Physics A: Mathematical and Theoretical **51** (2018), 105301.
- [3] P. Exner, T. Ichinose, *Note on a Product Formula Related to Quantum Zeno Dynamics*, Ann. Henri Poincaré **22** (2021), 1669–1697.
- [4] P. Facchi, M. Ligabò, *Quantum Zeno effect and dynamics*, Journal of Mathematical Physics **51** (2010), 022103.
- [5] J. M. Raimond, C. Sayrin, S. Gleyzes, I. Dotsenko, M. Brune, S. Haroche, P. Facchi, and S. Pascazio, *Phase Space Tweezers for Tailoring Cavity Fields by Quantum Zeno Dynamics*, Phys. Rev. Lett. **105** (2010), 213601.
- [6] J. M. Raimond, P. Facchi, B. Peaudecerf, S. Pascazio, C. Sayrin, I. Dotsenko, S. Gleyzes, M. Brune, S. Haroche, *Quantum Zeno dynamics of a field in a cavity*, Physical Review A **86** (2012), 032120.
- [7] F. D. Cunden, P. Facchi, M. Ligabò, *Classical limit of a truncated momentum operator*, in preparation (2022).

## The NLS equation with a point interaction in two and three dimensions

DIEGO NOJA

(joint work with Claudio Cacciapuoti and Domenico Finco)

The NLS equation with a power non-linearity and point interaction is described by the Cauchy problem

$$i\partial_t\psi = H_\alpha\psi + g|\psi|^{p-1}\psi \quad p > 1, \quad g = \pm 1$$

where  $H_\alpha$  is the abstract Schrödinger-like singular perturbation of the Laplacian known as *point interaction* or *delta potential*, sometimes improperly written as  $H_\alpha = -\Delta + \alpha\delta$ . The model is well studied in dimension one but only recently the two and three dimensional cases have been considered by several authors. In the talk, after a review of the definition, construction and properties of point interactions especially relevant to the applications in the nonlinear setting, the well-posedness of the Cauchy problem will be treated, as regards the known facts and reviewing also some still open problems. Moreover, some asymptotic properties of the model will be discussed, and in particular as regards the blow-up for strong non-linearities in dimension two. Namely, it will be shown that in two dimensions and in the super-critical regime  $p > 3$  large sets of initial data undergo blow-up and that the standing waves are strongly unstable, in the sense that arbitrarily close (in the energy norm) to any standing wave there exist blowing-up initial data.

### REFERENCES

- [1] R.Adami, F. Boni, R.Carlone, L.Tentarelli, Ground states for the planar NLSE with a point defect as minimisers of the constrained energy, *Calc.Var.&PDE* **61**, (2022) 195
- [2] C.Cacciapuoti, D.Finco, D.Noja, *Well posedness of the nonlinear Schrödinger equation with isolated singularities*, *JDE* **305** (2021) 288-318
- [3] H.D. Cornean, A.Michelangeli, K.Yajima, *Two-dimensional Schrödinger operators with point interactions: Threshold expansions, zero modes and  $L^p$ -boundedness of wave operators*, *Rev. Math. Phys.* **31** (2019), 1950012
- [4] D.Finco, D.Noja *Blow-up for the nonlinear Schrödinger equation with a point interaction in dimension two*, (2022), arXiv:2209.09537
- [5] N.Fukaya, V.Georgiev, M.Ikeda, *On stability and instability of standing waves for 2d-nonlinear Schrödinger equations with point interaction* *JDE*, **321** (2022), 258-295
- [6] V.Georgiev, A.Michelangeli, R. Scandone, *Standing waves and global well-posedness for the 2d Hartree equation with a point interaction* arXiv:2204.05053v1 (2022)
- [7] M. Ohta, T. Yamaguchi, *Strong instability of standing waves for nonlinear Schrödinger equations with a delta potential*, *RIMS Kokyuroku Bessatsu* **B 56** (2016) 79-92

## On the square of Laplacian with inverse square potential

MARIO RASTRELLI

(joint work with Vladimir S. Georgiev)

The aim of the talk is to show an explicit characterisation of the domain of the scaling-invariant perturbation of the Laplacian, the operator  $A = -\Delta + \frac{\beta}{|x|^2}$ , and of its square. For the domain of  $A$ , we require  $\beta > \frac{n(n-4)}{4}$  that is the necessary condition for essentially self-adjointness of the operator [8, 9]. For  $A^2$  it is still not clear what happens when  $\beta = 8 + \frac{n(n-4)}{4}$ . It is also more difficult to give an expression of the domain in even dimensions, when  $n < 8$ , due to the problems that arise with weighted Rellich inequalities. The description of  $D(A)$  gives to us a Rellich type inequality that can be used to study Schrödinger operators with singular potential of the inverse-square type [5], that hold even if  $n = 3, 4$ . The characterisation of  $D(A^2)$  improve this last result, giving to us a weighted version of the inequality.

### REFERENCES

- [1] Balinsky, Alexander A. and Evans, W. Desmond and Lewis, Roger T. *The Analysis and Geometry of Hardy's Inequality*, Springer (2010),
- [2] Bennett, D. M., *An Extension of Rellich's inequality*, Proceedings of the American Mathematical Society Vol. 106, No. 4, (1989), pp. 987–983.
- [3] B. Cassano, L. Cossetti, L. Fanelli, *Improved Hardy-Rellich inequalities*, arXiv:2106.09804v2 [**math.AP**] (2021) .
- [4] Cazacu, Cristian, *A new proof of the Hardy-Rellich inequality in any dimension*, Proc. Roy. Soc. Edinburgh Sect. A, 150 (2020), no. 6, 2894-2904 , <https://doi.org/10.48550/arXiv.1809.07506>
- [5] Georgiev, V. and Kubo, H, *Global solvability for nonlinear wave equations with singular potential*, arxiv, 2022,
- [6] Georgiev, V. and Kubo, H, *On the Rellich type inequality for Schrödinger operators with singular potential*, Proceeding ISAAC, Special Volume on Harmonic Analysis and PDEs (2022)
- [7] Georgiev, V. and Rastrelli, M., *On the square of Laplacian with inverse square potential*, Proceeding NTADES (2022)
- [8] Kalf, H. and Schmincke, U.-W. and Walter, J. and Wüst, R., *On the spectral theory of Schrödinger and Dirac operators with strongly singular potentials*, Spectral theory and differential equations, pp. 182–226, Springer, Berlin (1975).
- [9] Simon, B., *Essential self-adjointness of Schrödinger operators with singular potentials*, Arch. Rational Mech. Anal. Vol. 52, (1973), pp. 44–48.

## Non-linear Schrödinger equations with point interactions: results and perspectives

RAFFAELE SCANDONE

(joint work with Vladimir Georgiev, Alessandro Michelangeli)

In this talk I review recent advances in the study of non-linear Schrödinger equations with delta-like potentials. After introducing a suitable functional framework, the notion of sub-critical non-linearity, and a class of dispersive-type estimates, I

discuss some results concerning the global well-posedness and the existence (and symmetries) of standing waves. The last part of the talk is devoted to open questions and future research perspectives.

## Generalised solutions to non-linear Schrödinger equations with singularities

IVANA VOJNOVIĆ

(joint work with Nevena Dugandžija, Alessandro Michelangeli)

We consider Schrödinger equations of Hartree and cubic type in three spatial dimensions and its approximations of singular, point-like perturbations.

As approximants to the Hartree equation, we analyse the equation of the form

$$(1) \quad i\partial_t u_\varepsilon = -\Delta u_\varepsilon + V_\varepsilon u_\varepsilon + (w * |u_\varepsilon|^2)u_\varepsilon,$$

for  $\varepsilon \in (0, 1]$ . Here  $V_\varepsilon$  is a real-valued potential and is meant to represent a singular, delta-like profile centred at  $x = 0$ .

We assume that

$$(2) \quad V_\varepsilon(x) := \frac{1}{\varepsilon^\sigma} V\left(\frac{x}{\varepsilon}\right),$$

for a given measurable function  $V : \mathbb{R}^3 \rightarrow \mathbb{R}$  and a given  $\sigma \geq 0$ .

The corresponding nets of approximate solutions represent generalised solutions for the singular-perturbed Schrödinger equation. The behaviour of such nets is investigated for  $\sigma \in [0, 3]$ .

We also study a generalised solution in the Colombeau algebra  $\mathcal{G}_{C^1, H^2}$  for cubic and Hartree equation with delta potential, which correspond to the case  $\sigma = 3$  in (2). In the case of the Hartree equation with delta potential compatibility between the Colombeau solution and the solution of the classical Hartree equation is established. More precisely, we prove that

$$(3) \quad \lim_{\varepsilon \downarrow 0} \|u_\varepsilon - u\|_{L^\infty([0, T], L^2(\mathbb{R}^3))} = 0,$$

where Colombeau solution is represented by net  $(u_\varepsilon)$  and  $u$  is the unique solution in  $C(\mathbb{R}, L^2(\mathbb{R}^3))$  to the Cauchy problem

$$i\partial_t u = -\Delta u + (w * |u|^2)u$$

with initial datum  $a \in L^2(\mathbb{R}^3)$ .

## REFERENCES

- [1] S. Albeverio, F. Gesztesy, R. Høegh-Krohn, and H. Holden, *Solvable models in quantum mechanics*, AMS Chelsea Publishing, Providence, RI, (2005)
- [2] N. Dugandžija, A. Michelangeli and I. Vojnović, *Generalised solutions to linear and non-linear Schrödinger-type equations with point defect: Colombeau and non-Colombeau regimes*, arXiv:2209.07207
- [3] N. Dugandžija and M. Nedeljkov, *Generalized solution to multidimensional cubic Schrödinger equation with delta potential*, *Monatsh. Math.*, 190 (2019), 481–499.

- [4] N. Dugandžija and I. Vojnović, *Nonlinear Schrödinger equation with singularities*, to appear in: *Qualitative Properties of Dispersive PDEs*, V. Georgiev, A. Michelangeli, and R. Scandone editors, Springer-INdAM Series, (2022)
- [5] A. Michelangeli, A. Olgiati, and R. Scandone, *Singular Hartree equation in fractional perturbed Sobolev spaces*, *J. Nonlinear Math. Phys.*, 25 (2018), 558–588.

## Operators on Groups and Non-Commutative Fourier Transforms

JENS WIRTH

(joint work with Jonas Brinker, Michael Ruzhansky)

The study of differential and more general pseudo-differential operators is usually based on Euclidean Fourier transform

$$\widehat{f}(\xi) = \int_{\mathbb{R}^n} e^{-2\pi i x \cdot \xi} f(x) dx, \quad f \in \mathcal{S}(\mathbb{R}^n),$$

which allows to treat differential operators  $p(D) = \sum_{|\alpha| \leq m} a_\alpha D^\alpha$ ,  $D = \frac{i}{2\pi} \partial$  in terms of their symbols  $p(\xi)$ . More generally, for  $\sigma_A(x, \xi) \in C_b^\infty(\mathbb{R}^n) \widehat{\otimes} \mathcal{O}_M(\mathbb{R}^n)$  one considers operators of the form

$$Af(x) = \int_{\mathbb{R}^n} e^{2\pi i x \cdot \xi} \sigma_A(x, \xi) \widehat{f}(\xi) d\xi.$$

For details on pseudo-differential operators and their calculus see [1] and [2]. Although, (almost) any operator can be treated in such a way, the translation invariance of the constant coefficient differential operators is built into the calculus.

Sometimes it is advantageous to treat operators based on different symmetries and therefore use non-commutative analogues of the classical Fourier transform.

**The non-commutative Fourier transform.** For a locally compact type-I group  $G$  we denote by  $\widehat{G}$  the set of equivalence classes of irreducible unitary representations  $\xi : G \rightarrow \mathcal{L}(H_\xi)$  and for each  $[\xi] \in \widehat{G}$  define

$$\widehat{f}(\xi) = \int_G f(x) \xi(x)^* dx$$

as non-commutative Fourier transform. For sufficiently nice functions  $\widehat{f}(\xi)$  is of trace class and there exists a unique measure on  $\widehat{G}$ , the Plancherel measure  $\mu$ , such that the inversion formula

$$f(x) = \int_{\widehat{G}} \text{trace}(\widehat{f}(\xi) \xi(x)) d\mu(\xi)$$

holds true. Again this can be used as starting point to develop a pseudo-differential calculus, for details see e.g. [6]. It is worth looking at some examples to see how such a calculus will look like and what benefit it brings.

**Compact groups.** If  $G$  is compact, irreducible representations  $[\xi] \in \widehat{G}$  are finite-dimensional, i.e.  $H_\xi \simeq \mathbb{C}^{d_\xi}$ , and the inversion of non-commutative Fourier transform reduces to Peter–Weyl theorem

$$L^2(G) = \bigoplus_{[\xi] \in \widehat{G}} \text{span}\{\xi_{ij} \mid \xi(x) = (\xi_{ij}(x))_{1 \leq i, j \leq d_\xi}\}$$

in the form of series representations

$$f(x) = \sum_{[\xi] \in \widehat{G}} d_\xi \text{trace}(\widehat{f}([\xi])\xi(x)).$$

For  $G$  a compact Lie group any linear continuous operator  $A : C^\infty(G) \rightarrow \mathcal{D}'(G)$  can be represented in terms of a matrix-valued symbol

$$\sigma_A(x, \xi) = \xi(x)^*(A\xi)(x),$$

where  $A$  is applied to the matrix entries  $\xi_{ij}$ , as series

$$Af(x) = \sum_{[\xi] \in \widehat{G}} d_\xi \text{trace}(\widehat{f}([\xi])\sigma_A(x, \xi)\xi(x)).$$

Again symbols encode properties of the operator and allow for a full symbolic calculus under appropriate conditions, see [4] and [5]. To mention a typical example, Hörmander class pseudo-differential operators defined in local coordinates allow a global characterisation in terms of such global symbols. The operator  $A$  is pseudo-differential of order  $m$  and type  $(\rho, \delta)$  if

$$\sup_{x \in G} \|\partial_x^\alpha \Delta_\xi^\beta \sigma_A(x, \xi)\|_{\text{op}} \leq C_{\alpha, \beta} \langle \xi \rangle^{m - \rho|\beta| + \delta|\alpha|}$$

holds true with  $\langle \xi \rangle = \sqrt{1 + \lambda_\xi}$  given in terms of the Laplace eigenvalue  $\lambda_\xi$  on the eigenfunctions  $\xi_{ij}$ , an admissible selection of left-invariant vector fields  $\partial_x$  and an admissible selection of first order difference operators  $\Delta_\xi$ . First order difference operators by itself are defined in terms of the non-commutative Fourier transform,

$$\Delta \widehat{f} = \widehat{qf},$$

using differentiable functions  $q : G \rightarrow \mathbb{C}$  vanishing to first order in the identity element  $1 \in G$  and admissibility for a selection of such difference operators essentially means that their differentials in the identity span the full cotangent space.

**The Heisenberg group.** On the Heisenberg group  $\mathbb{H}_n = \mathbb{R}^{2n+1}$  with group law

$$(x, \xi, \tau) \bullet (y, \eta, \sigma) = (x + y, \xi + \eta, \tau + \sigma + \frac{1}{2}(y \cdot \eta - x \cdot \xi))$$

irreducible representations are characterised by Stone–von Neumann theorem. Of interest for us are only the Schrödinger representations

$$\rho_\lambda(x, \xi, \eta) = e^{2\pi i \tau + \pi i \lambda x \cdot \xi} M_\xi T_{\lambda x}$$

parametrised by  $\lambda \in \mathbb{R}^\times = \mathbb{R} \setminus \{0\}$  and given in terms of translations  $T_x$  and modulations  $M_\xi$  on the Hilbert space  $L^2(\mathbb{R}^n)$ . The Haar measure of the Heisenberg group is given by Lebesgue measure and the Fourier inversion formula is given by

$$f(x) = \int_{\mathbb{R}^\times} \text{trace}(\widehat{f}(\lambda)\rho_\lambda(x, \xi, \tau))|\lambda|^n d\lambda, \quad \widehat{f}(\lambda) = \widehat{f}(\rho_\lambda).$$

For more details, in particular on the resulting pseudo-differential calculus, see [7]. The main difference to the case of compact groups is that symbols  $\sigma_A$  of pseudo-differential operators  $A$  are by itself unbounded operators on the representation space  $L^2(\mathbb{R}^n)$  and should better by itself be seen as Weyl quantised pseudo-differential operators. The following table collects examples for this:

operator on $\mathbb{H}_n$	symbol as operator on $\mathbb{R}^n$	Weyl symbol
$\partial_{x_j} - \frac{1}{2}\xi_j\partial_\tau$ $\partial_{\xi_j} + \frac{1}{2}x_j\partial_\tau$ $\partial_\tau$	$\sqrt{ \lambda }\partial_{y_j}$ $i\sqrt{\lambda}y_j$ $i\lambda$	$\sqrt{ \lambda }\eta_j$ $i\sqrt{\lambda}y_j$ $i\lambda$
$\mathcal{L}_{sub}$	$ \lambda ^2 \sum_{j=1}^n (\partial_{u_j}^2 - u_j^2)$	$- \lambda ^2 \sum_{j=1}^n (y_j^2 + \eta_j^2)$

Symbols of pseudo-differential operators are again characterised in terms of smoothness and difference conditions. Now difference operators are in fact also differential operators acting on Weyl symbols

function on $\mathbb{H}_n$	difference operator on $\mathbb{R}^n$	difference operators on symbols
$x_j$ $\xi_j$ $\tau$	$ \lambda ^{-1/2} \text{ad } y_j$ $-i\lambda^{-1/2} \text{ad } \partial_{y_j}$ $\dots$	$ \lambda ^{-1/2} \partial_{\eta_j}$ $-i\lambda^{-1/2} \partial_{y_j}$ $\partial_\lambda - \frac{1}{2\lambda} \sum_{j=1}^n (y_j \partial_{y_j} + \eta_j \partial_{\eta_j})$

and in essence from these operators it can be seen that symbols of Hörmander class pseudo-differential operators on  $\mathbb{R}^n$  are in fact itself parameter-dependent pseudo-differential operators with symbols from Shubin classes.

**Smooth orbit characterisations.** Properties of operators  $A$  on a Hilbert space  $H$  can be encoded into smoothness properties of the orbit

$$x \mapsto \pi(x)^* A \pi(x)$$

with respect to unitary representations  $\pi : G \rightarrow \mathcal{L}(H)$ . It is well-known that  $C^\infty$ -smoothness of such orbits with respect to Schrödinger representations characterise pseudo-differential operators with symbols from  $S_{0,0}^0$ , see [3]. In [8], [9] this is generalised to ultra-differential smoothness for actions of compact and also for homogeneous groups.

REFERENCES

[1] L. Hörmander, *The analysis of linear partial differential operators. II. Differential operators with constant coefficients.* Springer-Verlag, Berlin, 1983.  
 [2] L. Hörmander, *The analysis of linear partial differential operators. III. Pseudodifferential operators.* Springer-Verlag, Berlin, 1985.  
 [3] H. O. Cordes. *The technique of pseudodifferential operators,* London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1995.

- 
- [4] M. Ruzhansky, V. Turunen, J. Wirth, *Hörmander class of pseudo-differential operators on compact Lie groups and global hypoellipticity*. J. Fourier Anal. Appl. 20:3 (2014), 476–499.
  - [5] M. Ruzhansky, J. Wirth, *Global functional calculus for operators on compact Lie groups*. J. Funct. Anal. 267:1 (2014), 144–172.
  - [6] M. Măntoiu, M. Ruzhansky, *Pseudo-differential operators, Wigner transform and Weyl systems on type I locally compact groups*. Doc. Math. 22 (2017), 1539–1592.
  - [7] V. Fischer, M. Ruzhansky, *Quantization on Nilpotent Lie groups*, Vol. 134 of Progress in Mathematics, Birkhäuser 2016
  - [8] J. Brinker, J. Wirth, *Gelfand triples for the Kohn-Nirenberg quantization on homogeneous Lie groups*. Advances in harmonic analysis and partial differential equations, 51–97, Birkhäuser/Springer, 2020
  - [9] J. Brinker, *Quantization of Algebras Defined by Ultradifferentiable Group Actions*, Dissertation, University of Stuttgart 2022

## Participants

**Prof. Dr. Riccardo Adami**

Dipartimento di Matematica  
Politecnico di Torino  
Corso Duca degli Abruzzi, 24  
10129 Torino  
ITALY

**Prof. Dr. h.c. Sergio Albeverio**

Institut für Angewandte Mathematik +  
HCM  
Universität Bonn  
Endenicher Allee 60  
53115 Bonn  
GERMANY

**Daniele Barbera**

Dip. di Matematica "L.Tonelli"  
Universita di Pisa  
Largo Bruno Pontecorvo, 5  
56127 Pisa  
ITALY

**Prof. Dr. Jacopo Bellazzini**

Dip. di Matematica "L.Tonelli"  
Universita di Pisa  
Largo Bruno Pontecorvo, 5  
56127 Pisa  
ITALY

**Dr. Ivan Beschastnyi**

Departamento de Matematica  
Universidade de Aveiro  
Campus Santiago  
Aveiro 3810-193  
PORTUGAL

**Dr. Filippo Boni**

Dipartimento di Matematica e  
Applicazioni "Renato Caccioppoli"  
Università degli Studi di Napoli Federico  
II  
Via Cintia, Monte S. Angelo  
80126 Napoli  
ITALY

**Prof. Dr. Ugo Boscin**

Laboratoire Jacques-Louis Lions  
Sorbonne Université  
4, Place Jussieu  
75005 Paris Cedex  
FRANCE

**Tommaso Cortopassi**

Scuola Normale Superiore di Pisa  
Piazza dei Cavalieri 7  
56126 Pisa  
ITALY

**Dr. Lucrezia Cossetti**

Fakultät für Mathematik  
Karlsruher Institut für Technologie(KIT)  
76128 Karlsruhe  
GERMANY

**Prof. Dr. Scipio Cuccagna**

Dip. di Matematica e Informatica  
Universita di Trieste  
Via Alfonso Valerio 12/1  
34127 Trieste  
ITALY

**Prof. Dr. Fabio Cunden**

Dipartimento di Matematica  
Universita di Bari  
70125 Bari  
ITALY

**Prof. Dr. Marko Erceg**  
Department of Mathematics  
University of Zagreb  
Bijenicka 30  
10000 Zagreb  
CROATIA

**Prof. Dr. Luca Fanelli**  
Universidad del Pais Vasco  
Facultad de Ciencias  
Departamento de Matematicas  
Apt. 644  
48940 Bilbao, Bizkaia  
SPAIN

**Dr. Valentina Franceschi**  
Dipartimento di Matematica  
Universita di Padova  
Via Trieste, 63  
35121 Padova  
ITALY

**Dr. Noriyoshi Fukaya**  
Department of Mathematics  
Tokyo University of Science  
1-3 Kagurazaka, Shinjuku-ku  
Tokyo 162-8601  
JAPAN

**Dr. Matteo Gallone**  
SISSA  
International School for Advanced  
Studies  
Via Beirut n. 2-4  
34014 Trieste  
ITALY

**Prof. Dr. Vladimir S. Georgiev**  
Dip. di Matematica  
Universita di Pisa  
Largo Bruno Pontecorvo, 5  
56127 Pisa  
ITALY

**Prof. Dr. Masahiro Ikeda**  
Information Science Lab.  
RIKEN Institute  
Wako  
Saitama 351-01  
JAPAN

**Dr. Jinyeop Lee**  
Mathematisches Institut  
Ludwig-Maximilians-Universität  
München  
Theresienstr. 39  
80333 München  
GERMANY

**Prof. Dr. Marilena Ligabò**  
Dipartimento di Matematica  
Universita di Bari  
70125 Bari  
ITALY

**Prof. Dr. Sandra Lucente**  
Dipartimento Interateneo di Fisica  
Università di Bari  
via amendola 173  
70125 Bari  
ITALY

**Prof. Dr. Alessandro Michelangeli**  
Institut für Angewandte Mathematik  
IAM  
& Hausdorff Center for Mathematics  
HCM  
Rheinische  
Friedrich-Wilhelms-Universität Bonn  
Endenicher Allee 60  
53115 Bonn  
GERMANY

**Prof. Dr. Diego Noja**

Associate professor of Mathematical  
Physics  
Dipartimento di Matematica e  
Applicazioni  
Università di Milano Bicocca  
Via Cozzi 55  
20125 Milano  
ITALY

**Prof. Dr. Tohru Ozawa**

No.310, Building 55N  
Department of Applied Physics  
Waseda University  
3-4-1, Okubo, Shinjuku  
Tokyo 1698555  
JAPAN

**Dr. Eugenio Pozzoli**

Mathématiques  
Université de Dijon  
B.P. 47870  
9 Avenue Savary  
21078 Dijon  
FRANCE

**Dr. Mario Rastrelli**

Dip. di Matematica "L.Tonelli"  
Università di Pisa  
Largo Bruno Pontecorvo, 5  
56127 Pisa  
ITALY

**Prof. Dr. Luca Rizzi**

SISSA  
International School for Advanced  
Studies  
via Bonomea 265  
34136 Trieste  
ITALY

**Dr. Raffaele Scandone**

Gran Sasso Science Institute  
Viale F. Crispi 7  
67100 L'Aquila  
ITALY

**Prof. Dr. Nikolay Tzvetkov**

Département de Mathématiques  
CY Cergy Paris University  
Site Saint-Martin, BP 222  
2, avenue Adolphe Chauvin  
95302 Cergy-Pontoise  
FRANCE

**Prof. Dr. Nicola Visciglia**

Dipartimento di Matematica  
Università di Pisa  
Largo Bruno Pontecorvo, 5  
56127 Pisa  
ITALY

**Prof. Dr. Ivana Vojnović**

Department of Mathematics and  
Informatics, Faculty of Sciences,  
University of Novi Sad  
Trg Dositeja Obradovica 4  
21000 Novi Sad  
SERBIA

**Prof. Dr. Jens Wirth**

Institut für Analysis, Dynamik und  
Modellierung  
Universität Stuttgart  
Pfaffenwaldring 57  
70569 Stuttgart  
GERMANY

**Prof. Dr. Kenji Yajima**

Department of Mathematics  
Faculty of Science  
Gakushuin University  
Toshimaku-ku, 1-5-1 Mejiro  
Tokyo 171-8588  
JAPAN

