



Mathematisches
Forschungsinstitut
Oberwolfach



Oberwolfach Preprints

OWP 2016 - 12

RICHARD M. ARON, FRÉDÉRIC BAYART,
PAUL M. GAUTHIER, MANUEL MAESTRE AND
VASSILI NESTORIDIS

Dirichlet Approximation and Universal Dirichlet
Series

Mathematisches Forschungsinstitut Oberwolfach gGmbH
Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

Starting in 2007, the MFO publishes a preprint series which mainly contains research results related to a longer stay in Oberwolfach. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 30 free hard copies (DIN A4, black and white copy) by surface mail.

Of course, the full copy right is left to the authors. The MFO only needs the right to publish it on its website *www.mfo.de* as a documentation of the research work done at the MFO, which you are accepting by sending us your file.

In case of interest, please send a **pdf file** of your preprint by email to *rip@mfo.de* or *owlf@mfo.de*, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany

Tel +49 7834 979 50
Fax +49 7834 979 55
Email admin@mfo.de
URL www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.
Copyright of the content is held by the authors.

DIRICHLET APPROXIMATION AND UNIVERSAL DIRICHLET SERIES

RICHARD M. ARON, FRÉDÉRIC BAYART, PAUL M. GAUTHIER, MANUEL MAESTRE, AND VASSILI NESTORIDIS

ABSTRACT. We characterize the uniform limits of Dirichlet polynomials on a right half plane. We extend the approximation theorems of Runge, Mergelyan and Vitushkin to the Dirichlet setting with respect to the Euclidean distance and to the chordal one, as well. We also strengthen the notion of Universal Dirichlet series.

1. INTRODUCTION

This paper is concerned with Dirichlet series, and in particular with approximation results on such series:

- Existence of universal (Dirichlet) series,
- Mergelyan-type and Vitushkin-type (Dirichlet) theorems, and
- Runge-type (Dirichlet) theorems.

Let us introduce some notation. As usual, we shall denote by \mathbb{C} the field of complex numbers, by $\mathbb{C}_+ = \{s \in \mathbb{C} : \operatorname{Re} s > 0\}$ and by $\mathbb{C}_- = \{s \in \mathbb{C} : \operatorname{Re} s < 0\}$. If Ω is an open subset of \mathbb{C} , then $H(\Omega)$ denotes the space of all holomorphic functions on Ω . Given K a compact subset of \mathbb{C} , we denote by $A(K)$ the space of all functions $f : K \rightarrow \mathbb{C}$ that are continuous on K and holomorphic in the interior of K . We are interested in ordinary Dirichlet series, namely series of the form $D = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$.

In Section 2, we will first review some basic properties of Dirichlet series, as well as recalling some background material on the relation of Dirichlet series and infinite dimensional holomorphy. Section 3 is devoted to extending work of the second author [3], showing the existence of a dense G_δ set of Dirichlet series D , each converging in a strong sense to any given entire function. In Section 4, we show that for certain compact subsets $K \subset \overline{\mathbb{C}}$, every $f \in A(K)$ can be uniformly approximated on K by “rational Dirichlet” polynomials. In addition to Mergelyan-type results of this form, we also prove a Runge approximation theorem for rational Dirichlet polynomials. Our results in both §3 and §4 depend on an approximation result (Theorem 3.1), in which it will be shown that on a compact set K with connected complements, Dirichlet polynomials can simultaneously uniformly approximate functions in $A(K)$ and functions from a certain class of Dirichlet series on \mathbb{C}_+ . Finally, in §5, we turn to the extension of the previous results to the situation where uniform approximation is now with respect to the chordal metric.

2010 *Mathematics Subject Classification*. Primary 30K10; Secondary 46G20, 30E10.

Key words and phrases. Universal series, Runge theorem.

The first and fourth authors were supported by MINECO MTM2014-57838-C2-2-P and Prometeo II/2013/013; the third by NSERC and Entente France-Québec. Partially supported also by the “SQuaREs” program at the American Institute of Mathematics, Palo Alto and the “Research in Pairs” program at the Mathematisches Forschungsinstitut, Oberwolfach.

2. BACKGROUND REVIEW; CLOSURE OF DIRICHLET POLYNOMIALS ON A HALF PLANE

To any Dirichlet series $D = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$, one can associate the following abscissas:

$$\sigma_c(D) = \inf\{\operatorname{Re} s : \sum_{n=1}^{\infty} \frac{a_n}{n^s} \text{ is convergent}\}.$$

$$\sigma_b(D) = \inf\{\sigma : \sum_{n=1}^{\infty} \frac{a_n}{n^s} \text{ is bounded on } \operatorname{Re} s \geq \sigma\}.$$

$$\sigma_u(D) = \inf\{\sigma : \sum_{n=1}^{\infty} \frac{a_n}{n^s} \text{ is uniformly convergent on } \operatorname{Re} s \geq \sigma\}.$$

$$\sigma_a(D) = \inf\{\operatorname{Re} s : \sum_{n=1}^{\infty} \frac{a_n}{n^s} \text{ is absolutely convergent}\}.$$

It is known that

$$\sigma_c(D) \leq \sigma_b(D) = \sigma_u(D) \leq \sigma_a(D) \leq \sigma_c(D) + 1.$$

The proof of inequalities relating $\sigma_c(D)$, $\sigma_u(D)$ and $\sigma_a(D)$ can be found in [21, Section 4.1, pag. 98] or in [8]. The equality $\sigma_b(D) = \sigma_u(D)$ was obtained by Bohr, and a proof can be found in [21, Theorem 6.2.3, pag. 145].

Recall that \mathcal{H}_{∞} is the Banach space of all Dirichlet series $\sum_{n=1}^{\infty} \frac{a_n}{n^s}$ that converge to a bounded function $D(s)$ on \mathbb{C}_+ , endowed with the supremum norm

$$\|D\|_{\infty} = \sup_{s \in \mathbb{C}_+} \left| \sum_{n=1}^{\infty} \frac{a_n}{n^s} \right|.$$

Definition 2.1. Let us denote by $\mathcal{A}(\mathbb{C}_+)$ the set of all Dirichlet series $D(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$ which are convergent on \mathbb{C}_+ and define a uniformly continuous function on that half plane.

Proposition 2.2. $\mathcal{A}(\mathbb{C}_+)$ is a closed subspace of \mathcal{H}_{∞} .

Proof. Since the uniform limit of a sequence of uniformly continuous functions is uniformly continuous, it is enough to check that $\mathcal{A}(\mathbb{C}_+)$ is a subset of \mathcal{H}_{∞} . Let $D(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s} \in \mathcal{A}(\mathbb{C}_+)$. By hypothesis there exists $\delta > 0$ such that $|D(s_1) - D(s_2)| < 1$ for every $s_1, s_2 \in \mathbb{C}_+$ such that $|s_1 - s_2| < \delta$. Let us take $k \in \mathbb{N}$ with $\frac{1}{k} < \delta$. Since $\sigma_c(D) \leq 0$, we have $\sigma_a(D) \leq 1$. Thus

$$|D(s)| = \left| \sum_{n=1}^{\infty} \frac{a_n}{n^s} \right| \leq \sum_{n=1}^{\infty} \frac{|a_n|}{n^{1+\frac{1}{k}}} := M < \infty,$$

for every $s \in \mathbb{C}$ such that $\operatorname{Re} s \geq 1 + \frac{1}{k}$. On the other hand if $0 < \operatorname{Re} s < 1 + \frac{1}{k}$, let j_0 be the smallest non-negative integer such that $\operatorname{Re} s + \frac{j_0}{k} \geq 1 + \frac{1}{k}$. Clearly $j_0 \leq k + 1$ and

$$|D(s)| \leq \sum_{j=0}^{j_0-1} \left| D\left(s + \frac{j}{k}\right) - D\left(s + \frac{j+1}{k}\right) \right| + \left| D\left(s + \frac{j_0}{k}\right) \right| \leq j_0 + M \leq k + 1 + M.$$

□

Theorem 2.3. Given $f : \mathbb{C}_+ \rightarrow \mathbb{C}$, the following are equivalent.

- (1) f is the uniform limit on \mathbb{C}_+ of a sequence of Dirichlet polynomials.
- (2) f is represented by a Dirichlet series pointwise on \mathbb{C}_+ and f is uniformly continuous on \mathbb{C}_+ .

Proof. (1) \Rightarrow (2) Let $f : \mathbb{C}_+ \rightarrow \mathbb{C}$, be the uniform limit on \mathbb{C}_+ of a sequence (P_k) of Dirichlet polynomials $P_k(s) = \sum_{n=1}^{n_k} \frac{b_{n,k}}{n^s}$. Since \mathcal{H}_∞ is a Banach space, f belongs to \mathcal{H}_∞ . Thus, f can be pointwise represented as $f(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$ and as the uniform limit of uniformly continuous functions, f too is uniformly continuous.

(2) \Rightarrow (1) Conversely, let $f(s) = \sum_{n=1}^{\infty} \frac{a_n}{n^s}$ pointwise on \mathbb{C}_+ with f uniformly continuous on \mathbb{C}_+ . Given $\varepsilon > 0$, there exists $\delta > 0$ such that if $s_1, s_2 \in \mathbb{C}_+$ satisfy $|s_1 - s_2| < \delta$, then $|f(s_1) - f(s_2)| < \varepsilon/2$. Since, by Proposition 2.2, f is bounded on \mathbb{C}_+ , we have $\sigma_u(f) = \sigma_b(f) \leq 0$. Hence, given $\delta/2$, the polynomials $P_k(s) = \sum_{n=1}^k \frac{a_n}{n^s}$ converge uniformly to f on $[\operatorname{Re} s \geq \delta/2]$. Thus if we set $Q_k(s) = P_k(s + \delta/2) = \sum_{n=1}^k \frac{a_n}{n^{\delta/2}} \frac{1}{n^s}$, then $\{Q_k(s)\}$ is a sequence of Dirichlet polynomials converging uniformly to $f(s + \delta/2)$ on \mathbb{C}_+ and we obtain

$$\left| f(s) - \sum_{n=1}^k \frac{a_n}{n^{\delta/2}} \frac{1}{n^s} \right| \leq \left| f(s) - f\left(s + \frac{\delta}{2}\right) \right| + \left| f\left(s + \frac{\delta}{2}\right) - \sum_{n=1}^k \frac{a_n}{n^{\delta/2}} \frac{1}{n^s} \right| < \frac{\varepsilon}{2} + \left\| f\left(\cdot + \frac{\delta}{2}\right) - Q_k \right\|,$$

for every $s \in \mathbb{C}_+$ and every k . Now choosing k_0 such that $\|f(\cdot + \frac{\delta}{2}) - Q_k\| \leq \frac{\varepsilon}{2}$ for $k \geq k_0$, we arrive at

$$\|f - Q_k\| \leq \varepsilon,$$

for every $k \geq k_0$. □

Let B_{c_0} denote the open unit ball in the space c_0 of sequences which converge to zero and for a function $f : B_{c_0} \rightarrow \mathbb{C}$, set $\|f\|_{B_{c_0}} = \sup\{|f(x)| : x \in B_{c_0}\}$. Endowed with this norm, the space of all functions on B_{c_0} which are uniformly continuous and complex Fréchet differentiable is a Banach algebra, which we denote by $\mathcal{A}_u(B_{c_0})$.

Theorem 2.4. $\mathcal{A}(\mathbb{C}_+)$ is isometrically isomorphic to $\mathcal{A}_u(B_{c_0})$.

Hedenmalm, Lindqvist and Seip in [16] obtained that \mathcal{H}_∞ , the Banach space of all Dirichlet series bounded and convergent on \mathbb{C}_+ is isometrically isometric to $H_\infty(B_{c_0})$, the Banach space of all bounded and complex Fréchet differentiable functions on the open unit ball of c_0 . Even though their proof makes use of other isometries, a careful analysis yields our claim, taking into consideration that B_{c_0} is denoted by $\mathbb{D}^\infty \cap c_0(\mathbb{N})$ in [16], and using our formula (2.1), below. On the other hand, in the very recent preprint [4], another proof of that isometry can be found, but again the proof of our statement is only implicit there and one needs to invoke (2.1). We present here an explicit proof of this fact since we shall need it in the proof Theorem 2.6.

Proof. Let $\mathbb{N}_0^{(\mathbb{N})}$ denote the set of sequences of non-negative integers with finite support. That is, $\mathbb{N}_0^{(\mathbb{N})}$ consists of all $\alpha = (\alpha_1, \dots, \alpha_N, 0 \dots)$ where $N \in \mathbb{N}$ and $\alpha_j \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$ for $j = 1, \dots, N$. By z^α we denote the monomial defined on $\mathbb{C}^\mathbb{N}$ by $z^\alpha = z_1^{\alpha_1} \dots z_N^{\alpha_N}$. Obviously z^α belongs to $\mathcal{A}_u(B_{c_0})$ since it is a complex Fréchet differentiable function when restricted to c_0 that is uniformly continuous on B_{c_0} . But z^α also can be considered as a monomial in N -variables defined on \mathbb{C}^N .

A function $P : c_0 \rightarrow \mathbb{C}$ is called an m -homogeneous polynomial if there exists a continuous m -linear form $A : c_0^m \rightarrow \mathbb{C}$, such that $P(x) = A(x, \dots, x)$ for every $x \in c_0$. Clearly $P \in \mathcal{A}_u(B_{c_0})$. It is known (see e.g [1]) that every function on $\mathcal{A}_u(B_{c_0})$ is the uniform limit of linear combinations of continuous homogeneous polynomials. In other words

$$\mathcal{A}_u(B_{c_0}) = \overline{\operatorname{span}\{P_m, m\text{-homogeneous polynomial} : m \in \mathbb{N}_0\}}^{\|\cdot\|_{B_{c_0}}}.$$

But in [7] it was shown that every homogeneous polynomial on c_0 is weakly uniformly continuous when restricted to the unit ball of c_0 and, by [13, Proposition 2.8, p. 90] and the fact that the

canonical basis of c_0 is shrinking, every such polynomial is the uniform limit on B_{c_0} of a sequence of polynomials of the form $P(z) = \sum_{\alpha \in J} c_\alpha z^\alpha$ where J is a finite subset of $\mathbb{N}_0^{(\mathbb{N})}$ and $c_\alpha \in \mathbb{C}$ for every $\alpha \in J$. Thus we have

$$(2.1) \quad \mathcal{A}_u(B_{c_0}) = \overline{\text{span}\{z^\alpha : \alpha \in \mathbb{N}_0^{(\mathbb{N})}\}}^{\|\cdot\|_{B_{c_0}}}.$$

The other ingredient needed for the proof is a deep result of Harald Bohr [9] (see also [21, pp. 115–117]). Let $p = (p_n)$ denote the increasing sequence of prime numbers. Given $\alpha = (\alpha_1, \dots, \alpha_k, 0 \dots) \in \mathbb{N}_0^{(\mathbb{N})}$ we set $p^\alpha = p_1^{\alpha_1} \dots p_k^{\alpha_k}$ and $N \in \mathbb{N}$, we denote by Λ_N the family of multi-indexes $\alpha \in \mathbb{N}_0^{(\mathbb{N})}$ such that $p^\alpha \leq N$. Then

$$\sup_{\text{Re } s > 0} \left| \sum_{n=1}^N \frac{a_n}{n^s} \right| = \sup_{z \in \mathbb{D}^k} \left| \sum_{\alpha \in \Lambda_N} a_{p^\alpha} z^\alpha \right|,$$

for every $a_1, \dots, a_N \in \mathbb{C}$ and all N , where the number k of variables is the subscript of the biggest prime number p_k less than or equal to N (in other words, k is the number of primes less than or equal to N). This result yields that the correspondence:

$$\frac{1}{n^s} \longmapsto z^\alpha, \quad n = p^\alpha,$$

generates a mapping

$$\phi : \text{span} \left\{ \frac{1}{n^s} : n \in \mathbb{N} \right\} \subset \mathcal{A}(C_+) \longrightarrow \text{span} \left\{ z^\alpha : \alpha \in \mathbb{N}_0^{(\mathbb{N})} \right\} \subset \mathcal{A}_u(B_{c_0}),$$

which is an isometry and is both linear and multiplicative. Thus it can be extended to a surjective isometry

$$\tilde{\phi} : \overline{\text{span} \left\{ \frac{1}{n^s} : n \in \mathbb{N} \right\}}^{\|\cdot\|_\infty} \longrightarrow \overline{\text{span} \left\{ z^\alpha : \alpha \in \mathbb{N}_0^{(\mathbb{N})} \right\}}^{\|\cdot\|_{B_{c_0}}}.$$

The conclusion follows from Theorem 2.3 which states that $\mathcal{A}(C_+) = \overline{\text{span} \left\{ \frac{1}{n^s} : n \in \mathbb{N} \right\}}^{\|\cdot\|_\infty}$, and from (2.1). □

Remark 2.5. Since we can extend any uniformly continuous function on C_+ to its closure $\overline{C_+}$, any function in $\mathcal{A}(C_+)$ extends uniformly to a uniformly continuous function on $\overline{C_+}$. However, there exist functions $f : \overline{C_+} \rightarrow \mathbb{C}$ that are uniformly continuous on $\overline{C_+}$ and pointwise representable for a certain $\sum_{n=1}^\infty \frac{a_n}{n^s}$ for every $s \in C_+$, but for which nevertheless there is some $t \in \mathbb{R}$ so that $\sum_{n=1}^\infty \frac{a_n}{n^{it}}$ does not converge. That is, in general $f(it)$ is not representable by this Dirichlet series.

Indeed, there exists $g(z) = \sum_{n=0}^\infty b_n z^n$ in the disk algebra such that $\sum_{n=0}^\infty b_n$ is not convergent. Hence, by considering g in $\mathcal{A}_u(B_{c_0})$ and taking $f = \tilde{\phi}^{-1}(g) \in \mathcal{A}(C_+)$, we have $f(s) = \sum_{l=1}^\infty b_{l-1} \frac{1}{(2^{l-1})^s}$ for every $s \in C_+$. But

$$\sum_{l=1}^\infty b_{l-1} \frac{1}{(2^{l-1})^0} = \sum_{n=0}^\infty b_n$$

does not converge and hence cannot represent the uniformly continuous unique extension of f at 0.

Given a non-empty set of indices I , we consider the compact Hausdorff space $\overline{\mathbb{D}}^I$, endowed with the product topology. Recently in [17], Makridis and Nestoridis introduced the space $A(\overline{\mathbb{D}}^I)$ of all functions $f : \overline{\mathbb{D}}^I \rightarrow \mathbb{C}$ continuous on $\overline{\mathbb{D}}^I$, endowed with the product topology, that are separately

holomorphic on \mathbb{D}^I . The space $A(\overline{\mathbb{D}}^I)$ is a Banach subalgebra of the space $C(\overline{\mathbb{D}}^I)$ of all continuous functions on $\overline{\mathbb{D}}^I$ endowed with the supremum norm. We now show that whenever I is an infinite and countable set, this space is going to be isometrically isomorphic to the space of Dirichlet series $\mathcal{A}(\mathbb{C}_+)$.

Theorem 2.6. *For $I = \mathbb{N}$, the Banach algebra $A(\overline{\mathbb{D}}^{\mathbb{N}})$ is isometrically isomorphic to $\mathcal{A}(\mathbb{C}_+)$ and also to $\mathcal{A}_u(B_{c_0})$.*

Proof. The mapping

$$\psi : \text{span}\{z^\alpha : \alpha \in \mathbb{N}_0^{(\mathbb{N})}\} \subset \mathcal{A}_u(B_{c_0}) \longrightarrow A(\overline{\mathbb{D}}^{\mathbb{N}}),$$

generated by setting $\psi(z^\alpha) = z^\alpha$, is an isometry which is into, linear and multiplicative. Hence ψ can be extended to another linear and multiplicative surjective isometry

$$\tilde{\psi} : \overline{\text{span}\{z^\alpha : \alpha \in \mathbb{N}_0^{(\mathbb{N})}\}}^{\|\cdot\|_{B_{c_0}}} \longrightarrow \overline{\text{span}\{z^\alpha : \alpha \in \mathbb{N}_0^{(\mathbb{N})}\}}^{\|\cdot\|_{C(\overline{\mathbb{D}}^I)}}.$$

But we have already shown that

$$\mathcal{A}_u(B_{c_0}) = \overline{\text{span}\{z^\alpha : \alpha \in \mathbb{N}_0^{(\mathbb{N})}\}}^{\|\cdot\|_{B_{c_0}}},$$

Moreover, by [17, Proposition 3.1], given $f \in A(\overline{\mathbb{D}}^{\mathbb{N}})$, there exists a sequence (P_n) of polynomials of the form

$$P(z) = \sum_{\alpha \in J} c_\alpha z^\alpha,$$

where J is a finite subset of $\mathbb{N}_0^{(\mathbb{N})}$, such that the sequence (P_n) converges uniformly on $\overline{\mathbb{D}}^{\mathbb{N}}$ to f . Hence,

$$A(\overline{\mathbb{D}}^{\mathbb{N}}) = \overline{\text{span}\{z^\alpha : \alpha \in \mathbb{N}_0^{(\mathbb{N})}\}}^{\|\cdot\|_{C(\overline{\mathbb{D}}^I)}}.$$

We have obtained that $A(\overline{\mathbb{D}}^{\mathbb{N}}) \stackrel{1}{\cong} \mathcal{A}_u(B_{c_0})$. Finally, by Theorem 2.4, $\mathcal{A}_u(B_{c_0})$ is isometrically isomorphic to $\mathcal{A}(\mathbb{C}_+)$. \square

Corollary 2.7. *Given a function $f : \mathbb{C} \rightarrow \mathbb{C}$, the following are equivalent.*

- (1) *On every (right) half-plane, f is the uniform limit of Dirichlet polynomials.*
- (2) *There exists a Dirichlet series $\sum_{n=1}^{\infty} a_n \frac{1}{n^s}$, which converges to f at every point of \mathbb{C} .*

Proof. (2) \Rightarrow (1). Assume that $f(s) = \sum_{n=1}^{\infty} a_n \frac{1}{n^s}$ for all points s of \mathbb{C} . Since $\sigma_a(f) \leq \sigma_c(f) + 1$ we have that $\sum_{n=1}^{\infty} |a_n| \frac{1}{n^{\text{Re } s}} < \infty$ for every s . Hence the sequence of Dirichlet polynomials $\sum_{n=1}^N a_n \frac{1}{n^s}$ converges to f uniformly on every half-plane.

(1) \Rightarrow (2). Fix k a non-positive integer. With a suitable change of variables, Theorem 2.3 implies that there exists a Dirichlet series $\sum_{n=1}^{\infty} a_n(k) \frac{1}{n^s}$ such that

$$f(s) = \sum_{n=1}^{\infty} a_n(k) \frac{1}{n^s},$$

for every s in \mathbb{C} with $\text{Re } s \geq k$. Thus all Dirichlet series of f coincide on \mathbb{C}_+ . Now the uniqueness of coefficients of a convergent Dirichlet series on \mathbb{C}_+ implies that $a_n(k) = a_n(l)$ for every non positive integers k and l and every $n \in \mathbb{N}$. Defining $a_n = a_n(k)$, we obtain $f(s) = \sum_{n=1}^{\infty} a_n \frac{1}{n^s}$, for every $s \in \mathbb{C}$. \square

3. UNIVERSAL DIRICHLET SERIES

The existence of the first universal Dirichlet series was established by the second author in [3] under the assumption that the compact sets under consideration were ‘‘admissible.’’ Further results on universal Dirichlet series can be found in [19], [12], [18]. Now, we improve most of those results by relaxing the assumption that the compact set be admissible.

For $\sigma > 0$ we denote $\|\sum_{n=1}^{\infty} a_n n^{-s}\|_{\sigma} = \sum_{n=1}^{\infty} |a_n| n^{-\sigma}$, and

$$D_a(\mathbb{C}_+) = \left\{ \sum_{n=1}^{\infty} a_n n^{-s} : \left\| \sum_{n=1}^{\infty} a_n n^{-s} \right\|_{\sigma} < \infty, \text{ for all } \sigma > 0 \right\}.$$

We endow $D_a(\mathbb{C}_+)$ with the Fréchet topology induced by the semi-norms $\|\cdot\|_{\sigma}, \sigma > 0$.

Theorem 3.1. *Let $K \subset \overline{\mathbb{C}_-}$ be compact with connected complement, $f \in D_a(\mathbb{C}_+)$, $g \in A(K)$, $\sigma > 0$ and $\varepsilon > 0$. Then there exists a Dirichlet polynomial $h = \sum_{n=1}^N a_n n^{-s}$ such that $\|h - g\|_{C(K)} < \varepsilon$ and $\|h - f\|_{\sigma} < \varepsilon$.*

The proof of this theorem, inspired by [2] and by [6, Section 11.5], needs some preparation. We first give two lemmas on entire functions of exponential type.

Lemma 3.2. *Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be an entire function of exponential type and let α be greater than the type. There exists a positive constant C depending only on f , such that if $u > 1$ with $|f(u)| > 0$ and N is an integer satisfying $N \geq e^2 \alpha(u+1)$, then there exists an interval $I \subset [u-1, u+1]$ of length $1/(2N^2)$ such that*

$$x \in I \implies |f(x)| \geq \frac{|f(u)|}{2} + C e^{-N}.$$

A proof of this lemma can be found in [21, Lemma 7.2.7].

Lemma 3.3. *Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a non-zero entire function of exponential type and let $\delta \in (0, 1)$. Assume that*

$$\limsup_{x \rightarrow +\infty} \frac{\log |f(x)|}{x} \geq 0.$$

Then

$$\sum_{n \geq 1} \frac{|f(\log n)|}{n^{1-\delta}} = +\infty.$$

Proof. Choose α greater than the type of f and $0 < \delta < 1$. Let (x_j) be a sequence of positive real numbers going to infinity such that for every $j \geq 1$, $|f(x_j)| \geq e^{-\delta x_j/2}$, $x_j > 1$ and $\lfloor x_j \rfloor^2 \geq e^2 \alpha(x_j + 1)$. Let $N_j = \lfloor x_j \rfloor^2$. By Lemma 3.2, we can find an interval $I_j = [y_j, y_j + 1/(2N_j^2)]$ contained in $[x_j - 1, x_j + 1]$ such that $x \in I_j \implies |f(x)| \geq e^{-\delta x_j/2}/2$.

By choosing x_1 sufficiently large, we may assume that

$$(3.1) \quad e^{y_j + \frac{1}{2N_j^2}} - e^{y_j} > 1.$$

Indeed,

$$e^{y_j + \frac{1}{2N_j^2}} - e^{y_j} \geq \min_{t \in [x_j - 1, x_j + 1]} \left(e^{t + \frac{1}{2N_j^2}} - e^t \right) = e^{x_j - 1 + \frac{1}{2N_j^2}} - e^{x_j - 1} \geq e^{x_j - 1 + (x_j - 1)^{-4}/2} - e^{x_j - 1} = \frac{e^{(x_j - 1)^{-4}/2} - 1}{e^{1 - x_j}},$$

which tends to $+\infty$ by l'Hôpital's rule, establishing (3.1). Now,

$$\begin{aligned}
\sum_{\log n \in I_j} \frac{|f(\log n)|}{n^{1-\delta}} &\gg \sum_{\log n \in I_j} \frac{e^{-\delta x_j/2}}{e^{(1-\delta)y_j}} \\
&\gg e^{-(1-\delta/2)y_j} \text{card}\{n : \log n \in I_j\} \\
&= e^{-(1-\delta/2)y_j} \text{card}\{n : n \in [e^{y_j}, e^{y_j + \frac{1}{2N_j^2}}]\} \\
&\gg e^{-(1-\delta/2)y_j} \left(e^{y_j + \frac{1}{2N_j^2}} - e^{y_j} \right) \quad \text{by (3.1)} \\
&\gg \frac{e^{\delta y_j/2}}{y_j^4}.
\end{aligned}$$

But this last quantity goes to infinity as j goes to infinity. \square

We also need a result from the geometry of function spaces. A proof of it can be found in [6, Lemma 11.11].

Lemma 3.4. *Let X be a locally convex topological vector space and let $(x_n)_{n \geq 1}$ be a sequence in X . Assume that $\sum_{n=1}^{+\infty} |\langle x^*, x_n \rangle| = +\infty$ for every nonzero continuous linear functional $x^* \in X^*$. Then, for every $N \in \mathbb{N}$, the set $\{\sum_{n=N}^M a_n x_n; M \geq N, |a_n| \leq 1\}$ is dense in X .*

Proof of Theorem 3.1. Let $K \subset \overline{\mathbb{C}_-}$ be compact with connected complement, let $P \in D_a(\mathbb{C}_+)$ be a Dirichlet polynomial, $g \in A(K)$, $\sigma > 0$ and $\varepsilon > 0$. We set $\delta = \sigma/2$ and we choose N bigger than the degree of P and such that

$$\sum_{n \geq N} \frac{1}{n^{1+\delta}} < \varepsilon.$$

Let ϕ be a nonzero continuous linear functional on $A(K)$. By the Hahn-Banach and the Riesz representation theorems, there exists a nonzero (complex) measure μ with support contained in K such that, for any $u \in A(K)$,

$$\langle \phi, u \rangle = \int_K u d\mu.$$

We intend to show that $\sum_{n \geq 1} |\langle \phi, n^{-s} \rangle| = +\infty$. Observe that

$$\phi(n^{-s}) = \int_K e^{-s \log n} d\mu = \mathcal{L}_\mu(\log n),$$

where \mathcal{L}_μ is the Laplace transform of μ . Observe that \mathcal{L}_μ is nonzero. Indeed, differentiating under the integral sign and evaluating at zero,

$$\mathcal{L}_\mu^{(k)}(0) = (-1)^k \langle \phi, z^k \rangle.$$

Since K has connected complement, polynomials are dense in $A(K)$ and since $\phi \neq 0$, it follows that $\mathcal{L}_\mu^{(k)}(0) \neq 0$ for some k . On the other hand, it is well-known (see for instance [6, Lemma 11.15]) that \mathcal{L}_μ is an entire function of exponential type and that

$$\limsup_{x \rightarrow +\infty} \frac{\log |\mathcal{L}_\mu(x)|}{x} \geq 0.$$

Applying Lemma 3.3, we get

$$\sum_{n \geq 1} |\langle \phi, n^{-s-1+\delta} \rangle| = \sum_{n \geq 1} \frac{|\mathcal{L}_\mu(\log n)|}{n^{1-\delta}} = +\infty.$$

By Lemma 3.4, this implies that $\{\sum_{n=N}^M a_n n^{-(1-\delta)} n^{-s}; M \geq N, |a_n| \leq 1\}$ is dense in $A(K)$. In particular, there exists $h_0 = \sum_{n=N}^M a_n n^{-(1-\delta)} n^{-s}, |a_n| \leq 1$ such that

$$\|h_0 - (g - P)\|_{C(K)} < \varepsilon.$$

Setting $h = h_0 + P$, we thus have $\|h - g\|_{C(K)} < \varepsilon$ and

$$\|h - P\|_\sigma = \|h_0\|_\sigma \leq \sum_{n=N}^{+\infty} \frac{1}{n^{1-\delta+\sigma}} < \varepsilon.$$

We have shown the theorem for the case that f is a Dirichlet polynomial. The general case follows trivially. \square

Remark 3.5. In the above theorem, translating K and reducing σ if necessary, we may in fact choose $K \subset \{s : \operatorname{Re} s < \sigma\}$. Using Cauchy's formula, we deduce that, for every compact $L \subset \overline{\mathbb{C}_-}$, for every $f \in D_a(\mathbb{C}_+)$, g entire, $\sigma, \varepsilon > 0$, and $N \in \mathbb{N}$, there exists a Dirichlet polynomial h such that

$$\sup_{0 \leq l \leq N} \|h^{(l)} - g^{(l)}\|_{C(L)} < \varepsilon \text{ and } \|h - f\|_\sigma < \varepsilon.$$

A consequence of Theorem 3.1 is the following result.

Theorem 3.6. *There exists a Dirichlet series D , absolutely convergent in \mathbb{C}_+ , with partial sums $S_N D$, such that, for every entire function g there exists a sequence (λ_n) , so that for every ℓ in $\{0, 1, 2, \dots\}$ the derivatives $S_{\lambda_n} D^{(\ell)}$ converge to $g^{(\ell)}$ uniformly on each compact subset of the closed left half plane, as n goes to infinity. The set of such Dirichlet series is dense and a G_δ in the space of Dirichlet series absolutely convergent in \mathbb{C}_+ , and it contains a dense vector subspace of it, apart from 0.*

Proof. Let $K_m = [-m, 0] \times [-m, m]$ and let (g_k) be an enumeration of all polynomials with coefficients in $\mathbb{Q} + i\mathbb{Q}$. We endow the space $H(\mathbb{C})$ of entire functions with the following distance:

$$d_m(f, g) = \sum_{l=0}^{+\infty} 2^{-l} \min\left(1, \|f^{(l)} - g^{(l)}\|_{C(K_m)}\right).$$

For $n, m, s \geq 1$, we define

$$E(n, k, m, s) = \{h \in D_a(\mathbb{C}_+) : d_m(S_n h, g_k) < 1/s\}.$$

By Remark 3.5, the set $\bigcup_n E(n, k, m, s)$ is dense and it is also clearly open in $D_a(\mathbb{C}_+)$. We claim that each D in $\bigcap_{k, m, s} \bigcup_n E(n, k, m, s)$ satisfies the conclusion of the theorem. Indeed, let g be any entire function. For each $m \geq 1$, there exists k_m such that $d_m(g, g_{k_m}) < 4^{-m}/2$. Then, since $D \in \bigcup_n E(n, k_m, m, 2 \cdot 4^m)$, we know from the triangle inequality that there exists an integer λ_m such that $d_m(S_{\lambda_m} D, g) < 4^{-m}$. By the definition of d_m , this implies that, for any $l \leq m$,

$$\|S_{\lambda_m} D^{(l)} - g^{(l)}\|_{C(K_m)} < 2^{-m}.$$

Since any compact set of $\overline{\mathbb{C}_-}$ is contained in all K_m for any m large enough, we are done. The construction of a dense vector subspace follows from the methods of [5]. \square

This result improves [5, Theorem 10] where the set K is assumed to be admissible, while in Theorem 3.6, the set K is any compact subset of $\overline{\mathbb{C}_-}$ with connected complement.

Remark 3.7. We should mention that S. Gardiner and M. Manolaki have recently been applying potential theory techniques to study properties of universal Dirichlet series in [15]. This promising approach has yielded results that apparently cannot be answered using standard complex analysis.

4. MERGELYAN AND RUNGE TYPE THEOREMS IN THE DIRICHLET SETTING

Another consequence of Theorem 3.1 is the following result.

Lemma 4.1. *Let $K \subset \mathbb{C}$ be compact with connected complement, $f \in A(K)$ and $\varepsilon > 0$. Then there exists a Dirichlet polynomial $P(s) = \sum_{n=1}^N a_n \frac{1}{n^s}$ such that $\|f - P\|_{C(K)} < \varepsilon$.*

It follows easily from Lemma 4.1 and Runge's theorem on polynomial approximation that if $K \subset \mathbb{C}$ is an arbitrary compact set, then the set of uniform limits of Dirichlet polynomials on K is the same as the set $P(K)$ of uniform limits of (algebraic) polynomials on K . The family $P(K)$ is known to consist precisely of the set of functions on K having an extension in $A(\widehat{K})$ where \widehat{K} is the complement of the unbounded component of $K^c = \mathbb{C} \setminus K$. Indeed, if a function f defined on K is the uniform limit of a sequence $\{p_n\}$ of polynomials, then the sequence $\{p_n\}$ is uniformly Cauchy on K and so, by the maximum principle it is also uniformly Cauchy on \widehat{K} . Thus, the sequence $\{p_n\}$ converges uniformly on \widehat{K} to some function \widehat{f} , which is in $A(\widehat{K})$ and whose restriction to K is f . Conversely, if f extends to a function $\widehat{f} \in A(\widehat{K})$, then, by Mergelyan's theorem, \widehat{f} is the uniform limit on \widehat{K} of a sequence of polynomials and *a fortiori* f is the uniform limit on K of the same sequence of polynomials.

Another easy consequence of Lemma 4.1 is the following result.

Theorem 4.2. *Let $\Omega \subset \mathbb{C}$ be an open simply connected set and $H(\Omega)$ be the space of holomorphic functions on Ω endowed with the topology of uniform convergence on compacta. Then, Dirichlet polynomials are dense in $H(\Omega)$.*

Mergelyan's theorem, initially valid for compact sets with connected complement, can be extended to compact sets K such that K^c has a finite numbers of components, where the approximation will not be achieved by polynomials but by rational functions (See [22, Exercise 1, Chapter 20, p.394]). In analogy with this result we prove the following one.

Theorem 4.3. *Let $\Omega \subset \mathbb{C}$ be a domain bounded by a finite number of disjoint Jordan curves. Let $\{z_1, \dots, z_N\}$ be a set containing exactly one point from every bounded component of $\overline{\Omega}^c$. Let $f \in A(\Omega)$ and $\varepsilon > 0$. Then there exist Dirichlet polynomials P_0, P_1, \dots, P_N so that the "rational Dirichlet" function*

$$R(s) = P_0(s) + P_1 \left(\frac{1}{s - z_1} \right) + \dots + P_N \left(\frac{1}{s - z_N} \right)$$

satisfies

$$\|f - R\|_{C(\overline{\Omega})} < \varepsilon.$$

Proof. Let V_j , $j = 0, \dots, N$ be the components of $(\mathbb{C} \cup \{\infty\}) \setminus \overline{\Omega}$ and $\infty \in V_0$. By the well known "Laurent Decomposition" (see for example [11]),

$$f = f_0 + f_1 + \dots + f_N$$

where $f_j \in H(V_j^c)$ for $j = 0, \dots, N$ and $f_1(\infty) = \dots = f_N(\infty) = 0$.

The complement of V_0^c is V_0 , which is connected. So by Lemma 4.1 there exists a Dirichlet polynomial P_0 so that

$$\|P_0 - f\|_{C(\bar{\Omega})} \leq \|P_0 - f\|_{C(V_0^c)} < \frac{\varepsilon}{N+1}.$$

If we do the inversion $z \rightarrow \frac{1}{z-z_j}$ ($j = 1, \dots, N$) then the image of V_j^c is a compact set with connected component. Thus, by Lemma 4.1 the function $f_j(\frac{1}{w} + z_j)$ can be approximated by a Dirichlet polynomial $P_j(w)$ so that

$$\|f_j(z) - P_j(\frac{1}{z-z_j})\|_{C(\bar{\Omega})} \leq \|f_j(z) - P_j(\frac{1}{z-z_j})\|_{C(V_j^c)} < \frac{\varepsilon}{N+1}.$$

The triangle inequality yields the result. \square

For a compact set $K \subset \mathbb{C}$, clearly the set $P(K)$ of functions $f : K \rightarrow \mathbb{C}$ which can be uniformly approximated by polynomials is contained in the class $A(K)$. Mergelyan's theorem gives a characterization of those compact sets $K \subset \mathbb{C}$ such that $P(K) = A(K)$. Let us call such sets *Mergelyan sets*. Mergelyan's theorem then states that K is a Mergelyan set if and only if K^c is connected. The analogous problem for rational approximation was solved by Vitushkin (see, for example [10]). For compact $K \subset \mathbb{C}$, denote by $R(K)$ the set of functions $f : K \rightarrow \mathbb{C}$ which are uniform limits of rational functions. Again, clearly $R(K) \subset A(K)$. Vitushkin characterized, in terms of continuous analytic capacity, those compact $K \subset \mathbb{C}$ for which $R(K) = A(K)$. Let us call such sets *Vitushkin sets*. Since rational functions are well-defined on the Riemann sphere, we could also consider uniform approximation of functions $f : K \rightarrow \mathbb{C}$, defined on compact subsets $K \subset \mathbb{C} \cup \{\infty\}$, by rational functions whose poles lie outside of K . However, for simplicity, we shall confine our study to compact subsets of \mathbb{C} .

An example of a Vitushkin set is the closure $\bar{\Omega}$ of a domain $\Omega \subset \mathbb{C}$ bounded by finitely many disjoint Jordan curves. We can use this example to formulate an analogue of the previous theorem for arbitrary Vitushkin sets. If P is a Dirichlet polynomial and $a \in \mathbb{C}$ we define the rational Dirichlet function $P_a(s) = P(1/(s-a))$ and if $a = \infty$, we set $P_\infty = P$.

Theorem 4.4. *Let $K \subset \mathbb{C}$ be a Vitushkin compactum. Let Z be a set which meets every component of $\bar{\mathbb{C}} \setminus K$. Let $f \in A(K)$ and $\varepsilon > 0$. Then, there exist finitely many points $z_1, \dots, z_n \in Z$ and Dirichlet polynomials P_{z_1}, \dots, P_{z_n} so that the "rational Dirichlet" function*

$$R(s) = P_{z_1}(s) + \dots + P_{z_n}(s)$$

satisfies

$$\|f - R\|_{C(K)} < \varepsilon.$$

Proof. By Vitushkin's theorem, there is a rational function g such that $\|f - g\|_{C(K)} < \varepsilon/2$. Since rational functions composed with Möbius transformations are again rational functions, we may assume that $K \subset \mathbb{C}$. Let $\Omega \subset \mathbb{C}$ be a domain bounded by finitely many disjoint Jordan curves, such that Ω contains K and all poles of h lie outside of $\bar{\Omega}$. By the previous theorem, there is a rational Dirichlet function R such that $\|g - R\|_{C(\bar{\Omega})} < \varepsilon/2$. The triangle inequality yields the desired approximation. \square

The usual notation for the set of uniform limits of rational functions having no poles on K is $R(K)$ and we have defined K to be a Vitushkin set if $R(K) = A(K)$. Let us denote by $RD(K)$ the uniform limits of Dirichlet rational functions having no singularities on K . We have shown that $RD(K) = A(K)$ if $R(K) = A(K)$, that is, if the compact set K is a Vitushkin compactum. In fact, this

condition is also necessary, since for the converse we may approximate Dirichlet polynomials by regular polynomials using Runge's theorem.

We now describe some examples. Theorem 4.3 asserts that the closure $\overline{\Omega}$ of a domain Ω bounded by finitely many disjoint Jordan curves is a Vitushkin set. There are many other interesting examples of Vitushkin sets. For instance, if the diameters of the complementary components of K (even though there may be infinitely many such components) are bounded away from zero or if the compact set has area measure zero, then it is a Vitushkin set.

If we consider any arbitrary domain $\Omega \subset \mathbb{C}$, then there is an exhausting family of compact subsets K_m , $m = 1, 2, \dots$ of Ω , such that every K_m is bounded by finite number of disjoint Jordan curves. Thus, K_m can be chosen as $\overline{\Omega}$ in Theorem 4.3 and Theorem 4.3 can be applied. Let A be a set containing a point in each component of $(\mathbb{C} \cup \{\infty\}) \setminus \overline{\Omega}$. (In fact let $A_m = \{z_{1,m}, \dots, z_{N_m,m}\}$ be a set containing exactly one point from each bounded component of the set K_m^c . Then it suffices to take $A = \cup_{m=1}^{\infty} A_m$.) Then one easily derives from Theorem 4.3 the following Runge type theorem.

Theorem 4.5. *Let $\Omega \subset \mathbb{C}$ be a domain and let $A = \cup_{m=1}^{\infty} A_m$ be as above. We consider $H(\Omega)$ endowed with the topology of uniform convergence on compacta. Then "rational Dirichlet" functions of the form*

$$R(s) = P_0(s) + \sum_{j=1}^N P_j \left(\frac{1}{s - z_j} \right),$$

$N \in \mathbb{N}$, $z_j \in A$ and $P_j(s) = \sum_{n=1}^{N_j} a_{n,j} \frac{1}{n^s}$, are dense in $H(\Omega)$.

In addition to considering uniform approximation by Dirichlet polynomials on compacta, one can raise the question of uniform approximation on (unbounded) closed sets. We restrict our comments to the case where the closed set is a line. Let E be a straight line in \mathbb{C} . Denote by $X(E)$ the family of uniform limits of Dirichlet polynomials on E .

1) If $E = \mathbb{R}$, it is easy to see that $X(\mathbb{R})$ coincides with the set of Dirichlet polynomials. Indeed, the only Dirichlet polynomials which are bounded on the real line are the constant polynomials. Thus, if a sequence $\{P_j\}$ of Dirichlet polynomials is uniformly Cauchy on the real line, then for some natural number n_0 , we have that $P_{n_0} - P_j$ is a constant c_j , for each $j \geq n_0$, so $P_{n_0} - P_j$ converges to a constant c . Thus, if P_j converges uniformly on the real line to a function f , then f is the Dirichlet polynomial $c - P_{n_0}$. The same holds for any line E which is not vertical.

2) If $E = i\mathbb{R}$, then by Theorem 2.3, one can see that functions in $X(i\mathbb{R})$ are the boundary values of functions in $\mathcal{A}(\mathbb{C}_+)$. That is, $\varphi \in X(i\mathbb{R})$ if and only if there exists a function $f \in \mathcal{A}(\mathbb{C}_+)$, such that

$$\varphi(i\tau) = \lim_{\sigma \rightarrow 0} f(\sigma + i\tau), \quad -\infty < \tau < +\infty.$$

5. SPHERICAL APPROXIMATION

In this section we try to extend the previous results when the uniform approximation is not meant with respect to the usual Euclidean distance but instead is meant with respect to the chordal distance χ .

We recall that the chordal distance χ is defined as follows.

$$\chi(a, b) = \begin{cases} \frac{|a-b|}{\sqrt{1+|a|^2}\sqrt{1+|b|^2}} & \text{for } a, b \in \mathbb{C}, \\ \frac{1}{\sqrt{1+|a|^2}} & \text{for } a \in \mathbb{C}, b = \infty. \end{cases}$$

Obviously $\chi(a, b) \leq |a - b|$ for $a, b \in \mathbb{C}$.

Let us start with a compact set $K \subset \mathbb{C}$ with connected complement. By Mergelyan's Theorem every Dirichlet polynomial can be approximated by polynomials uniformly in $|\cdot|$. Since $\chi(a, b) \leq |a - b|$ for all $a, b \in \mathbb{C}$, it follows that every Dirichlet polynomial can be approximated χ -uniformly on K by polynomials. Lemma 4.1 implies that every polynomial can be approximated on K by Dirichlet polynomials. Since $\chi(a, b) \leq |a - b|$, it follows that every polynomial can be approximated χ -uniformly on K by Dirichlet polynomials. Thus we have the following proposition.

Proposition 5.1. *Let $K \subset \mathbb{C}$ be a compact set with connected complement. Then the set of χ -uniform limits on K of Dirichlet polynomials coincides with the set of χ -uniform limits on K of polynomials.*

If K is a closed Jordan domain, the set of χ -uniform limits on K of polynomials is precisely the family of mappings

$$\tilde{A}(K) = \{f : K \rightarrow \mathbb{C} \cup \{\infty\} \text{ continuous, } f \equiv \infty \text{ or } f(K^\circ) \subset \mathbb{C} \text{ and } f|_{K^\circ} \text{ holomorphic}\}.$$

(See [14] and the references therein.) Combining this with the above proposition, we obtain the following result, which was proved in [20].

Theorem 5.2. *Let K be the closure of a Jordan domain. Then the set of χ -uniform limits on K of Dirichlet polynomials is precisely $\tilde{A}(K)$.*

For an arbitrary compact subset $K \subset \mathbb{C}$, let $\tilde{A}(K)$ be the family of continuous mappings $f : K \rightarrow \mathbb{C} \cup \{\infty\}$, such that for every component V of K° either $f|_V \equiv \infty$ or $f(V) \subset \mathbb{C}$ and $f|_V$ is holomorphic. In case K is a closed Jordan domain, this is consistent with our previous definition of $\tilde{A}(K)$. It is well known that if $f : K \rightarrow \mathbb{C} \cup \{\infty\}$ is a χ -uniform limit of functions holomorphic on K , then $f \in \tilde{A}(K)$. In particular, this is the case if f is a χ -uniform limit on K of Dirichlet polynomials on K (see for example [14]). The converse has been proved in the particular case where K is the closure of a Jordan domain according to Theorem 5.2, but it is open in the general case of compact sets K with connected complement (see [14]). Thus we have the

Question 5.3. *Let $K \subset \mathbb{C}$ be a compact set with K° connected. Let $f \in \tilde{A}(K)$ and $\varepsilon > 0$; is it then true that there exists a Dirichlet polynomial $P(s) = \sum_{n=1}^N a_n \frac{1}{n^s}$ such that $\chi(f(s), P(s)) < \varepsilon$ for all $s \in K$?*

By Proposition 5.1, this problem is equivalent to that of finding, for every $\varepsilon > 0$, an algebraic polynomial P such that $\chi(f(s), P(s)) < \varepsilon$ for all $s \in K$. This, in turn, is equivalent to finding, for every $\varepsilon > 0$, a rational function R , pole-free on K , such that $\chi(f(s), R(s)) < \varepsilon$ for all $s \in K$. Indeed, one direction is obvious, since every polynomial is a rational, pole-free function on K . Suppose, conversely, that there is a rational function R , pole-free on K , such that $\chi(f(s), R(s)) < \varepsilon/2$, for every $s \in K$. By Runge's theorem, there is a polynomial P such that $|R(s) - P(s)| < \varepsilon/2$, for every $s \in K$. Hence, for every $s \in K$, we have

$$\chi(f(s), P(s)) \leq \chi(f(s), R(s)) + \chi(R(s), P(s)) < \chi(f(s), R(s)) + |R(s) - P(s)| < \varepsilon,$$

which establishes the claim.

Combining Theorem 5.2 with a Laurent Decomposition ([11]), we obtain the following Theorem 5.4. We omit the proof, which is identical with the proof of Theorem 3.3 in [14], the only difference being that instead of polynomials we use Dirichlet polynomials.

Theorem 5.4. *Let $\Omega \subset \mathbb{C}$ be a domain bounded by a finite set of disjoint Jordan curves. Let $K = \overline{\Omega}$ and A be a set containing a point from every bounded component of $\mathbb{C} \setminus K$. Let $f : K \rightarrow \mathbb{C} \cup \{\infty\}$ be a continuous mapping. Then the following are equivalent.*

- (1) The function f is the χ -uniform limit on K of a sequence of functions of the form $P_0(s) + \sum_{j=1}^N P_j(\frac{1}{s-z_j})$ where P_j are Dirichlet polynomials and $z_j \in A$, $j = 1, \dots, N$, $N \in \mathbb{N}$ and P_0 is a Dirichlet polynomial.
- (2) The function f is the χ -uniform limit on K of a sequence of functions of the form $P_0(s) + \sum_{j=1}^N P_j(\frac{1}{s-z_j})$ where $z_j \in \mathbb{C} \setminus K$, $j = 1, \dots, N$, $N \in \mathbb{N}$ and all P_j are Dirichlet polynomials.
- (3) $f \in \tilde{A}(K)$.

Since every domain has an exhaustion by compact sets K_m , $m = 1, 2, \dots$ of the form of the set K in Theorem 5.4 we obtain the following Runge type theorem.

Theorem 5.5. *Let Ω be a domain in \mathbb{C} and A be a set containing ∞ and a point from each bounded component of $\mathbb{C} \setminus \Omega$. Let $f \equiv \infty$ or $f : \Omega \rightarrow \mathbb{C}$ be holomorphic. Then f is the χ -uniform limit on each compact subset of Ω of functions of the form $h = P_0(s) + \sum_{j=1}^N P_j(\frac{1}{s-z_j})$, where P_j are Dirichlet polynomials for $j = 0, \dots, N$, $z_j \in A \setminus \{\infty\}$, for $j = 1, \dots, N$ and $N \in \mathbb{N}$.*

Conversely every χ -uniform limit on each compact subset of Ω of functions of the form of h is either a constant equal to ∞ or holomorphic in Ω .

Remark 5.6. Earlier, we made some remarks concerning uniform approximation on lines by Dirichlet polynomials. Now we consider χ -uniform approximation on the real line. Denote by $X_\chi(\mathbb{R})$ the χ -uniform limits on \mathbb{R} of Dirichlet polynomials. If $f \in X_\chi(\mathbb{R})$, what can we say about f apart from the fact that it is χ -continuous? The constant function $f \equiv \infty$ can be thus approximated by the Dirichlet polynomials $D_n(s) = n$. Trivially, every Dirichlet polynomial can be so approximated. Are there other examples?

Consider the special case of Dirichlet series $D(s)$ which are χ -uniformly convergent on \mathbb{R} . We claim that $\Sigma(s) = \sum_{n=1}^\infty n^{-s}$ is such a series. It χ -converges pointwise to the function $\Sigma(\sigma) = \zeta(\sigma)$, for $\sigma > 1$ and $\Sigma(\sigma) = \infty$, for $\sigma \leq 1$. The sum $\Sigma(\sigma)$ is χ -continuous on \mathbb{R} . We wish to show that the series converges χ -uniformly. Fix $\epsilon > 0$ and choose $\sigma_\epsilon > 1$ such that $\chi(\Sigma(\sigma), \infty) < \epsilon$, for $\sigma \leq \sigma_\epsilon$. Choose N_1 such that $\chi(\sum_{n=1}^N n^{-\sigma_\epsilon}, \infty) < \epsilon$, for all $N \geq N_1$. It follows that $\chi(\sum_{n=1}^N n^{-\sigma}, \infty) < \epsilon$, for all $\sigma \leq \sigma_\epsilon$ and all $N \geq N_1$. Choose N_2 such that $|\sum_{n=1}^N n^{-\sigma} - \zeta(\sigma)| < \epsilon$, for all $N \geq N_2$ and all $\sigma \geq \sigma_\epsilon$. Now set $N_\epsilon = \max\{N_1, N_2\}$. Then, $N \geq N_\epsilon$ implies that $\chi(\sum_{n=1}^\infty n^{-\sigma}, \Sigma(\sigma)) \leq 2\epsilon$, for all $\sigma \in \mathbb{R}$. We have shown that the Dirichlet series $\Sigma(\sigma)$ converges χ -uniformly on \mathbb{R} . The same holds for every Dirichlet series $D(s)$ with non-negative coefficients. It converges χ -uniformly on all of \mathbb{R} . It is finite for $\sigma > \sigma_c(D)$ and it is ∞ for $\sigma \leq \sigma_c(D)$. This holds even if $\sigma_c(D) = \pm\infty$.

REFERENCES

- [1] Aron; B. J. Cole; T. W. Gamelin. Spectra of algebras of analytic functions on a Banach space. *J. Reine Angew. Math.* 415 (1991), 51-93.
- [2] B. Bagchi, A joint universality theorem for Dirichlet L-functions, *Math. Z.* 181 (1982), 319-334.
- [3] F. Bayart, Topological and algebraic geometry of divergence and universality, *Studia Math.*, **167**, n0. 2 (2005), 161-181.
- [4] F. Bayart, A. Defant, L. Frerick, M. Maestre and P. Sevilla-Peris, Multipliers of Dirichlet series and monomial series expansions of holomorphic functions in infinitely many variables, preprint.
- [5] F. Bayart, K-G. Grosse-Erdmann, V. Nestoridis and C. Papadimitropoulos, Abstract theory of universal series and applications, *Proc. London Math. Soc.*, **96** (3), (2008), 417-463.
- [6] F. Bayart and É. Matheron, *Dynamics of linear operators*, Cambridge Tracts in Math, vol. 179, Cambridge University Press, 2009.
- [7] W. Bogdanowicz, On the weak continuity of the polynomial functionals defined on the space c_0 , *Bull. Acad. Polon. Sci., Cl. III.* **5** (1957), 243-246.

- [8] H. Boas, The football player and the infinite series, *Notices of the AMS*, **44 no. 11**, (1997), 1430–1435.
- [9] H. Bohr, Über die Bedeutung der Potenzreihen unendlich vieler Variabeln in der Theorie der Dirichletschen Reihen $\sum \frac{a_n}{n^s}$. *Nachr. Ges. Wiss. Göttingen, Math. Phys. Kl.*, (1913), 441–488.
- [10] A. Boivin and P. M. Gauthier, Holomorphic and harmonic approximation on Riemann surfaces. *Approximation, complex analysis, and potential theory* (Montreal, QC, 2000), 107–128, NATO Sci. Ser. II Math. Phys. Chem., **37**, Kluwer Acad. Publ., Dordrecht, 2001.
- [11] G. Costakis, V. Nestoridis and I. Papadoperakis, Universal Laurent Series, *Proc. Edinb. Math. Soc.(2)* **48 no. 3**, (2005), 571–583.
- [12] O. Demanze, A. Mouze, Growth of coefficients of universal Dirichlet series, *Ann. Polon. Math.* **91** (2007), 11–23.
- [13] S. Dineen, *Complex analysis on infinite-dimensional spaces*. Springer Monographs in Mathematics. Springer-Verlag London Ltd., London, 1999.
- [14] M. Fragouloupoulou, V. Nestodiris and I. Papadoperakis, Some results on spherical approximation, *Bull. London Math. Soc.*, **45**, (2013), 1171–1180.
- [15] S. J. Gardiner and M. Manolaki, Boundary behaviour of Dirichlet series with applications to universal series, arXiv:1602.03323.
- [16] H. Hedenmalm, P. Lindqvist, and K. Seip, A Hilbert space of Dirichlet series and systems of dilated functions in $L^2(0, 1)$ *Duke Math. J.*, **86**, (1997), 1–37.
- [17] K. Makridis and V. Nestoridis, Sets of uniqueness for uniform limits of polynomials in several complex variables. *J. Math. Anal. Appl.* **432 no. 2** (2015), 994–1004.
- [18] A. Mouze, Universality and summability of Dirichlet series, *Complex Variables and Elliptic Equations*, **54** (2009), 57–70.
- [19] V. Nestoridis, C. Papadimitropoulos, Abstract theory of universal series and applications to Dirichlet series, *C.R. Math. Acad. Sci. Paris*, **341 no.9**, (2005), 539–543.
- [20] V. Nestoridis, C. Papadoperakis, A remark on the two extensions of the disc algebra and Mergelyan’s theorem, arxiv:1104.0833.
- [21] H. Queffélec and M. Queffélec, *Diophantine approximation and Dirichlet series*, HRI lecture notes, vol. 2, New Delhi, India : Hindustan Book Agency, 2013.
- [22] W. Rudin, *Real and Complex Analysis*, Third edition, McGraw-Hill Inc., 1987.

DEPARTMENT OF MATHEMATICAL SCIENCES, KENT STATE UNIVERSITY, KENT OH 44242, USA
E-mail address: aron@math.kent.edu

LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ BLAISE PASCAL, BP 10448, F-63000 CLERMONT-FERRAND, FRANCE
E-mail address: Frederic.Bayart@math.univ-bpclermont.fr

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, MONTRÉAL QC, H3C3J7, CANADA
E-mail address: gauthier@dms.umontreal.ca

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, UNIVERSIDAD DE VALENCIA, DOCTOR MOLINER 50, 46100 BURJASOT (VALENCIA), SPAIN
E-mail address: manuel.maestre@uv.es

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ATHENS, 157 84 PANEPISTEMIOPOLIS, ATHENS, GREECE
E-mail address: vnestor@math.uoa.gr