

Oberwolfach Preprints



OWP 2009 - 22

CRAIG A. TRACY AND HAROLD WIDOM

On the Distribution of a Second Class Particle in
the Asymmetric Simple Exclusion Process

Mathematisches Forschungsinstitut Oberwolfach gGmbH
Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

Starting in 2007, the MFO publishes a preprint series which mainly contains research results related to a longer stay in Oberwolfach. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 30 free hard copies (DIN A4, black and white copy) by surface mail.

Of course, the full copy right is left to the authors. The MFO only needs the right to publish it on its website *www.mfo.de* as a documentation of the research work done at the MFO, which you are accepting by sending us your file.

In case of interest, please send a **pdf file** of your preprint by email to *rip@mfo.de* or *owlf@mfo.de*, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany

Tel +49 7834 979 50
Fax +49 7834 979 55
Email admin@mfo.de
URL www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.
Copyright of the content is held by the authors.

July 20, 2009

On the Distribution of a Second Class Particle in the Asymmetric Simple Exclusion Process

Craig A. Tracy

*Department of Mathematics
University of California
Davis, CA 95616, USA
email: tracy@math.ucdavis.edu*

Harold Widom

*Department of Mathematics
University of California
Santa Cruz, CA 95064, USA
email: widom@ucsc.edu*

Abstract

We give an exact expression for the distribution of the position $X(t)$ of a single second class particle in the asymmetric simple exclusion process (ASEP) where initially the second class particle is located at the origin and the first class particles occupy the sites $\mathbb{Z}^+ = \{1, 2, \dots\}$.

1 Introduction

The asymmetric simple exclusion process (ASEP) [2, 3] is one of the simplest models of nonequilibrium statistical mechanics and has been called the “default stochastic model for transport phenomena” [8]. A useful concept in exclusion processes is that of a *second class particle*:¹

Imagine that the particles in the system are each called either first class or second class. The evolution is the same as before, except that if a second class particle attempts to go to a site occupied by a first class particle, it is not allowed to do so, while if a first class particle attempts to move to a site occupied by a second class particle, the two particles exchange positions. In other words, a first class particle has priority over a second class particle. This rule has no effect on whether or not a given site is occupied at a given time. The advantage, though, is that viewed by itself, the collection of first class particles is Markovian, and has the same law as the exclusion process. The collection of second class particles is clearly not Markovian. However, the collection of first and second class particles is Markovian, and again evolves like an exclusion process.

Here we consider ASEP on the integer lattice \mathbb{Z} with jumps one step to the right with rate p and jumps one step to the left with rate $q = 1 - p$. We assume a leftwards drift, i.e. $q > p$. We further assume that the system has one second class particle initially located at the origin and first class particles initially located at sites in

$$Y = \{0 < y_1 < y_2 < \dots\} \subset \mathbb{Z}^+.$$

With the above initial condition, we denote by $X(t)$ the position of the second class particle at time t . The purpose of this note is to give an exact expression for the probability that the second class particle is at position x at time t , i.e. $\mathbb{P}_Y(X(t) = x)$. (The subscript Y denotes the sites of the initial configuration of the first class particles.) Our main result is for $Y = \mathbb{Z}^+$ and is given below in (9) and in a slightly different form in (11).

¹The following quote is taken from Liggett [3].

2 A Basic Lemma

The single second class particle located at $X(t)$ can be viewed as the (single) discrepancy under *basic coupling* between two asymmetric simple exclusion processes η_t and ζ_t where $\zeta_t(X(t)) = 1$ and $\eta_t(X(t)) = 0$ and initially $\{x : \zeta_0(x) = 1\} = Y' = \{0\} \cup Y$ and $\{x : \eta_0(x) = 1\} = Y$ [2, 3].

The following identity we first learned from H. Spohn [5] but presumably it has a long history:

$$\mathbb{P}_Y(X(t) = x) = \mathbb{P}_{Y'}(\zeta_t(x) = 1) - \mathbb{P}_Y(\eta_t(x) = 1). \quad (1)$$

For the convenience of the reader, we give a short proof of (1). Let ζ_t and η_t be as above evolving together under the basic coupling [2, 3]. Recall that the coupled processes satisfy $\eta_t \leq \zeta_t$ for all $t > 0$ since they satisfy this inequality at $t = 0$ [2, 3].² Define

$$\begin{aligned} \mathcal{J}_\eta(x, t) &:= \sum_{z \leq x} \eta_t(z) = \text{number of particles in configuration } \eta_t \text{ with positions } \leq x, \\ \mathcal{J}_\zeta(x, t) &:= \sum_{z \leq x} \zeta_t(z) = \text{number of particles in configuration } \zeta_t \text{ with positions } \leq x, \\ \mathcal{I}(x, t) &= \begin{cases} 1 & \text{if } X(t) \leq x, \\ 0 & \text{if } X(t) > x. \end{cases} \end{aligned}$$

By counting

$$\mathcal{J}_\zeta(x, t) = \mathcal{J}_\eta(x, t) + \mathcal{I}(x, t). \quad (2)$$

Since

$$\begin{aligned} \mathbb{E}_{Y'}(\mathcal{J}_\zeta(x, t)) &= \sum_{z \leq x} \mathbb{E}_{Y'}(\zeta_t(z)) = \sum_{z \leq x} \mathbb{P}_{Y'}(\zeta_t(z) = 1), \\ \mathbb{E}_Y(\mathcal{J}_\eta(x, t)) &= \sum_{z \leq x} \mathbb{E}_Y(\eta_t(z)) = \sum_{z \leq x} \mathbb{P}_Y(\eta_t(z) = 1), \\ \mathbb{E}_Y(\mathcal{I}(x, t)) &= \mathbb{P}_Y(X(t) \leq x) = \sum_{z \leq x} \mathbb{P}_Y(X(t) = z), \end{aligned}$$

the expectation of (2) gives

$$\sum_{z \leq x} \mathbb{P}(X(t) = z) = \sum_{z \leq x} \mathbb{P}(\zeta_t(z) = 1) - \sum_{z \leq x} \mathbb{P}(\eta_t(z) = 1)$$

²Given two configurations $\eta, \zeta \in \{0, 1\}^{\mathbb{Z}}$ we say $\eta \leq \zeta$ if $\eta(x) \leq \zeta(x)$ for all $x \in \mathbb{Z}$.

from which (1) follows.

3 Probability for a site to be occupied in ASEP

For ASEP with particles initially at Y we denote by $x_m(t)$ the position of the m th left-most particle at time t (so $x_m(0) = y_m$). In Theorem 5.2 of [6] the authors gave an exact expression for $\mathbb{P}_Y(x_m(t) = x)$. To state this result we first recall the definition of the τ -binomial coefficients. For $0 \leq \tau := p/q < 1$ define for each $n \in \mathbb{Z}^+$

$$[n] = \frac{1 - \tau^n}{1 - \tau}, \quad [n]! = [n][n-1] \cdots [1], \quad [0]! := 1, \quad \begin{bmatrix} n \\ k \end{bmatrix} = \frac{[n]!}{[k]![n-k]!}, \quad 0 \leq k \leq n,$$

and if $k > n$ we set $\begin{bmatrix} n \\ k \end{bmatrix} = 0$. Equation (5.12) of [6] can be written in the following way^{3,4}

$$\mathbb{P}_Y(x_m(t) = x) = \sum_{k=1}^{|Y|} \sum_{\substack{S \subset Y \\ |S|=k}} c_{m,k} \tau^{\sigma(S,Y)} \int_{\mathcal{C}_R} \cdots \int_{\mathcal{C}_R} I(x, k, \xi) \prod_{i=1}^k \xi_i^{-s_i} d^k \xi \quad (3)$$

where, if $S := \{s_1, \dots, s_k\}$ then

$$\begin{aligned} c_{m,k} &= q^{k(k-1)/2} (-1)^{m+1} \tau^{m(m-1)/2} \tau^{-km} \begin{bmatrix} k-1 \\ k-m \end{bmatrix}, \\ \sigma(S, Y) &= \# \{(s, y) : s \in S, y \in Y, \text{ and } y \leq s\} \\ &= \text{sum of the positions of the elements of } S \text{ in } Y, \\ I(x, k, \xi) &= \prod_{1 \leq i < j \leq k} \frac{\xi_j - \xi_i}{p + q \xi_i \xi_j - \xi_i} \left(1 - \prod_{i=1}^k \xi_i \right) \prod_{i=1}^k \frac{\xi_i^{x-1} e^{\varepsilon(\xi_i)t}}{1 - \xi_i}, \\ \varepsilon(\xi) &= \frac{p}{\xi} + q\xi - 1 \end{aligned}$$

³We make some changes in the notation in (5.12) of [6]. The (p, q) -binomial coefficient $\begin{bmatrix} n \\ k \end{bmatrix}$ of [6] equals $q^{k(n-k)}$ times the τ -binomial coefficient $\begin{bmatrix} n \\ k \end{bmatrix}$ defined above. The second change is a little more subtle. The sum in (5.12) is over all finite subsets $S \subset \{1, 2, \dots, |Y|\}$ with $|S| \geq m$. If $S = \{s_1, \dots, s_k\}$ the subset $Y_S := \{y_{s_1}, \dots, y_{s_k}\}$ and the factor $\prod_{i \in S} \xi_i^{-y_i}$ appears in the integrand of (5.12). Thus we can equivalently sum over all finite subsets $S \subset Y$ where now the factor $\prod_{1 \leq i \leq k} \xi_i^{-s_i}$ appears in the integrand. The factor $\sigma(S) = \sum_{i \in S} i$ of (5.12) becomes $\sigma(S, Y)$ given above.

⁴All contour integrals are to be given a factor of $1/2\pi i$.

and \mathcal{C}_R is a circle of radius R centered at the origin with $R \gg 1$ so that all (finite) singularities of the integrand are enclosed by \mathcal{C}_R . Observe that $c_{m,k} = 0$ when $m > k$.

Since

$$\mathbb{P}_Y(\eta_t(x) = 1) = \sum_{m=1}^{|Y|} \mathbb{P}_Y(x_m(t) = x), \quad (4)$$

we sum the right side of (3) over all $m \leq k$. To carry out this sum recall the τ -binomial theorem

$$\sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix} (-1)^j z^j \tau^{j(j-1)/2} = (1-z)(1-z\tau) \cdots (1-z\tau^{n-1}).$$

Using this a simple calculation shows

$$\sum_{m=1}^k (-1)^{m+1} \tau^{m(m-1)/2} \tau^{-km} \begin{bmatrix} k-1 \\ k-m \end{bmatrix} = (-1)^{k+1} \tau^{-k(k+1)/2} \prod_{j=1}^{k-1} (1-\tau^j).$$

Thus

$$\begin{aligned} \mathbb{P}_Y(\eta_t(x) = 1) &= \sum_{k=1}^{|Y|} (-1)^{k+1} q^{k(k-1)/2} \tau^{-k(k+1)/2} \prod_{j=1}^{k-1} (1-\tau^j) \\ &\quad \times \sum_{\substack{S \subset Y \\ |S|=k}} \tau^{\sigma(S,Y)} \int_{\mathcal{C}_R} \cdots \int_{\mathcal{C}_R} I(x, k, \xi) \prod_{i=1}^k \xi_i^{-s_i} d^k \xi. \end{aligned} \quad (5)$$

REMARK: The above formula holds for either $|Y|$ finite or infinite. For $|Y| = N$, the integral of order N in (5) is gotten from the summand $S = Y$. Since $\sigma(Y, Y) = N(N+1)/2$, we get for the coefficient of this integral

$$(-1)^{N+1} q^{N(N-1)/2} \prod_{j=1}^{N-1} = (-1)^{N+1} \prod_{j=1}^{N-1} (q^j - p^j). \quad (6)$$

4 Probability for a site to be occupied by a second class particle

As above, suppose that our initial configuration consists of a second class particle at site 0 and first class particles at sites in Y . As above, set $Y' = Y \cup \{0\}$. The process ζ_t has

initially its particles at sites in Y' . We apply formula (5) to the initial configurations Y' and Y and by (1) we subtract to obtain $\mathbb{P}_Y(X(t) = x)$. If $|Y'| = N$ there is one N -dimensional integral that comes from the expansion of $\mathbb{P}_{Y'}(\zeta_t(x) = 1)$ when $S = Y'$. The coefficient of the integral of highest order equals (6).

We now consider the special case of *step initial condition*; that is, $Y = \mathbb{Z}^+$, and use Corollary (5.13) of [6] to obtain a more compact expression for $\mathbb{P}_{\mathbb{Z}^+}(x_m(t) = x)$. To find $\mathbb{P}_{\mathbb{Z}^+}(\eta_t(x) = 1)$ we again apply (4) but use (5.13) of [6]. As above we interchange the sums over k and m , use the τ -binomial theorem ([4], pg. 26), to conclude

$$\mathbb{P}_{\mathbb{Z}^+}(\eta_t(x) = 1) = - \sum_{k \geq 1} \frac{q^{k^2}}{k!} \prod_{j=1}^{k-1} (1 - \tau^j) \int_{\mathcal{C}_R} \cdots \int_{\mathcal{C}_R} \tilde{J}_k(x, \xi) d\xi_1 \cdots d\xi_k \quad (7)$$

where

$$\tilde{J}_k(x, \xi) = \prod_{i \neq j} \frac{\xi_j - \xi_i}{p + q\xi_i\xi_j - \xi_i} (1 - \prod_i \xi_i) \prod_i \frac{\xi_i^{x-1} e^{\varepsilon(\xi_i)t}}{(1 - \xi_i)(q\xi_i - p)}.$$

We can get the corresponding formula for $Y' = \mathbb{Z}^+ \cup \{0\}$ by observing that there is a one-one correspondence between subsets $S' \subset Y'$ and subsets $S \subset Y$ given by $S = S' + 1$. Then $\sigma(S', Y') = \sigma(S, Y)$ and, with obvious notation, $\prod \xi_i^{-s_i'} = \prod \xi_i \cdot \prod \xi_i^{-s_i}$. It follows that for the difference $\mathbb{P}_{Y'}(\zeta_t(x) = 1) - \mathbb{P}_Y(\eta_t(x) = 1)$ we multiply the integrand $\tilde{J}_k(x, \xi)$ in (7) by $\prod \xi_i - 1$.

Thus

$$\mathbb{P}_{\mathbb{Z}^+}(X(t) = x) = \sum_{k \geq 1} \frac{q^{k^2}}{k!} \prod_{j=1}^{k-1} (1 - \tau^j) \int_{\mathcal{C}_R} \cdots \int_{\mathcal{C}_R} \tilde{J}_k(x, \xi) d\xi_1 \cdots d\xi_k \quad (8)$$

where

$$\tilde{J}_k(x, \xi) = \prod_{i \neq j} \frac{\xi_j - \xi_i}{p + q\xi_i\xi_j - \xi_i} (1 - \prod_i \xi_i)^2 \prod_i \frac{\xi_i^{x-1} e^{\varepsilon(\xi_i)t}}{(1 - \xi_i)(q\xi_i - p)}.$$

From this it follows that the distribution function is (on \mathcal{C}_R , $|\xi^{-1}| \ll 1$)

$$\mathbb{P}_{\mathbb{Z}^+}(X(t) \leq x) = \sum_{k \geq 1} \frac{q^{k^2}}{k!} \prod_{j=1}^{k-1} (1 - \tau^j) \int_{\mathcal{C}_R} \cdots \int_{\mathcal{C}_R} J_k(x, \xi) d\xi_1 \cdots d\xi_k \quad (9)$$

where

$$J_k(x, \xi) = \prod_{i \neq j} \frac{\xi_j - \xi_i}{p + q\xi_i\xi_j - \xi_i} (\prod_i \xi_i - 1) \prod_i \frac{\xi_i^x e^{\varepsilon(\xi_i)t}}{(1 - \xi_i)(q\xi_i - p)}.$$

Since

$$\frac{1}{p + q\xi\xi' - \xi} = \frac{1}{\xi(\xi' - 1)} + O(\tau), \quad \tau \rightarrow 0,$$

the TASEP limit of $J_k(x, \xi)$ is

$$J_k^{\text{TASEP}}(x, \xi) := \lim_{\tau \rightarrow 0} J_k(x, \xi) = \prod_{i \neq j} (\xi_j - \xi_i) \left(\prod \xi_i - 1 \right) \prod_i \frac{\xi_i^x e^{\varepsilon(\xi_i)t}}{(\xi_i(1 - \xi_i))^k}$$

where now $\varepsilon(\xi) = \xi - 1$; and hence,

$$\lim_{\tau \rightarrow 0} \mathbb{P}_{\mathbb{Z}^+}(X(t) \leq x) = \sum_{k \geq 1} \frac{1}{k!} \int_{\mathcal{C}_R} \cdots \int_{\mathcal{C}_R} J_k^{\text{TASEP}}(x, \xi) d\xi_1 \cdots d\xi_k. \quad (10)$$

Expression (9) for the distribution function can be simplified somewhat. Define the kernel

$$K_{x,t}(\xi, \xi') = q \frac{(\xi')^x e^{\varepsilon(\xi')t}}{p + q\xi\xi' - \xi},$$

and the associated operator $K_{x,t}$ on $L^2(\mathcal{C}_R)$ by

$$f(\xi) \longrightarrow \int_{\mathcal{C}_R} K_{x,t}(\xi, \xi') f(\xi') d\xi', \quad \xi \in \mathcal{C}_R.$$

Then using the identity [7]

$$\det \left(\frac{1}{p + q\xi_i \xi_j - \xi_i} \right)_{1 \leq i, j \leq k} = (-1)^k (pq)^{k(k-1)/2} \prod_{i \neq j} \frac{\xi_j - \xi_i}{p + q\xi_i \xi_j - \xi_i} \prod_i \frac{1}{(1 - \xi_i)(q\xi_i - p)}$$

we have

$$\begin{aligned} \mathbb{P}_{\mathbb{Z}^+}(X(t) \leq x) &= \sum_{k \geq 1} \tau^{-k(k-1)/2} \prod_{j=1}^{k-1} (1 - \tau^j) \times \\ &\quad \frac{(-1)^k}{k!} \int_{\mathcal{C}_R} \cdots \int_{\mathcal{C}_R} \left[\det (K_{x+1,t}(\xi_i, \xi_j))_{1 \leq i, j \leq k} - \det (K_{x,t}(\xi_i, \xi_j))_{1 \leq i, j \leq k} \right] \\ &= \sum_{k \geq 1} \tau^{-k(k-1)/2} \prod_{j=1}^{k-1} (1 - \tau^j) \int_{\mathcal{C}_R} \frac{1}{\lambda^{k+1}} [\det(I - \lambda K_{x+1,t}) - \det(I - \lambda K_{x,t})] d\lambda \end{aligned} \quad (11)$$

where $\det(I - \lambda K_{x,t})$ is the Fredholm determinant and the last line follows from the Fredholm expansion.

REMARKS:

1. One cannot interchange the sum and the integration in (11) as was possible in an analogous calculation in [7]. This is the case even though (9) converges absolutely for all $0 \leq \tau \leq 1$ (recall one may take $R \gg 1$). Thus we do not have a representation of $\mathbb{P}_{\mathbb{Z}^+}(X(t) \leq x)$ as a single integral whose integrand involves the above Fredholm determinants as was the case in [7].
2. ASEP with first and second class particles is integrable in the sense that the Yang-Baxter equations are satisfied [1]. Using this integrable structure, it is possible to compute directly, i.e. without using the basic lemma (1), $\mathbb{P}_{\mathbb{Z}^+}(X(t) = x)$ using methods similar to that of [6]. We have carried this out to the extent that (6) was computed by this approach. However, this route is much more involved than the one presented here.

ACKNOWLEDGEMENTS: The authors wish to thank Professor Dr. Herbert Spohn for communicating to them the basic lemma. This research was done during a stay within the Research in Pairs Program at the Mathematisches Forschungsinstitut Oberwolfach from May 17 – May 30, 2009. The authors thank Professor Dr. Gert-Martin Greul and the staff for their hospitality during the authors' stay. This work was supported by the National Science Foundation under grants DMS-0553379 (first author) and DMS-0552388 (second author).

References

- [1] Alcaraz, F.C., Bariev, R.Z.: Exact solution of asymmetric diffusions with second-class particles of arbitrary size. *Braz. J. Phys.* **30** (2000), 13–26.
- [2] Liggett, T.M.: *Interacting Particle Systems*. Berlin, Springer-Verlag, 2005 [Reprint of the 1985 Edition].
- [3] Liggett, T.M.: *Stochastic Interacting Systems: Contact, Voter and Exclusion Processes*. Berlin, Springer-Verlag, 1999.
- [4] Macdonald, I.G.: *Symmetric Functions and Hall Polynomials*, Oxford, Clarendon Press (1995).
- [5] Spohn, H.: private communication, May 30, 2009.

- [6] Tracy, C.A., Widom, H.: Integral formulas for the asymmetric simple exclusion process. Commun. Math. Phys. **279** (2008), 815–844.
- [7] Tracy, C.A., Widom, H.: A Fredholm determinant representation in ASEP. J. Stat. Phys. **132** (2008), 291–300.
- [8] Yau, H.-T: $(\log t)^{2/3}$ law of the two dimensional asymmetric simple exclusion process. Ann. Math. **159** (2004), 377–405.