


Oberwolfach Preprints



OWP 2009 - 02

P. BLASIAK, G. H. E. DUCHAMP, A. HORZELA, K. A.
PENSON AND A. I. SOLOMON

HEISENBERG-WEYL ALGEBRA REVISITED:
COMBINATORICS OF WORDS AND PATHS

Mathematisches Forschungsinstitut Oberwolfach gGmbH
Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

Starting in 2007, the MFO publishes a preprint series which mainly contains research results related to a longer stay in Oberwolfach. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 30 free hard copies (DIN A4, black and white copy) by surface mail.

Of course, the full copy right is left to the authors. The MFO only needs the right to publish it on its website *www.mfo.de* as a documentation of the research work done at the MFO, which you are accepting by sending us your file.

In case of interest, please send a **pdf file** of your preprint by email to *rip@mfo.de* or *owlf@mfo.de*, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany

Tel +49 7834 979 50
Fax +49 7834 979 55
Email admin@mfo.de
URL www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.
Copyright of the content is hold by the authors.

Heisenberg–Weyl Algebra Revisited: Combinatorics of Words and Paths

P Blasiak^a, G H E Duchamp^b, A Horzela^a, K A Penson^c
and A I Solomon^{c,d}

^a H. Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences
ul. Radzikowskiego 152, PL 31342 Kraków, Poland

^b LIPN, UMR 7030, Université Paris 13 - CNRS
99, av. J-B Clément, F-93430 Villetaneuse, France

^c Laboratoire de Physique Théorique de la Matière Condensée
Université Pierre et Marie Curie, CNRS UMR 7600
Tour 24 - 2ième ét., 4 pl. Jussieu, F 75252 Paris Cedex 05, France

^d The Open University, Physics and Astronomy Department
Milton Keynes MK7 6AA, United Kingdom

E-mail:

pawel.blasiak@ifj.edu.pl, andrzej.horzela@ifj.edu.pl,
ghed@lipn-univ.paris13.fr, penson@lptl.jussieu.fr,
a.i.solomon@open.ac.uk

Abstract. The Heisenberg–Weyl algebra, which underlies virtually all physical representations of Quantum Theory, is considered from the combinatorial point of view. We provide a concrete model of the algebra in terms of paths on a lattice with some decomposition rules. We also discuss the rook problem on the associated Ferrers board; this is related to the calculus in the normally ordered basis. From this starting point we explore a combinatorial underpinning of the Heisenberg–Weyl algebra, which offers novel perspectives, methods and applications.

1. Introduction

From a modern viewpoint the formalism and structure of Quantum Theory is founded on the theory of Hilbert space [1, 2]. The physical content of the theory consists of representing physical quantities as operators which satisfy some algebraic relations. Virtually all correspondence schemes come endowed with the Heisenberg–Weyl algebra structure, be it the canonical quantization scheme, the occupation number representation in quantum mechanics, or the second quantization formalism of quantum field theory. This derives from the analogy with classical mechanics whose Poissonian structure is reflected in the commutator of position and momentum observables [3]. Ubiquitous and profound, the Heisenberg–Weyl algebra has become the hallmark of non-commutativity in Quantum Theory.

An exemplary model of the Heisenberg–Weyl algebra involves combinations of derivative D , multiplication X and identity I operators acting on the space of polynomials. Physical examples are the position \hat{x} and momentum \hat{p} operators in the space of square integrable functions, or the annihilation a and creation a^\dagger operators in Fock space. Here, without loss of generality, we conform to the notation $\{a, a^\dagger\}$ for the generators of the (associative) algebra \ddagger , satisfying

$$aa^\dagger = a^\dagger a + I, \tag{1}$$

where I is the multiplicative identity. We shall be interested in combinatorial aspects of this relation and discuss one of the ensuing models of the Heisenberg–Weyl algebra.

Indeed, the combinatorial properties of Eq.(1) were recognized early and successfully applied to the domain of algebraic enumeration, principally concerning the action of the operators X and D on generating series. From this point of view these operators are auxiliary constructions facilitating enumeration of discrete structures [4, 5, 6, 7, 8, 9, 10].

However, in this note we adopt another approach, which leads towards a combinatorial model of the algebra itself. Starting from the definition of the Heisenberg–Weyl algebra as the algebra of words in a and a^\dagger supplemented by the relation of Eq. (1) we will recast it in the language of paths on a lattice with some decomposition rules. We shall also consider a convenient choice of basis, here taken to be normally ordered monomials, which permits a direct link to the combinatorics of words and the related algebra of paths. In this way algebraic problems may be expressed in the more concrete form of the decomposition and enumeration of paths. For illustration we consider the normal ordering procedure which reduces to the familiar rook problem on the Ferrers board and then derive the structure constants of the algebra by a simple combinatorial argument.

\ddagger We do not attach much weight to this particular realization, however, as we shall study algebraic properties only, for which the underlying Fock space plays no role. Our considerations hold true for any representation of the Heisenberg–Weyl algebra.

2. Heisenberg–Weyl algebra

In this note we consider an *algebra* \mathcal{A} to be a linear vector space over a field \mathbb{K} with bilinear multiplication law

$$\mathcal{A} \times \mathcal{A} \ni (b, c) \longrightarrow bc \in \mathcal{A} \quad (2)$$

which is associative and possesses a unit element I . More precisely, it is called an *associative algebra with unit* as distinct from an algebraic structure lacking associativity or a unit (*e.g.* Lie algebra). A *basis* of an algebra is a basis for its vector space structure. Each basis $(b_i)_{i \in \Lambda}$ defines a unique family $\gamma_{ij}^k \in \mathbb{K}$ such that for every ordered pair $(i, j) \in \Lambda \times \Lambda$ the set of $k \in \Lambda$ such that $\gamma_{ij}^k \neq 0$ is finite and

$$b_i b_j = \sum_{k \in \Lambda} \gamma_{ij}^k b_k. \quad (3)$$

The γ_{ij}^k are called the *structure constants* of the algebra \mathcal{A} with respect to the basis $(b_i)_{i \in \Lambda}$, from which the multiplication law can be uniquely recovered.

The *Heisenberg–Weyl algebra*, denoted by \mathcal{H} , is the algebra generated by a, a^\dagger , satisfying the relation of Eq. (1). Elements of the algebra $A \in \mathcal{H}$ are linear combinations of finite products of a and a^\dagger of the form

$$A = \sum_{\mathbf{r}, \mathbf{s}} \alpha_{\mathbf{r}, \mathbf{s}} a^{\dagger r_1} a^{s_1} a^{\dagger r_2} a^{s_2} \dots a^{\dagger r_k} a^{s_k}, \quad (4)$$

where $\mathbf{r} = (r_1, \dots, r_k)$ and $\mathbf{s} = (s_1, \dots, s_k)$ are nonnegative integer multi-indexes (with the convention $a^0 = a^{\dagger 0} = I$). This representation is ambiguous, however, due to the commutation relation Eq. (1) which yields different representations of the same element of the algebra, *e.g.* $aa^\dagger = a^\dagger a + I$. The problem can be resolved by fixing the preferred order of the generators a and a^\dagger . Conventionally, it is done by choosing the *normally ordered* form in which all annihilators stand to the right of creators. In this case, each element of the algebra \mathcal{H} is uniquely written in the normally ordered form as

$$A = \sum_{r, s} \beta_{rs} a^{\dagger r} a^s. \quad (5)$$

Hence the normally ordered monomials $a^{\dagger r} a^s$ constitute a natural basis for the Heisenberg–Weyl algebra

$$\text{BASIS OF } \mathcal{H} : \quad b_{(r, s)} = a^{\dagger r} a^s, \quad (6)$$

indexed by pairs of integers $r, s = 0, 1, 2, \dots$, and Eq. (5) is the expansion of the element A in this basis. We should note that the normally ordered representation of the elements of the algebra suggests itself as the simplest one [11]. It is important and commonly used in practical applications in quantum optics [12, 13, 14] or quantum field theory [15, 16]. Working in this basis entails the reshuffling of a and a^\dagger to the normally ordered form, which in general is a nontrivial task [17]. This brings up the issue of efficient calculation methods and intuitive schemes providing insight into the ordering procedure itself. Below, we provide a combinatorial model for the Heisenberg–Weyl algebra and then propose a resolution of the problem from this starting-point.

3.2. Normal ordering procedure

It has been shown [18, 19, 20] that the normal ordering of a word \mathbf{w} in a and a^\dagger satisfying Eq. (1) reduces to the rook problem on the associated Ferrers board $\mathcal{B}_{\mathbf{w}}$. Namely,

$$\mathbf{w} = \sum_{k \geq 0} r_{\mathcal{B}_{\mathbf{w}}}(k) \mathbf{w}^{(k)} \quad (11)$$

where $\mathbf{w}^{(k)}$ are normally ordered monomials $a^{\dagger r} a^s$ obtained from \mathbf{w} by crossing out k pairs of a and a^\dagger and then reshuffling the rest as if they were commuting variables (called the double dot operation used in quantum field theory). For example, for a word in Eq. (7) we have $\mathbf{w}^{(0)} = a^{\dagger 5} a^4$, $\mathbf{w}^{(1)} = a^{\dagger 4} a^3$, $\mathbf{w}^{(2)} = a^{\dagger 3} a^2$, etc., and hence its normally ordered form reads (see Eq. (8))

$$\mathbf{w} = a^{\dagger 5} a^4 + 10 a^{\dagger 4} a^3 + 23 a^{\dagger 3} a^2 + 9 a^{\dagger 2} a . \quad (12)$$

A rigorous proof of Eq. (11) relies on the observation that each word can be reduced to the sum of two simpler ones by choosing the places in which a precedes a^\dagger (which correspond to step-forming cells of the previous section) and reshuffling them according to Eq. (1), *i.e.* $aa^\dagger \rightarrow I + a^\dagger a$. For example, for a word of Eq. (7) there are three choices which exactly correspond to possible decompositions of the associated Ferrers diagram $\mathcal{B}_{\mathbf{w}}$ in Fig. 2. We note that although there are various possible decomposition schemes it can be shown that the result is unique.

In short, the normal ordering of a word reduces to the enumeration of possible non-capturing rook arrangements on the associated Ferrers board. The problem can be systematically handled by successive decompositions of the board. Moreover, one can devise simple algorithms based on the recursive rule given in Eq. (10). The methods described in this paper may be extended to the q -deformed case (see *e.g.* [21, 20]).

3.3. Combinatorial realization of the Heisenberg–Weyl algebra

We observed in Sect. 3.1 that each word in two letters, here taken as a and a^\dagger , can be encoded as a path, see Fig. 1. This establishes an isomorphism between the algebra \mathcal{W} of words in two letters, and the algebra \mathcal{P} of paths. In \mathcal{W} multiplication is given by simple concatenation of words with the unit being the void word, while in \mathcal{P} multiplication is given by concatenation of paths. In both cases we shall indicate the unit, which is the void word or path respectively, by the symbol \emptyset . Both algebras are free. The Heisenberg–Weyl algebra arises by imposing on \mathcal{W} the relation of Eq. (1), *i.e.* $\mathcal{H} = \mathcal{W}/\{aa^\dagger = a^\dagger a + I\}$. In \mathcal{P} this relation takes the symbolic form

$$\Gamma = \lrcorner + \ddot{:}, \quad (13)$$

by which we mean that a given staircase path is equivalent to the sum of two staircase paths obtained by

- (i) replacing an *upper-left-hand* corner (Γ) by a *lower-right-hand* corner (\lrcorner), and
- (ii) removing the row and column which intersect in the given cell ($\ddot{:}$).

Note that this reduction is exactly equivalent to the decomposition of the associated




Figure 3. Path defined by the word $\mathbf{w} = a^{\dagger r} a^s a^{\dagger k} a^l$ and the associated Ferrers board.

Ferrers board induced by the rook problem. In fact, any path can be uniquely decomposed into a finite sum of paths without steps (*i.e.* paths pertaining to monomials $a^{\dagger r} a^s$). The latter constitute the basis in $\mathcal{P}/_{\{\Gamma=\lrcorner+\cdot\}}$ corresponding to the normally ordered basis in \mathcal{H} .

In this way, we obtain the combinatorial model of the Heisenberg–Weyl algebra as the algebra of paths with the relation Eq. (13), *i.e.*

$$\mathcal{H} = \mathcal{W}/_{\{aa^{\dagger} = a^{\dagger}a + I\}} \cong \mathcal{P}/_{\{\Gamma=\lrcorner+\cdot\}}. \quad (14)$$

With this equivalence the algebraic structure of \mathcal{H} and its calculus is recast in combinatorial terms, thus providing different perspectives and making accessible intuitive combinatorial arguments.

3.4. Example: Structure constants

To illustrate the above equivalence we show how to calculate the structure constants Eq. (3) of the Heisenberg–Weyl algebra by simple combinatorial enumeration. In the form suited for our purposes here we multiply two elements of the basis of Eq. (6), say $b_{(r,s)}$ and $b_{(k,l)}$, defining the structure constants $\gamma_{(r,s)(k,l)}^{(p,q)}$ in the form

$$b_{(r,s)} b_{(k,l)} = \sum_{p,q} \gamma_{(r,s)(k,l)}^{(p,q)} b_{(p,q)}. \quad (15)$$

Expansion in this basis of the product $b_{(r,s)} b_{(k,l)}$ essentially comes from the normal ordering of the word

$$\mathbf{w} = a^{\dagger r} a^s a^{\dagger k} a^l. \quad (16)$$

Following the scheme of Section 3.1 we draw the associated path and subsequently read off the Ferrers board $\mathcal{B}_{\mathbf{w}}$ which has a simple rectangular form, see Fig. 3. Enumeration

of non-capturing rook arrangements on the board is a direct result of the observations:

- (1) the maximum number of rooks on the board is $\min\{s, k\}$, and
- (2) there are as many possible arrangements of i rooks as unordered choices of i columns and rows in the board \mathcal{B}_w .

Hence the rook numbers are

$$r_{\mathcal{B}_w}(i) = i! \binom{s}{i} \binom{k}{i}, \quad (17)$$

for $i = 0 \dots \min\{s, k\}$ and zero otherwise. Consequently, Eq. (11) gives the normally ordered form of the word \mathfrak{w}

$$a^{\dagger r} a^s a^{\dagger k} a^l = \sum_{i=0}^{\min\{k, s\}} i! \binom{s}{i} \binom{k}{i} a^{\dagger r+k-i} a^{s+l-i}. \quad (18)$$

Finally, from Eq. (15) one may read off the structure constants of the algebra \mathcal{H} in the normally ordered basis. For fixed (r, s) and (k, l) the only non-vanishing γ s are

$$\gamma_{(r, s)(k, l)}^{(r+k-i, s+l-i)} = i! \binom{s}{i} \binom{k}{i} \quad \text{for } i = 0 \dots \min\{k, s\}. \quad (19)$$

Note that the right hand side of Eq. (19) is, in fact, independent of r and l since the outer elements are not included in the commutation.

4. Summary

We considered the Heisenberg–Weyl algebra \mathcal{H} starting from the (two generator) free algebra of words \mathcal{W} and imposing the relation of Eq. (1), *i.e.*

$$\mathcal{W} \xrightarrow{[a, a^{\dagger}] = I} \mathcal{H}$$

We showed that words can be uniquely encoded as staircase paths on a plane, thus providing a realization of \mathcal{W} as a combinatorial algebra of paths \mathcal{P} . This allowed us to construct a model of \mathcal{H} in terms of paths with the decomposition rule of Eq. (13) in \mathcal{P} which reflects the defining relation of Eq. (1) in \mathcal{W} . The following diagram illustrates the whole scheme

$$\begin{array}{ccc} \mathcal{W} & \xleftrightarrow{\sim} & \mathcal{P} \\ \downarrow & & \downarrow \\ \mathcal{W}/_{aa^{\dagger} = a^{\dagger}a + I} & \xleftrightarrow{\sim} & \mathcal{P}/_{\{\Gamma = \lrcorner + ::\}} \end{array}$$

We further looked at the normally ordered basis in \mathcal{H} and the corresponding basis in $\mathcal{P}/_{\{\Gamma = \lrcorner + ::\}}$. We pointed out that the decomposition rule of Eq. (13), reducing the number of steps in a path, is closely related to the familiar rook problem on the associated Ferrers board. This permits a graphical illustration of the calculus in this basis and, more generally, of the normal ordering problem.

In this note we have advocated a combinatorial approach to the Heisenberg–Weyl algebra by showing that it can be conceived as having a purely combinatorial origin.

This gives a new perspective on the whole algebraic framework which is easily amenable to the sophisticated methods of discrete mathematics. Finally, we should mention other models of the algebra based on various discrete structures such as set partitions, graphs or urn models [22, 23, 24] as well as results deriving from combinatorial methodology, see [17] and references therein.

Acknowledgments

We acknowledge our appreciation of the warm hospitality and support of the Mathematisches Forschungsinstitut Oberwolfach and the Laboratoire d’Informatique de l’Université Paris-Nord in Villetaneuse where most of this research was carried out. P.B and A.H. wish to acknowledge support from the Polish Ministry of Science and Higher Education under Grants No. N202 061434 and N202 107 32/2832.

References

- [1] C. J. Isham. *Lectures on Quantum Theory: Mathematical and Structural Foundations*. Imperial College Press, London, 1995.
- [2] A. Peres. *Quantum Theory: Concepts and Methods*. Kluwer Academic Publishers, New York, 2002.
- [3] P. A. M. Dirac. *The Principles of Quantum Mechanics*. Oxford University Press, New York, 4th edition, 1982.
- [4] J. Riordan. *An Introduction to Combinatorial Analysis*. Wiley, New York, 1984.
- [5] R. P. Stanley. *Enumerative Combinatorics*. Cambridge University Press, Cambridge, 1999.
- [6] L. Comtet. *Advanced Combinatorics*. Reidel, Dordrecht, 1974.
- [7] H. S. Wilf. *Generatingfunctionology*. Academic Press, New York, 1994.
- [8] P. Flajolet and R. Sedgewick. *Analytic Combinatorics*. Cambridge University Press, Cambridge, 2008.
- [9] F. Bergeron, G. Labelle, and P. Leroux. *Combinatorial Species and Tree-like Structures*. Cambridge University Press, Cambridge, 1998.
- [10] V. Bryant. *Aspects of Combinatorics: A Wide-ranging Introduction*. Cambridge University Press, Cambridge, 1968.
- [11] K. E. Cahill and R. J. Glauber. Ordered expansions in boson amplitude operators. *Phys.Rev.*, 177:1857–1881, 1969.
- [12] R. J. Glauber. The quantum theory of optical coherence. *Phys. Rev.*, 130:2529–2539, 1963.
- [13] W. P. Schleich. *Quantum Optics in Phase Space*. Wiley, Berlin, 2001.
- [14] J. R. Klauder and B-S. Skagerstam. *Coherent States. Application in Physics and Mathematical Physics*. World Scientific, Singapore, 1985.
- [15] J. D. Bjorken and S. D. Drell. *Relativistic Quantum Fields*. McGraw-Hill, New York, 1993.
- [16] R. D. Mattuck. *A Guide to Feynman Diagrams in the Many-Body Problem*. Dover Publications, New York, 2nd edition, 1992.
- [17] P. Blasiak, A. Horzela, K. A. Penson, A. I. Solomon, and G. H. E. Duchamp. Combinatorics and Boson normal ordering: A gentle introduction. *Am. J. Phys.*, 75:639–646, 2007. arXiv:0704.3116 [quant-ph].
- [18] A. M. Navon. Combinatorics and fermion algebra. *Nuovo Cimento*, 16B:324–330, 1973.
- [19] A. I. Solomon, G. H. E. Duchamp, P. Blasiak, A. Horzela, and K. A. Penson. Normal order: Combinatorial graphs. In *Proceedings of 3rd International Symposium on Quantum Theory and*

- Symmetries (Cincinnati 2003)*, pages 527–536, Singapore, 2004. World Scientific. arXiv:quant-ph/0402082.
- [20] A. Varvak. Rook numbers and the normal ordering problem. *Jour. Comb. Th., A* 112:292–307, 2005. arXiv:math.CO/0402376.
- [21] J. Katriel and G. Duchamp. Ordering relations for q -boson operators, continued fraction techniques and the q -BCH enigma. *J. Phys. A : Math. Gen.*, 28:7209–7225, 1995.
- [22] M. A. Méndez, P. Blasiak, and K. A. Penson. Combinatorial approach to generalized Bell and Stirling numbers and boson normal ordering problem. *J. Math. Phys.*, 46:083511, 2005. arXiv:quant-ph/0505180.
- [23] P. Blasiak and A. Horzela. Graphs for Quantum Theory. 2008. arXiv:0710.0266 [quant-ph].
- [24] P. Blasiak. Urn Models & Operator Ordering Procedures. *Oberwolfach Preprints*, OWP 2008 - 06, 2008. Available at: <http://www.mfo.de/publications/owp>.