


# Oberwolfach Preprints



OWP 2010 - 10

SOPHIE MORIER-GENOUD AND VALENTIN OVSIENKO

A Series of Algebras Generalizing the Octonions  
and Hurwitz-Radon Identity

Mathematisches Forschungsinstitut Oberwolfach gGmbH  
Oberwolfach Preprints (OWP) ISSN 1864-7596

## Oberwolfach Preprints (OWP)

Starting in 2007, the MFO publishes a preprint series which mainly contains research results related to a longer stay in Oberwolfach. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 30 free hard copies (DIN A4, black and white copy) by surface mail.

Of course, the full copy right is left to the authors. The MFO only needs the right to publish it on its website *www.mfo.de* as a documentation of the research work done at the MFO, which you are accepting by sending us your file.

In case of interest, please send a **pdf file** of your preprint by email to *rip@mfo.de* or *owlf@mfo.de*, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

## Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)  
Schwarzwaldstrasse 9-11  
77709 Oberwolfach-Walke  
Germany

Tel +49 7834 979 50  
Fax +49 7834 979 55  
Email [admin@mfo.de](mailto:admin@mfo.de)  
URL [www.mfo.de](http://www.mfo.de)

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.  
Copyright of the content is held by the authors.

# A SERIES OF ALGEBRAS GENERALIZING THE OCTONIONS AND HURWITZ-RADON IDENTITY

SOPHIE MORIER-GENOUD AND VALENTIN OVSIENKO

ABSTRACT. We study non-associative twisted group algebras over  $(\mathbb{Z}_2)^n$  with cubic twisting functions. We construct a series of algebras that extend the classical algebra of octonions in the same way as the Clifford algebras extend the algebra of quaternions. We study their properties, give several equivalent definitions and prove their uniqueness within some natural assumptions. We then prove a simplicity criterion.

We present two applications of the constructed algebras and the developed technique. The first application is a simple explicit formula for the following famous square identity:  $(a_1^2 + \dots + a_N^2)(b_1^2 + \dots + b_{\rho(N)}^2) = c_1^2 + \dots + c_N^2$ , where  $c_k$  are bilinear functions of the  $a_i$  and  $b_j$  and where  $\rho(N)$  is the Hurwitz-Radon function. The second application is the relation to Moufang loops and, in particular, to the code loops. To illustrate this relation, we provide an explicit coordinate formula for the factor set of the Parker loop.

**Key Words:** Graded commutative algebras, non-associative algebras, Clifford algebras, octonions, square identities, Hurwitz-Radon function.

**Mathematics Subject Classification (2010):** 16W50, 15A66, 11E25, 94B25.

## CONTENTS

1. Introduction	2
2. Twisted algebras over $(\mathbb{Z}_2)^n$ : definitions and tools	5
2.1. Twisted group algebras	6
2.2. Quasialgebra structure	6
2.3. The pentagonal and the hexagonal diagrams	7
2.4. The signature	7
2.5. Isomorphic twisted algebras	8
2.6. Generating functions and anti-involutions	9
3. The series $\mathbb{O}_n$ and $\mathbb{M}_n$ : characterization	10
3.1. Symmetric quasialgebras	10
3.2. The generating functions of the algebras $\mathbb{O}_n$ and $\mathbb{M}_n$	11
3.3. Characterization of the algebras of the $\mathbb{O}$ - and $\mathbb{M}$ -series	12
3.4. Generators and relations	13
4. The series $\mathbb{O}_n$ and $\mathbb{M}_n$ : properties	15
4.1. Criterion of simplicity	15
4.2. The first algebras of the series	16
4.3. The commutation graph	17
5. Generating Functions	19
5.1. Existence of a generating function	19

5.2. Generating functions are cubic	20
5.3. Uniqueness of the generating function	21
5.4. From the generating function to the twisting function	21
6. Proof of the simplicity criterion	22
6.1. The idea of the proof	22
6.2. Central elements	22
6.3. Proof of Theorem 2, part (i)	24
6.4. Proof of Theorem 2, part (ii)	24
7. Hurwitz-Radon square identities	25
7.1. The explicit solution	26
7.2. The Euclidean norm	26
7.3. Proof of Theorem 3	27
8. Relation to code loops	28
Appendix: linear algebra and differential calculus over $\mathbb{Z}_2$	29
References	31

**Acknowledgments.** This work was completed at the Mathematisches Forschungsinstitut Oberwolfach (MFO). The first author has benefited from the award of a *Leibniz Fellowship*, the second author is also grateful to MFO for hospitality. We are pleased to thank Christian Duval, Alexey Lebedev, Dmitry Leites, John McKay and Sergei Tabachnikov for their interest and helpful comments.

## 1. INTRODUCTION

The starting idea of this work is the following naive question: *is there a natural way to multiply  $n$ -tuples of 0 and 1?*

Of course, it is easy to find such algebraic structures. The abelian group  $(\mathbb{Z}_2)^n$  provides such a multiplication, but the corresponding group algebra  $\mathbb{K}[(\mathbb{Z}_2)^n]$ , where  $\mathbb{K}$  is an arbitrary field, is not a simple algebra, since it is isomorphic to a direct sum of  $n$  copies of  $\mathbb{K}[\mathbb{Z}_2]$ . A much more interesting algebraic structure on  $\mathbb{K}[(\mathbb{Z}_2)^n]$  is given by the twisted product

$$(1.1) \quad u_x \cdot u_y = (-1)^{f(x,y)} u_{x+y},$$

where  $x, y \in (\mathbb{Z}_2)^n$  and  $f$  is a two-argument function on  $(\mathbb{Z}_2)^n$  with values in  $\mathbb{Z}_2 \cong \{0, 1\}$ . We use the standard notations  $u_x$  for the element of  $\mathbb{K}[(\mathbb{Z}_2)^n]$  corresponding to  $x \in (\mathbb{Z}_2)^n$ . The only difference between the above product and that of the group algebra  $\mathbb{K}[(\mathbb{Z}_2)^n]$  is the sign. Yet, the structure of the algebra changes completely.

Remarkably enough, the classical Clifford algebras can be obtained as twisted group algebras. The first example is the algebra of quaternions,  $\mathbb{H}$ . This example was found by many authors but probably first in [20]. The algebra  $\mathbb{H}$  is a twisted  $(\mathbb{Z}_2)^2$ -algebra. More precisely, consider the 4-dimensional vector space over  $\mathbb{R}$  spanned by  $(0, 0)$ ,  $(0, 1)$ ,  $(1, 0)$  and  $(1, 1)$  with the multiplication:

$$u_{(x_1, x_2)} \cdot u_{(y_1, y_2)} = (-1)^{x_1 y_1 + x_1 y_2 + x_2 y_2} u_{(x_1 + y_1, x_2 + y_2)}.$$

It is easy to check that the obtained twisted  $(\mathbb{Z}_2)^2$ -algebra is, indeed, isomorphic to  $\mathbb{H}$ , see also [22] for a different,  $(\mathbb{Z}_2)^3$ -grading on the quaternions.

Along the same lines, a Clifford algebra with  $n$  generators, is a  $(\mathbb{Z}_2)^n$ -graded algebra, see [5]. The (complex) Clifford algebra  $Cl_n$  is isomorphic to the twisted group algebras over  $(\mathbb{Z}_2)^n$  with the product

$$(1.2) \quad u_{(x_1, \dots, x_n)} \cdot u_{(y_1, \dots, y_n)} = (-1)^{\sum_{1 \leq i \leq j \leq n} x_i y_j} u_{(x_1 + y_1, \dots, x_n + y_n)},$$

where  $(x_1, \dots, x_n)$  is an  $n$ -tuple of 0 and 1. The above twisting function is bilinear and therefore is a 2-cocycle on  $(\mathbb{Z}_2)^n$ . The real Clifford algebras  $Cl_{p,q}$  are also twisted group algebras over  $(\mathbb{Z}_2)^n$ . The twisting function  $f$  in the real case contains an extra term  $\sum_{1 \leq i \leq p} x_i y_i$  corresponding to the signature.




FIGURE 1.  $(\mathbb{Z}_2)^3$ -grading on the octonions.

The algebra of octonions  $\mathbb{O}$  can also be viewed as a twisted group algebra [4]. It is isomorphic to  $\mathbb{R} [(\mathbb{Z}_2)^3]$  equipped with the following product:

$$u_{(x_1, x_2, x_3)} \cdot u_{(y_1, y_2, y_3)} = (-1)^{(x_1 x_2 y_3 + x_1 y_2 x_3 + y_1 x_2 x_3 + \sum_{1 \leq i \leq j \leq 3} x_i y_j)} u_{(x_1 + y_1, x_2 + y_2, x_3 + y_3)}.$$

Note that the twisting function in this case is a polynomial of degree 3, and does not define a 2-cocycle. This is equivalent to the fact that the algebra  $\mathbb{O}$  is not associative. The multiplication table on  $\mathbb{O}$  is usually represented by the Fano plane. The corresponding  $(\mathbb{Z}_2)^3$ -grading is given in Figure 1. We also mention that different group gradings on  $\mathbb{O}$  were studied in [14], we also refer to [6] for a survey on the octonions and Clifford algebras.

In this paper, we introduce two series of complex algebras,  $\mathbb{O}_n$  and  $\mathbb{M}_n$ , and of real algebras,  $\mathbb{O}_{p,q}$  and  $\mathbb{M}_{p,q}$ . The series  $\mathbb{O}_n$  and  $\mathbb{O}_{p,q}$  generalize the algebra of octonions in a similar way as the Clifford algebras generalize the algebra of quaternions. The situation



The relations of higher degree are then calculated inductively using the following simple “linearity law”. Given three monomials  $u, v, w$ , then

$$u \cdot (v \cdot w) = (-1)^{\phi(\deg u, \deg v, \deg w)} (u \cdot v) \cdot w,$$

where  $\phi$  is the trilinear function uniquely defined by the above relations of degree 3, see Section 3.4 for the details. For instance, one has

$$u_i \cdot ((u_j \cdot u_k) \cdot u_\ell) = (u_i \cdot (u_j \cdot u_k)) \cdot u_\ell,$$

for  $i \neq j \neq k \neq \ell$ , etc.

The presentation of  $\mathbb{M}_n$  is exactly the same as above, except that the generators of  $\mathbb{M}_n$  commute. We will prove two classification results characterizing the algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  algebras in an axiomatic way.

The algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  are closely related to the theory of Moufang loops and, in particular, to code loops, see [17, 13, 23] and references therein. Indeed, the homogeneous elements  $\pm u_x$ , where  $x \in (\mathbb{Z}_2)^n$  form a Moufang loop of rank  $2^{n+1}$ . As an application, we show in Section 8 how the famous Parker loop fits into our framework.

The main results of the paper contain three theorems and their corollaries.

- (1) Theorem 1 is a general characterization of non-associative twisted group algebras over  $(\mathbb{Z}_2)^n$  with symmetric non-associativity factor, in terms of *generating functions*. This notion is our main tool and the way to show why the cubic twisting functions are so special. It also relates the subject to Moufang (code) loops.
- (2) Theorem 2 answers the question for which  $n$  (and  $p, q$ ) the the constructed algebras are simple. The result is quite similar to that for the Clifford algebras, except that the algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  degenerate for one value of  $n$  over 4 and not 1 over 2 as  $C\ell_n$ .
- (3) Theorem 3 provides explicit formulæ of the Hurwitz-Radon square identities. The algebras  $\mathbb{O}_n$  (as well as  $\mathbb{M}_n$ ) are not composition algebras. However, they have natural Euclidean norm  $\mathcal{N}$ . We obtain a necessary and sufficient condition for elements  $u$  and  $v$  to satisfy  $\mathcal{N}(u \cdot v) = \mathcal{N}(u)\mathcal{N}(v)$ . Whenever we find two subspaces  $V, W \subset \mathbb{O}_n$  consisting of elements satisfying this condition, we obtain a square identity generalizing the famous “octonionic” 8-square identity.

Our main tools include variations on the cohomology of  $(\mathbb{Z}_2)^n$  and the linear algebra over  $(\mathbb{Z}_2)^n$ . A brief account on this subject is presented in Appendix. In particular, we introduce and extensively use the notion of generating function associated with an algebra. This is a (polynomial of degree 3) function on  $(\mathbb{Z}_2)^n$  with values in  $\mathbb{Z}_2$ . This is a very elementary object, but it contains the full structure of the algebra (except the signature in the real case). Note that generating functions are useful even for the classical Clifford and octonion algebras and simplify the proofs of known results.

## 2. TWISTED ALGEBRAS OVER $(\mathbb{Z}_2)^n$ : DEFINITIONS AND TOOLS

In this section, we recall the notion of twisted group algebra with a twisting function that is not necessarily a 2-cocycle, we also recall the related notion of graded quasialgebra introduced in [4]. Sections 2.1–2.4 contain a well-known and elementary material. In Section 2.5, we show that, in the complex case, the structure of quasialgebra determines

the twisted group algebra itself, up to isomorphism. Finally, in Section 2.6, we define the notion of generating function which is the main tool we use in this paper.

**2.1. Twisted group algebras.** The most general definition is the following. Let  $(G, +)$  be an abelian group. A *twisted group algebra*  $(\mathbb{K}[G], F)$  is the algebra spanned by the elements  $u_x$  for  $x \in G$  and equipped with the product

$$u_x \cdot u_y = F(x, y) u_{x+y},$$

where  $F : G \times G \rightarrow \mathbb{K}^*$  is an *arbitrary* two-argument function such that

$$F(0, \cdot) = F(\cdot, 0) = 1.$$

The algebra  $(\mathbb{K}[G], F)$  is always unital and it is associative if and only if  $F$  is a 2-cocycle on  $G$ . Twisted group algebras is a classical subject (see, e.g., [9, 7] and references therein).

We will be interested in the particular case of twisted algebras over  $G = (\mathbb{Z}_2)^n$  and the twisting function  $F$  of the form

$$F(x, y) = (-1)^{f(x, y)},$$

with  $f$  taking values in  $\mathbb{Z}_2 \cong \{0, 1\}$ . The function  $f$  will not necessarily be a 2-cocycle.

**2.2. Quasialgebra structure.** An arbitrary twisted group algebra  $\mathcal{A} = (\mathbb{K}[(\mathbb{Z}_2)^n], f)$  gives rise to two functions

$$\beta : (\mathbb{Z}_2)^n \times (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2, \quad \phi : (\mathbb{Z}_2)^n \times (\mathbb{Z}_2)^n \times (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$$

such that

$$(2.1) \quad u_x \cdot u_y = (-1)^{\beta(x, y)} u_y \cdot u_x,$$

$$(2.2) \quad u_x \cdot (u_y \cdot u_z) = (-1)^{\phi(x, y, z)} (u_x \cdot u_y) \cdot u_z,$$

for any homogeneous elements  $u_x, u_y, u_z \in \mathcal{A}$ . The function  $\beta$  obviously satisfies the following properties:  $\beta(x, y) = \beta(y, x)$  and  $\beta(x, x) = 0$ . Following [4], we call the structure  $\beta, \phi$  a *graded quasialgebra*.

The functions  $\beta$  and  $\phi$  can be expressed in terms of the twisting function  $f$ :

$$(2.3) \quad \beta(x, y) = f(x, y) + f(y, x),$$

$$(2.4) \quad \phi(x, y, z) = f(y, z) + f(x + y, z) + f(x, y + z) + f(x, y).$$

Note that (2.4) reads  $\phi = \delta f$ , where  $\delta$  is the usual coboundary operator, cf. [1]. In particular,  $\phi$  is a (trivial) 3-cocycle. Conversely, given the functions  $\beta$  and  $\phi$ , to what extent the corresponding function  $f$  is uniquely defined? We will give the answer to this question in Section 2.5.

**Example 2.1.** (a) For the Clifford algebra  $\mathcal{C}\ell_n$  (and for  $\mathcal{C}\ell_{p, q}$  with  $p + q = n$ ), the function  $\beta$  is bilinear:

$$\beta_{\mathcal{C}\ell}(x, y) = \sum_{i \neq j} x_i y_j.$$

The function  $\phi \equiv 0$  since the twisting function (1.2) is a 2-cocycle, this is of course equivalent to the associativity property. Every simple graded quasialgebra with bilinear  $\beta$  and  $\phi \equiv 0$  is a Clifford algebra, see [22].

(b) For the algebra of octonions  $\mathbb{O}$ , the function  $\beta$  is as follows:  $\beta(x, y) = 0$  if either  $x = 0$ , or  $y = 0$ , or  $x = y$ ; otherwise,  $\beta(x, y) = 1$ . The function  $\phi$  is the determinant of  $3 \times 3$  matrices:

$$\phi(x, y, z) = \det |x, y, z|,$$

where  $x, y, z \in (\mathbb{Z}_2)^3$ . This function is symmetric and trilinear.

**Remark 2.2.** The notion of graded quasialgebra was defined in [5] in a more general situation where  $G$  is an arbitrary abelian group and the functions that measure the defect of commutativity and associativity take values in  $\mathbb{K}^*$  instead of  $\mathbb{Z}_2$ . The “restricted version” we consider is very special and this is the reason we can say much more about it. On the other hand, many classical algebras can be treated within our framework.

**2.3. The pentagonal and the hexagonal diagrams.** Consider any three homogeneous elements,  $u, v, w \in \mathcal{A}$ . The functions  $\beta$  and  $\phi$  relate the different products,  $u(vw), (uv)w, (vu)w$ , etc. The hexagonal diagrams in Figure 2 represent different loops in  $\mathcal{A}$  that lead




FIGURE 2. Two hexagonal and the pentagonal commutative diagrams

to the following identities

$$(2.5) \quad \begin{aligned} \phi(x, y, z) + \beta(x, y + z) + \phi(y, z, x) + \beta(z, x) + \phi(y, x, z) + \beta(x, y) &= 0, \\ \phi(x, y, z) + \beta(z, y) + \phi(x, z, y) + \beta(z, x) + \phi(z, x, y) + \beta(x + y, z) &= 0. \end{aligned}$$

Note that these identities can be checked directly from (2.3) and (2.4). In a similar way, the products of any four homogeneous elements  $t, u, v, w$ , see the pentagonal diagrams of Figure 2 is equivalent to the condition

$$(2.6) \quad \phi(y, z, t) + \phi(x + y, z, t) + \phi(x, y + z, t) + \phi(x, y, z + t) + \phi(x, y, z) = 0,$$

which is nothing but the 3-cocycle condition  $\delta\phi = 0$ . We already knew this identity from  $\phi = \delta f$ .

Let us stress on the fact that these two commutative diagrams are tautologically satisfied and give no restriction on  $f$ .

**2.4. The signature.** We say that the twisting functions  $f$  and  $f'$  differs by a *signature* if one has

$$(2.7) \quad f(x, y) = f'(x, y) + x_{i_1}y_{i_1} + \cdots + x_{i_p}y_{i_p},$$

where  $p \leq n$  is an integer. The above extra term changes only the squares of the generators  $u_i$ , all the other relations remain unchanged. Note that  $f - f'$  is a non-trivial 2-cocycle.

The signature represents the main difference between the twisted group algebras over  $\mathbb{C}$  and  $\mathbb{R}$ . In the complex case, the corresponding algebras  $\mathcal{A}$  and  $\mathcal{A}'$  are isomorphic. Indeed, consider the following set of generators:

$$(2.8) \quad u_i = u_{(0, \dots, 0, 1, 0, \dots, 0)},$$

where 1 stands at  $i$ -th position. The map defined in these generators by

$$u_i \mapsto \begin{cases} \sqrt{-1} u_i, & i = i_1, \dots, i_p, \\ u_i, & \text{otherwise,} \end{cases}$$

is an isomorphism between  $\mathcal{A}$  and  $\mathcal{A}'$ . In the real case, the algebras  $\mathcal{A}$  and  $\mathcal{A}'$  can be non-isomorphic but can also be isomorphic.

The quasilgebra structures corresponding to the twisted group algebras  $\mathcal{A}$  and  $\mathcal{A}'$  are identically the same. The relations between the generators are also the same for  $\mathcal{A}$  and for  $\mathcal{A}'$ , except for the sign in  $u_i^2 = \pm 1$ .

**2.5. Isomorphic twisted algebras.** It is natural to ask under what condition two functions  $f$  and  $f'$  define isomorphic algebras. Unfortunately, we do not know the complete answer to this question and give here two conditions which are sufficient but certainly not necessary.

(a) If  $f - f' = \delta b$  is a coboundary, i.e.,  $b : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  is a function such that

$$(2.9) \quad f(x, y) - f'(x, y) = b(x + y) + b(x) + b(y),$$

then the corresponding twisted algebras are isomorphic. Indeed, the isomorphism is given by the map  $u_x \mapsto (-1)^{b(x)} u_x$ , for all  $x \in (\mathbb{Z}_2)^n$ .

(b) Given a group automorphism  $T : (\mathbb{Z}_2)^n \rightarrow (\mathbb{Z}_2)^n$ , the functions  $f$  and

$$f'(x, y) = f(T(x), T(y))$$

define isomorphic twisted group algebras via  $u_x \mapsto u_{T^{-1}(x)}$ . Note that the automorphisms of  $(\mathbb{Z}_2)^n$  are just arbitrary linear transformations.

We are ready to answer the question formulated in the end of Section 2.2.

**Proposition 2.3.** *Given two twisted algebras  $\mathcal{A} = (\mathbb{K}[(\mathbb{Z}_2)^n], f)$  and  $\mathcal{A}' = (\mathbb{K}[(\mathbb{Z}_2)^n], f')$ , the corresponding quasilgebra structures coincide, i.e.,  $\beta' = \beta$  and  $\phi' = \phi$ , if and only if*

$$f(x, y) - f'(x, y) = \delta b(x, y) + \sum_{1 \leq i \leq n} \lambda_i x_i y_i,$$

where  $b : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  is an arbitrary function, and  $\lambda_i$  are coefficients in  $\mathbb{Z}_2$ . In particular, if  $\mathbb{K} = \mathbb{C}$ , then  $\mathcal{A} \cong \mathcal{A}'$ .

*Proof.* If the quasilgebras structures coincide, then  $\phi' = \phi$  implies  $\delta f' = \delta f$ , so that  $f - f'$  is a 2-cocycle. We use the well-known information about the second cohomology space  $H^2((\mathbb{Z}_2)^n; \mathbb{Z}_2)$  (see, e.g., [1]). Every non-trivial 2-cocycle is a linear combination of the following bilinear maps:  $(x, y) \mapsto x_i y_i$ , for some  $i$  and  $(x, y) \mapsto x_k y_\ell$ , for some  $k < \ell$ . One deduces that  $f - f'$  is of the form

$$f(x, y) - f'(x, y) = \delta b(x, y) + \sum_{1 \leq i \leq n} \lambda_i x_i y_i + \sum_{k < \ell} \mu_{k\ell} x_k y_\ell.$$

Since  $\beta' = \beta$ , one observes that  $f - f'$  is symmetric, so that the last summand vanishes, while the second summand is nothing but the signature.

Conversely, if  $f$  and  $f'$  are related by the above expression, then the quasialgebra structures obviously coincide.  $\square$

**2.6. Generating functions and anti-involutions.** As usual, we will be interested in the case where  $\mathcal{A} = (\mathbb{K}[(\mathbb{Z}_2)^n], f)$  is a twisted group algebra. However, the following notion makes sense for an arbitrary  $G$ -graded quasialgebra.

**Definition 2.4.** Given a  $G$ -graded quasialgebra, a function  $\alpha : G \rightarrow \mathbb{Z}_2$  will be called a *generating function* if the binary function  $\beta$  and the ternary function  $\phi$  defined by (2.1) and (2.2) are both determined by  $\alpha$  via

$$(2.10) \quad \beta(x, y) = \alpha(x + y) + \alpha(x) + \alpha(y),$$

$$(2.11) \quad \begin{aligned} \phi(x, y, z) &= \alpha(x + y + z) \\ &\quad + \alpha(x + y) + \alpha(x + z) + \alpha(y + z) \\ &\quad + \alpha(x) + \alpha(y) + \alpha(z). \end{aligned}$$

Note that the identity (2.10) implies that  $\alpha$  vanishes on the zero element  $0 = (0, \dots, 0)$  of  $(\mathbb{Z}_2)^n$ , because the corresponding element  $1 := u_0$  is the unit of  $\mathcal{A}$  and therefore commutes with any other element of  $\mathcal{A}$ .

To the best of our knowledge, the notion of generating function has not been considered. This notion will be crucial for us, but it also seems to be useful in many cases, for instance, to encode the structure of the Clifford algebras in a simple and natural way.

**Remark 2.5.** The identity (2.10) means that  $\beta$  is the differential of  $\alpha$  in the usual sense of group cohomology (see, e.g., [1]). In particular,  $\beta$  is a 2-coboundary. The second identity (2.11) suggests the operator of “second derivation”,  $\delta_2$ , defined by the right-hand-side, so that the above identities then read:

$$\beta = \delta\alpha, \quad \phi = \delta_2\alpha.$$

The algebra  $\mathcal{A}$  is commutative if and only if  $\delta\alpha = 0$ ; it is associative if and only if  $\delta_2\alpha = 0$ . The cohomological meaning of the operator  $\delta_2$  will be discussed in Appendix.

The most important feature of the notion of generating function is the following. In the complex case, *the generating function contains the full information about the algebra*. In the real case, the generating function determines the algebra up to the signature. This immediately follows from Proposition 2.3.

Recall that an *anti-involution* on an algebra  $\mathcal{A}$  is a linear map  $a \mapsto \bar{a}$  from  $\mathcal{A}$  to  $\mathcal{A}$  such that  $\overline{ab} = \bar{b}\bar{a}$  and  $\bar{\bar{a}} = a$ . Every generating function defines an anti-involution of the following particular form:

$$(2.12) \quad \overline{u_x} = (-1)^{\alpha(x)} u_x,$$

**Proposition 2.6.** *If  $\alpha$  is a generating function, then the linear map defined by formula (2.12) is an anti-involution.*

*Proof.* Using (2.1) and (2.10), one has

$$\overline{u_x u_y} = (-1)^{\alpha(x+y)} u_x u_y = (-1)^{\alpha(x+y)+\beta(x,y)} u_y u_x = (-1)^{\alpha(x)+\alpha(y)} u_y u_x = \overline{u_y} \overline{u_x}.$$

Hence the result.  $\square$

**Example 2.7.** (a) In the simplest examples of the algebras  $\mathbb{H}$  and  $\mathbb{O}$ , the anti-involution consists in the sign inverting of purely imaginary quaternions and octonions, respectively. One can easily check that this involution is actually a generating function. It is amazing that such simple functions contain the full information about the structure of  $\mathbb{H}$  and  $\mathbb{O}$ .

(b) The generating function of  $C\ell_n$  is as follows:

$$(2.13) \quad \alpha_{C\ell}(x) = \sum_{1 \leq i \leq j \leq n} x_i x_j.$$

Indeed, one checks that the binary function  $\beta$  defined by (2.10) is exactly the skew-symmetrization of the function  $f = \sum_{1 \leq i \leq j \leq n} x_i y_j$ . The function  $\phi$  defined by (2.11) is identically zero, since  $\alpha$  is a quadratic polynomial.

### 3. THE SERIES $\mathbb{O}_n$ AND $\mathbb{M}_n$ : CHARACTERIZATION

In this section, we formulate our first main result. Theorem 1, concerns the general properties of twisted  $(\mathbb{Z}_2)^n$ -algebras with  $\phi = \delta f$  symmetric. This result distinguishes a class of algebras of which our algebras of  $\mathbb{O}$ - and  $\mathbb{M}$ -series are the principal representatives. We will also present several different ways to define the algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$ , as well as of  $\mathbb{O}_{p,q}$  and  $\mathbb{M}_{p,q}$ .

**3.1. Symmetric quasialgebras.** Any arbitrary twisted group algebra leads to a quasialgebra structure. One needs to assume some additional conditions on the “twisting” function  $f$  in order to obtain an interesting class of algebras.

We will be interested in the case where the function  $\phi = \delta f$ , see formula (2.4), is symmetric:

$$(3.1) \quad \phi(x, y, z) = \phi(y, x, z) = \phi(x, z, y).$$

This condition seems to be very natural: it means that, if three elements,  $u_x, u_y$  and  $u_z$  form a antiassociative triplet, i.e., one has  $u_x \cdot (u_y \cdot u_z) = -(u_x \cdot u_y) \cdot u_z$ , then this property is independent of the ordering of the elements in the triplet.

An immediate consequence of the identity (2.5) is that, if  $\phi$  is symmetric, then it is completely determined by  $\beta$ :

$$(3.2) \quad \begin{aligned} \phi(x, y, z) &= \beta(x + y, z) + \beta(x, z) + \beta(y, z) \\ &= \beta(x, y + z) + \beta(x, y) + \beta(x, z), \end{aligned}$$

as the “defect of linearity” in each argument.

The following statement is our main result about the general structure of a twisted group algebra  $\mathcal{A} = (\mathbb{K}[(\mathbb{Z}_2)^n], f)$ . We formulate this result in a slightly more general context of  $(\mathbb{Z}_2)^n$ -graded quasialgebra.

**Theorem 1.** *Given a  $(\mathbb{Z}_2)^n$ -graded quasialgebra  $\mathcal{A}$ , the following conditions are equivalent.*

- (i) *The function  $\phi$  is symmetric.*
- (ii) *The algebra  $\mathcal{A}$  has a generating function.*

This theorem will be proved in Section 5.1.

It is now natural to ask under what condition a function  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  is a generating function for some twisted group algebra. The following statement provides a necessary and sufficient condition.

**Proposition 3.1.** *Given a function  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$ , there exists a twisted group algebra  $\mathcal{A}$  such that  $\alpha$  is a generating function of  $\mathcal{A}$ , if and only if  $\alpha$  is a polynomial of degree  $\leq 3$ .*

This proposition will be proved in Section 5.2. Furthermore, we will show in Section 5.3 that the generating function can be chosen in a canonical way.

Theorem 1 has a number of consequences. In particular, it implies two more important properties of  $\phi$ . The function  $\phi$  is called *trilinear* if it satisfies

$$(3.3) \quad \phi(x + y, z, t) = \phi(x, z, t) + \phi(y, z, t),$$

and similarly in each argument. The function  $\phi$  is *alternate* if it satisfies

$$(3.4) \quad \phi(x, x, y) = \phi(x, y, x) = \phi(y, x, x) = 0,$$

for all  $x, y \in (\mathbb{Z}_2)^n$ .

**Corollary 3.2.** *If the function  $\phi$  is symmetric, then  $\phi$  is trilinear and alternate.*

Corollary 3.2 will be proved in Section 5.2.

**Remark 3.3.** Let us stress that an algebra satisfying all the above properties is *graded-alternative*, i.e.,  $u_x \cdot (u_x \cdot u_y) = u_x^2 \cdot u_y$  and  $(u_y \cdot u_x) \cdot u_x = u_y \cdot u_x^2$ , for all homogeneous elements  $u_x$  and  $u_y$ . This does not imply that the algebra is alternative. Let us mention that alternative graded quasialgebras were classified in [3].

Our next goal is to study two series of algebras with symmetric function  $\phi = \delta f$ . Let us notice that the Cayley-Dickson algebras are not of this type, cf. [4].

**3.2. The generating functions of the algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$ .** We already defined the complex algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$ , with  $n \geq 3$  and the real algebra  $\mathbb{O}_{p,q}$  and  $\mathbb{M}_{p,q}$  see Introduction, formulæ (1.3) and (1.4).

It is very easy to calculate the associated function  $\phi = \delta f$  which is exactly the same for  $f = f_{\mathbb{O}}$  or  $f_{\mathbb{M}}$ . One obtains

$$\phi(x, y, z) = \sum_{i \neq j \neq k} x_i y_j z_k.$$

This function is symmetric in  $x, y, z$  and Theorem 1 implies that the algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  have generating functions. The explicit formulæ are as follows:

$$(3.5) \quad \alpha_{\mathbb{O}}(x) = \sum_{1 \leq i < j < k \leq n} x_i x_j x_k + \sum_{1 \leq i < j \leq n} x_i x_j + \sum_{1 \leq i \leq n} x_i,$$

and

$$(3.6) \quad \alpha_{\mathbb{M}}(x) = \sum_{1 \leq i < j < k \leq n} x_i x_j x_k + \sum_{1 \leq i \leq n} x_i.$$

Note that the generating functions  $\alpha_{\mathbb{O}}$  and  $\alpha_{\mathbb{M}}$  are  $\mathfrak{S}_n$ -invariant with respect to the natural action of the group of permutations  $\mathfrak{S}_n$  on  $(\mathbb{Z}_2)^n$ .

Thanks to the  $\mathfrak{S}_n$ -invariance, we can give a very simple description of the above functions. Denote by  $|x|$  the *weight* of  $x \in (\mathbb{Z}_2)^n$  (i.e., the number of 1 entries in  $x$  written as an  $n$ -tuple of 0 and 1). The above generating functions, together with that of the Clifford algebras depend only on  $|x|$  and are 4-periodic:

$ x $	1	2	3	4	5	6	7	8	...
$\alpha_{Cl}$	1	1	0	0	1	1	0	0	...
$\alpha_{\mathbb{O}}$	1	1	1	0	1	1	1	0	...
$\alpha_{\mathbb{M}}$	1	0	0	0	1	0	0	0	...

This table is the most simple way to use the generating function in any calculation. One can deduce the explicit formulæ (2.13), (3.5) and (3.6) directly from the table (3.7).

**3.3. Characterization of the algebras of the  $\mathbb{O}$ - and  $\mathbb{M}$ -series.** Let us formulate two uniqueness results that allow us to give axiomatic definitions of the introduced algebras.

Recall that the group of permutations  $\mathfrak{S}_n$  acts on  $(\mathbb{Z}_2)^n$  in a natural way. We will characterize the algebras of the  $\mathbb{O}$ - and  $\mathbb{M}$ -series in terms of  $\mathfrak{S}_n$ -invariance. We observe that, in spite of the fact that the functions  $f_{\mathbb{O}}$  and  $f_{\mathbb{M}}$  are not  $\mathfrak{S}_n$ -invariant, the corresponding algebras are. However, we believe that  $\mathfrak{S}_n$ -invariance is a technical assumption and can be relaxed, see Appendix for a discussion.

The first uniqueness result is formulated directly in terms of the twisting function  $f$ . We study the unital twisted algebras  $\mathcal{A} = (\mathbb{K}[(\mathbb{Z}_2)^n], f)$  satisfying the following conditions.

- (1) The function  $f$  is a polynomial of degree 3.
- (2) The algebra  $\mathcal{A}$  is graded-alternative, see (3.4).
- (3) The set of relations between the generators (2.8) of  $\mathcal{A}$  is invariant with respect to the action of the group of permutations  $\mathfrak{S}_n$ .

**Proposition 3.4.** *The algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  are the only twisted  $(\mathbb{Z}_2)^n$ -algebras satisfying the above three conditions.*

*Proof.* Since the algebra  $\mathcal{A}$  is unital, we have  $f(0, \cdot) = f(\cdot, 0) = 0$ . This implies that  $f$  contains no constant term and no terms depending only on  $x$  (or only on  $y$ ) variables. The most general twisting function  $f$  of degree 3 is of the form

$$\begin{aligned}
 f(x, y) = & \sum_{i < j < k} \left( \lambda_{ijk}^1 x_i x_j y_k + \lambda_{ijk}^2 x_i y_j x_k + \lambda_{ijk}^3 y_i x_j x_k \right. \\
 & \left. + \mu_{ijk}^1 y_i y_j x_k + \mu_{ijk}^2 y_i x_j y_k + \mu_{ijk}^3 x_i y_j y_k \right) \\
 & + \sum_{i, j} \nu_{ij} x_i y_j,
 \end{aligned}$$

where  $\lambda_{ijk}^e, \mu_{ijk}^e$  and  $\nu_{ij}^e$  are arbitrary coefficients 0 or 1. Indeed, the expression of  $f$  cannot contain the monomials  $x_i x_j y_j$  and  $x_i y_i y_j$  because of the condition (2).

By Proposition 2.3, adding a coboundary to  $f$  does not change the algebra. We may assume that for any  $i < j < k$ , the coefficient  $\mu_{ijk}^1 = 0$  (otherwise, we add the coboundary of  $b(x) = x_i x_j x_k$ ).

We now compute  $\phi = \delta f$  and obtain:

$$\begin{aligned} \phi(x, y, z) = \sum_{i < j < k} & \left( (\lambda_{ijk}^1 + \mu_{ijk}^3) x_i y_j z_k + (\lambda_{ijk}^2 + \mu_{ijk}^3) x_i z_j y_k + (\lambda_{ijk}^1 + \mu_{ijk}^2) y_i x_j z_k \right. \\ & \left. + \lambda_{ijk}^2 y_i z_j x_k + (\lambda_{ijk}^3 + \mu_{ijk}^2) z_i x_j y_k + \lambda_{ijk}^3 x_i y_j z_k \right). \end{aligned}$$

We can assume that

$$u_i \cdot (u_j \cdot u_k) = -(u_i \cdot u_j) \cdot u_k, \quad i \neq j \neq k.$$

Indeed, if  $u_i \cdot (u_j \cdot u_k) = (u_i \cdot u_j) \cdot u_k$  for some values of  $i, j, k$  such that  $i \neq j \neq k$ , then (3) implies the same associativity relation for all  $i, j, k$ . Since  $\phi$  is trilinear, this means that  $\mathcal{A}$  is associative, so that  $\phi = 0$ . This can only happen if  $\lambda_{ijk}^e = \mu_{ijk}^e = 0$  for all  $i, j, k$ , so that  $\deg f = 2$ .

In other words, we obtain a system of equations  $\phi(x_i, y_j, z_k) = 1$  for all  $i, j, k$ . This system has a unique solution  $\lambda_{ijk}^1 = \lambda_{ijk}^2 = \lambda_{ijk}^3 = 1$  and  $\mu_{ijk}^2 = \mu_{ijk}^3 = 0$ .

Finally, if all of the generators commute, we obtain  $\nu_{ij} = \nu_{ji}$ , so that  $\nu_{ij} = 0$  up to a coboundary, so that  $f = f_{\mathbb{M}}$ . If all of the generators anticommute, again up to a coboundary, we obtain  $\nu_{ij} = 1$ , if and only if  $i < j$ , so that  $f = f_{\mathbb{O}}$ .  $\square$

The second uniqueness result is formulated in terms of the generating function.

**Proposition 3.5.** *The complex algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  and the real algebras  $\mathbb{O}_{p,q}$  and  $\mathbb{M}_{p,q}$  with  $p + q = n$ , are the only non-associative twisted  $(\mathbb{Z}_2)^n$ -algebras with an  $\mathfrak{S}_n$ -invariant generating function.*

*Proof.* By Proposition 3.1, we know that the generating function is a polynomial of degree 3. Every  $\mathfrak{S}_n$ -invariant polynomial  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  of degree 3 is a linear combination

$$\alpha = \lambda_3 \alpha_3 + \lambda_2 \alpha_2 + \lambda_1 \alpha_1 + \lambda_0 \alpha_0$$

of the following four functions:

$$\alpha_3(x) = \sum_{1 \leq i < j < k \leq n} x_i x_j x_k, \quad \alpha_2(x) = \sum_{1 \leq i < j \leq n} x_i x_j, \quad \alpha_1(x) = \sum_{1 \leq i \leq n} x_i, \quad \alpha_0(x) = 1.$$

Since  $\alpha(0) = 0$ , cf. Section 2.6, one obtains  $\lambda_0 = 0$ . The function  $\alpha_1$  does not contribute to the quasialgebra structure  $\beta = \delta \alpha$  and  $\phi = \delta_2 \alpha$ , so that  $\lambda_1$  can be chosen arbitrary. Finally,  $\lambda_3 \neq 0$  since otherwise  $\phi = 0$  and the corresponding algebra is associative. We obtain the functions  $\alpha_{\mathbb{O}} = \alpha_3 + \alpha_2 + \alpha_1$  and  $\alpha_{\mathbb{M}} = \alpha_3 + \alpha_1$  as the only possible  $\mathfrak{S}_n$ -invariant generating function that define non-associative algebras.  $\square$

Note that relaxing the non-associativity condition  $\phi \neq 0$ , will also recover the Clifford algebras  $Cl_n$  and  $Cl_{p,q}$  and the group algebra itself.

**3.4. Generators and relations.** Let us now give another definition of the complex algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  and of the real algebras  $\mathbb{O}_{p,q}$  and  $\mathbb{M}_{p,q}$ . We use a purely algebraic approach and present our algebras in terms of generators and relations.

Consider the generators (2.8). The following relations can be easily calculated directly using the functions (1.3) and (1.4).

One then has

$$(3.8) \quad u_i^2 = \begin{cases} 1, & i \leq p \\ -1, & \text{otherwise,} \end{cases}$$

where  $1 = u_{(0, \dots, 0)}$  is the unit. The rest of the relations is independent of the signature.

The main difference between the series  $\mathbb{O}$  and  $\mathbb{M}$  is that the generators *anticommute* in the  $\mathbb{O}$ -case and *commute* in the  $\mathbb{M}$ -case:

$$(3.9) \quad u_i \cdot u_j = -u_j \cdot u_i \quad \text{in } \mathbb{O}_n, \mathbb{O}_{p,q} \quad u_i \cdot u_j = u_j \cdot u_i \quad \text{in } \mathbb{M}_n, \mathbb{M}_{p,q}.$$

The third-order relations are determined by the function  $\phi$  and therefore these relations are the same for both series:

$$(3.10) \quad u_i \cdot (u_i \cdot u_j) = u_i^2 \cdot u_j,$$

$$(3.11) \quad u_i \cdot (u_j \cdot u_k) = -(u_i \cdot u_j) \cdot u_k,$$

where  $i \neq j \neq k$  in the second relation. Note that the antiassociativity relation in (3.11) is the reason why the algebras from the  $\mathbb{M}$  series generated by commuting elements, can, nevertheless, be simple.

The axiomatic definition characterizes a Clifford algebra as an algebra with  $n$  anticommuting generators satisfying the relations (3.8) and the identity of associativity. We will now give a very similar definition of the algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  (as well as  $\mathbb{O}_{p,q}$  and  $\mathbb{M}_{p,q}$ ). The associativity is replaced by the identity of quasialgebra.

Define a family of algebras  $\mathcal{A}$  with  $n$  generators  $u_1, \dots, u_n$ . Consider the monoid  $X_n$  of non-associative monomials in  $u_i$  and define a function  $\phi : X_n \times X_n \times X_n \rightarrow \mathbb{Z}_2$  satisfying the following two properties:

$$(1) \quad \phi(u_i, u_j, u_k) = \begin{cases} 1, & \text{if } i \neq j \neq k, \\ 0, & \text{otherwise.} \end{cases}$$

$$(2) \quad \phi(u \cdot u', v, w) = \phi(u, v, w) + \phi(u', v, w), \text{ and similar in each variable.}$$

Such function exists and is unique. Moreover,  $\phi$  is symmetric.

Define an algebra  $\mathcal{A}^{\mathbb{C}}$  or  $\mathcal{A}^{\mathbb{R}}$  (complex or real), generated by  $u_1, \dots, u_n$  that satisfies the relations (3.8) together with one of the following two relations. All the generators either anticommute:  $u_i \cdot u_j = -u_j \cdot u_i$ , where  $i \neq j$ , or commute:  $u_i \cdot u_j = u_j \cdot u_i$ , where  $i \neq j$ .

We will also assume the identity

$$u \cdot (v \cdot w) = (-1)^{\phi(u,v,w)} (u \cdot v) \cdot w,$$

for all monomials  $u, v, w$ .

**Proposition 3.6.** *If the generators anticommute, then  $\mathcal{A}^{\mathbb{C}} \cong \mathbb{O}_n$  and  $\mathcal{A}^{\mathbb{R}} \cong \mathbb{O}_{p,q}$ . If the generators commute, then  $\mathcal{A}^{\mathbb{C}} \cong \mathbb{M}_n$  and  $\mathcal{A}^{\mathbb{R}} \cong \mathbb{M}_{p,q}$ .*

*Proof.* By definition of  $\mathcal{A} = \mathcal{A}^{\mathbb{C}}$  (resp.  $\mathcal{A}^{\mathbb{R}}$ ), the elements

$$u_{i_1 \dots i_k} = u_{i_1} \cdot (u_{i_2} \cdot (\dots (u_{i_{k-1}} \cdot u_{i_k}) \dots)),$$

where  $i_1 < i_2 < \dots < i_k$ , form a basis of  $\mathcal{A}$ . Therefore,  $\dim \mathcal{A} = 2^n$ . The linear map sending the generators of  $\mathcal{A}$  to the generators (2.8) of  $\mathbb{O}_n$  or  $\mathbb{M}_n$  ( $\mathbb{O}_{p,q}$  or  $\mathbb{M}_{p,q}$ , respectively) is a homomorphism, since the function  $\phi$  corresponding to these algebras is symmetric and trilinear. It sends the above basis of  $\mathcal{A}$  to that of  $\mathbb{O}_n$  or  $\mathbb{M}_n$  ( $\mathbb{O}_{p,q}$  or  $\mathbb{M}_{p,q}$ , respectively).  $\square$

4. THE SERIES  $\mathbb{O}_n$  AND  $\mathbb{M}_n$ : PROPERTIES

In this section, we study properties of the algebras of the series  $\mathbb{O}$  and  $\mathbb{M}$ . The main result is Theorem 2 providing a criterion of simplicity. We describe the first algebras of the series and give the list of isomorphisms in lower dimensions. We also define a non-oriented graph encoding the structure of the algebra. Finally, we formulate open problems.

**4.1. Criterion of simplicity.** The most important property of the defined algebras that we study is the simplicity. Let us stress that we understand simplicity in the usual sense: an algebra is called *simple* if it contains no proper ideal. Note that in the case of commutative associative algebras, simplicity and division are equivalent notions, in our situation, the notion of simplicity is much weaker.

**Remark 4.1.** This notion should not be confounded with the notion of graded-simple algebra. The latter notion is much weaker and means that the algebra contains no graded ideal; however, this notion is rather a property of the grading and not of the algebra itself.

The following statement is the second main result of this paper. We will treat the complex and the real cases independently.

**Theorem 2.** (i) *The algebra  $\mathbb{O}_n$  (resp.  $\mathbb{M}_n$ ) is simple if and only if  $n \neq 4m$  (resp.  $n \neq 4m + 2$ ). One also has*

$$\mathbb{O}_{4m} \cong \mathbb{O}_{4m-1} \oplus \mathbb{O}_{4m-1}, \quad \mathbb{M}_{4m+2} \cong \mathbb{M}_{4m+1} \oplus \mathbb{M}_{4m+1}.$$

(ii) *The algebra  $\mathbb{O}_{p,q}$  is simple if and only if one of the following conditions is satisfied*

- (1)  $p + q \neq 4m$ ,
- (2)  $p + q = 4m$  and  $p, q$  are odd;

(iii) *The algebra  $\mathbb{M}_{p,q}$  is simple if and only if one of the following conditions is satisfied*

- (1)  $p + q \neq 4m + 2$ ,
- (2)  $p + q = 4m + 2$  and  $p, q$  are odd.

This theorem will be proved in Section 6.

The arguments developed in the proof of Theorem 2 allow us to link the complex and the real algebras in the particular cases below. Let us use the notation  $\mathbb{O}_n^{\mathbb{R}}$  and  $\mathbb{M}_n^{\mathbb{R}}$  when we consider the algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  as  $2^{n+1}$ -dimensional real algebras. We have the following statement.

**Corollary 4.2.** (i) *If  $p + q = 4m$  and  $p, q$  are odd, then  $\mathbb{O}_{p,q} \cong \mathbb{O}_{p+q-1}^{\mathbb{R}}$ .*

(ii) *If  $p + q = 4m + 2$  and  $p, q$  are odd, then  $\mathbb{M}_{p,q} \cong \mathbb{M}_{p+q-1}^{\mathbb{R}}$ .*

This statement is proved in Section 6.4.

**Remark 4.3.** To explain the meaning of the above statement, we notice that, in the case where the complex algebras split into a direct sum, the real algebras can still be simple. In this case, all the simple real algebras are isomorphic to the complex algebra with  $n - 1$  generators. In particular, all the algebras  $\mathbb{O}_{p,q}$  and  $\mathbb{O}_{p',q'}$  with  $p + q = p' + q' = 4m$  and  $p$  and  $p'$  odd are isomorphic to each other (and similarly for the  $\mathbb{M}$ -series). A very similar property holds for the Clifford algebras.

Theorem 2 immediately implies the following.

**Corollary 4.4.** *The algebras  $\mathbb{O}_n$  and  $\mathbb{M}_n$  with even  $n$  are not isomorphic.*

This implies, in particular, that the real algebras  $\mathbb{O}_{p,q}$  and  $\mathbb{M}_{p',q'}$  with  $p+q = p'+q' = 2m$  are not isomorphic.

**4.2. The first algebras of the series.** Let us consider the first examples of the introduced algebras. It is natural to ask if some of the introduced algebras can be isomorphic to the other ones.

**Proposition 4.5.** *(i) For  $n = 3$ , one has:*

$$\mathbb{O}_{3,0} \cong \mathbb{O}_{2,1} \cong \mathbb{O}_{1,2} \not\cong \mathbb{O}_{0,3}.$$

*The first three algebras are isomorphic to the algebra of split-octonions, while  $\mathbb{O}_{0,3} \cong \mathbb{O}$ .*

*(ii) For  $n = 4$ , one has:*

$$\mathbb{O}_{4,0} \cong \mathbb{O}_{2,2} \cong \mathbb{O}_{3,0} \oplus \mathbb{O}_{3,0}, \quad \mathbb{O}_{0,4} \cong \mathbb{O}_{0,3} \oplus \mathbb{O}_{0,3}.$$

*In particular,  $\mathbb{O}_{4,0}$  and  $\mathbb{O}_{2,2}$  are not isomorphic to  $\mathbb{O}_{0,4}$ .*

*Proof.* The above isomorphisms are combination of the general isomorphisms of type (a) and (b), see Section 2.5. The involved automorphisms of  $(\mathbb{Z}_2)^3$  and  $(\mathbb{Z}_2)^4$  are

$$\begin{array}{ll} x'_1 = x_1, & x'_1 = x_1, \\ x'_2 = x_1 + x_2, & x'_2 = x_1 + x_2, \\ x'_3 = x_1 + x_2 + x_3, & x'_3 = x_1 + x_2 + x_3, \\ & x'_4 = x_1 + x_2 + x_3 + x_4. \end{array}$$

Then, the twisting functions of the above isomorphic algebras coincide modulo coboundary.  $\square$

Let us notice that the very first algebras of the  $\mathbb{O}$ -series are all obtained as a combination of the algebras of octonions and split-octonions. In this sense, we do not obtain new algebras among them.

The next algebras,  $\mathbb{O}_5$  and  $\mathbb{M}_5$ , as well as all of the real algebras  $\mathbb{O}_{p,q}$  and  $\mathbb{M}_{p,q}$  with  $p+q = 5$ , are not combinations of the classical algebras. Since these algebras are simple, they are not direct sums of lower-dimensional algebras. The next statement shows that these algebras are not tensor products of classical algebras. Note that the only ‘‘candidate’’ for an isomorphism of this kind is the tensor product of the octonion algebra and the algebra of complex  $(2 \times 2)$ -matrices.

**Proposition 4.6.** *Neither of the algebras  $\mathbb{O}_5$  and  $\mathbb{M}_5$  is isomorphic to the tensor product of the octonion algebra  $\mathbb{O}$  and the algebra  $\mathbb{C}[2]$  of complex  $(2 \times 2)$ -matrices:*

$$\mathbb{O}_5 \not\cong \mathbb{O} \otimes \mathbb{C}[2], \quad \mathbb{M}_5 \not\cong \mathbb{O} \otimes \mathbb{C}[2].$$

*Proof.* Let us consider the element  $u = u_{(1,1,1,1,0)}$  in  $\mathbb{O}_5$  and the element  $u = u_{(1,1,0,0,0)}$  in  $\mathbb{M}_5$ . Each of these elements has a very big centralizer  $Z_u$  of  $\dim Z_u = 24$ . Indeed, the above element of  $\mathbb{O}_5$  commutes with itself and with any homogeneous element  $u_x$  of the weight  $|x| = 0, 1, 3, 5$  as well as 6 elements such that  $|x| = 2$ . The centralizer  $Z_u$  is the vector space spanned by these 24 homogeneous elements, and similarly in the  $\mathbb{M}_5$  case. We will show that the algebra  $\mathbb{O} \otimes \mathbb{C}[2]$  does not contain such an element.

Assume, *ad absurdum*, that an element  $u \in \mathbb{O} \otimes \mathbb{C}[2]$  has a centralizer of dimension  $\geq 24$ . It follows that  $Z_u$  contains at least two independent elements of the form

$$z_1 = e_1 \otimes 1 + 1 \otimes m_1, \quad z_2 = e_2 \otimes 1 + 1 \otimes m_2,$$

where  $e_1$  and  $e_2$  are pure imaginary octonions and  $m_1$  and  $m_2$  are traceless matrices. Indeed, the space spanned by the elements of this form is 10-dimensional. Without loss of generality, we can assume that one of the following holds

- (1) the generic case:  $e_1, e_2$  and  $m_1, m_2$  are linearly independent and pairwise anticommute,
- (2)  $e_2 = 0$  and  $m_1, m_2$  are linearly independent and anticommute,
- (3)  $m_2 = 0$  and  $e_1, e_2$  are linearly independent and anticommute.

In the case (1), an easy computation shows that the element  $u$  is proportional to

$$u = 1 \otimes 1 + e_1 \otimes m_1 + e_2 \otimes m_2 + e_1 e_2 \otimes m_1 m_2.$$

This element belongs to a subalgebra  $\mathbb{C}[4] = \mathbb{C}[2] \otimes \mathbb{C}[2] \subset \mathbb{O} \otimes \mathbb{C}[2]$ . The centralizer of  $u$  is of codimension  $\geq 10$ . Indeed, the codimension of  $Z_u$  inside  $\mathbb{C}[4]$  is at least 6. Furthermore, the 4-dimensional space of the elements  $e_3 \otimes 1$ , where  $e_3$  anticommutes with  $e_1, e_2$  is transversal to  $Z_u$ . Hence,  $\dim Z_u \leq 22$ . In the case (2),  $u$  is proportional to  $e_1 \otimes 1$  and one easily checks that  $Z_u = e_1 \otimes \mathbb{C}[2] \oplus 1 \otimes \mathbb{C}[2]$  is of dimension 8. In the case (3),  $u$  is proportional to  $1 \otimes m_1$  so that  $Z_u = \mathbb{O} \otimes 1 \oplus \mathbb{O} \otimes m_1$  is of dimension 16. In each case, we got a contradiction.  $\square$

In the  $\mathbb{M}$ -case, we have the following isomorphism.

**Proposition 4.7.** *One has*

$$\mathbb{M}_{1,2} \cong \mathbb{M}_{0,3},$$

*Proof.* This isomorphism can be obtained by the following automorphism of  $(\mathbb{Z}_2)^3$ .

$$x'_1 = x_1 + x_2 + x_3, \quad x'_2 = x_2, \quad x'_3 = x_3$$

This algebra is not isomorphic to  $\mathbb{O}_{0,3}$  or  $\mathbb{O}_{3,0}$ .  $\square$

**4.3. The commutation graph.** We associate a non-oriented graph, that we call the commutation graph, to every twisted group algebra in the following way. The vertices of the graph are the elements of  $(\mathbb{Z}_2)^n$ . The (non-oriented) edges  $x - y$  join the elements  $x$  and  $y$  such that  $u_x$  and  $u_y$  anticommute.

**Proposition 4.8.** *Given a complex algebra  $(\mathbb{C}[(\mathbb{Z}_2)^n], f)$  with symmetric function  $\phi = \delta f$ , the commutation graph completely determines the structure of  $\mathcal{A}$ .*

*Proof.* In the case where  $\phi$  is symmetric, formula (3.2) and Proposition 2.3 imply that the graph determines the structure of the algebra  $\mathcal{A}$ , up to signature.  $\square$

This means, two complex algebras,  $\mathcal{A}$  and  $\mathcal{A}'$  corresponding to the same commutation graph are isomorphic. We do not know if the converse statement is true. However, it is easy to show the following. Given two algebras,  $\mathcal{A}$  and  $\mathcal{A}'$  with different commutation graphs, there is no isomorphism of  $\mathcal{A}$  and  $\mathcal{A}'$  that preserves the  $(\mathbb{Z}_2)^n$ -grading.

**Example 4.9.** The algebra  $\mathbb{M}_3$  is the first non-trivial algebra of the series  $\mathbb{M}_n$ . The corresponding commutation graph is presented in Figure 3, together with the graph of the Clifford algebra  $\mathcal{Cl}_3$ .




FIGURE 3. The algebras  $\mathcal{Cl}_3$  and  $\mathbb{M}_3$ .

The algebra  $\mathcal{Cl}_3$  is not simple:  $\mathcal{Cl}_3 = \mathbb{C}[2] \oplus \mathbb{C}[2]$ . It contains a central element  $u_{(1,1,1)}$  corresponding to a “singleton” in Figure 3.

**Remark 4.10.** (a) The defined planar graph is *dual trivalent*, that is, every edge represented by a projective line or a circle, see Figure 3, contains exactly 3 elements. Indeed, any three homogeneous elements  $u_x, u_y$  and  $u_{x+y}$  either commute or anticommute with each other. This follows from the tri-linearity of  $\phi$ .

(b) We also notice that the superposition of the graphs of  $\mathcal{Cl}_n$  and  $\mathbb{M}_n$  is precisely the graph of the algebra  $\mathbb{O}_n$ . We thus obtain the following “formula”:  $\mathcal{Cl} + \mathbb{M} = \mathbb{O}$ .




FIGURE 4. The commutation graph of  $\mathbb{M}_4$ .

**Example 4.11.** The commutation graph of the algebra  $\mathbb{M}_4$  is presented in Figure 4.

The commutation graph of the Clifford algebra  $Cl_4$  is presented in Figure 5. Note that both algebras,  $\mathbb{M}_4$  and  $Cl_4$  are simple. The superposition of the graphs of  $\mathbb{M}_4$  and  $Cl_4$  cancels all the edges from  $(1, 1, 1, 1)$ . Therefore, the element  $(1, 1, 1, 1)$  is a singleton in the graph of the algebra  $\mathbb{O}_4$ . This corresponds to the fact that  $u_{(1,1,1,1)}$  in  $\mathbb{O}_4$  is a central, in particular,  $\mathbb{O}_4$  is not simple.




FIGURE 5. The commutation graph of  $Cl_4$ .

The planar graph provides a nice way to visualize the algebra  $(\mathbb{K}[(\mathbb{Z}_2)^n], f)$ .

### 5. GENERATING FUNCTIONS

In this section we prove Theorem 1 and its corollaries. Our main tool is the notion of generating function. We show that the structure of all the algebras we consider in this paper is determined (up to signature) by a single function of one argument  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$ . This of course simplifies the understanding of these algebras.

**5.1. Existence of a generating function.** Given a  $(\mathbb{Z}_2)^n$ -graded quasialgebra, let us prove that there exists a generating function  $\alpha$  if and only if the ternary map  $\phi$  is symmetric.

The condition that  $\phi$  is symmetric is of course necessary for existence of  $\alpha$ , cf. formula (2.11), let us prove that this condition is, indeed, sufficient.

**Lemma 5.1.** *If  $\phi$  is symmetric, then  $\beta$  is a 2-cocycle:  $\delta\beta = 0$ .*

*Proof.* If  $\phi$  is symmetric then the identity (3.2) is satisfied. In particular, the sum of the two expressions of  $\phi$  gives:

$$\beta(x + y, z) + \beta(x, y + z) + \beta(x, y) + \beta(y, z) = 0$$

which is nothing but the 2-cocycle condition  $\delta\beta = 0$ . □

Using the information about the second cohomology space  $H^2((\mathbb{Z}_2)^n; \mathbb{Z}_2)$ , as in the proof of Proposition 2.3, we deduce that  $\beta$  is of the form

$$\beta(x, y) = \delta\alpha(x, y) + \sum_{i \in I} x_i y_i + \sum_{(k, \ell) \in J} x_k y_\ell,$$

where  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  is an arbitrary function and where  $I$  is a subset of  $\{1, \dots, n\}$  and  $J$  is a subset of  $\{(k, \ell) \mid k < \ell\}$ . Indeed, the second and the third terms are the most general non-trivial 2-cocycles on  $(\mathbb{Z}_2)^n$  with coefficients in  $\mathbb{Z}_2$ .

Furthermore, the function  $\beta$  satisfies two properties: it is symmetric and  $\beta(x, x) = 0$ . The second property implies that  $\beta$  does not contain the terms  $x_i y_i$ . The symmetry of  $\beta$  means that whenever there is a term  $x_k y_\ell$ , there is  $x_\ell y_k$ , as well. But,  $x_k y_\ell + x_\ell y_k$  is a coboundary of  $x_k x_\ell$ . We proved that  $\beta = \delta\alpha$ , which is equivalent to the identity (2.10).

Finally, using the equality (3.2), we also obtain the identity (2.11).

Theorem 1 is proved.

**5.2. Generating functions are cubic.** In this section, we prove Proposition 3.1. We show that a function  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  is a generating function of a  $(\mathbb{Z}_2)^n$ -graded quasialgebra if and only if  $\alpha$  is a polynomial of degree  $\leq 3$ .

The next statement is an application of the pentagonal diagram in Figure 2.

**Lemma 5.2.** *A generating function  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  satisfies the equation  $\delta_3\alpha = 0$ , where the map  $\delta_3$  is defined by*

$$(5.1) \quad \begin{aligned} \delta_3\alpha(x, y, z, t) := & \alpha(x + y + z + t) \\ & + \alpha(x + y + z) + \alpha(x + y + t) + \alpha(x + z + t) + \alpha(y + z + t) \\ & + \alpha(x + y) + \alpha(x + z) + \alpha(x + t) + \alpha(y + z) + \alpha(y + t) + \alpha(z + t) \\ & + \alpha(x) + \alpha(y) + \alpha(z) + \alpha(t). \end{aligned}$$

*Proof.* This follows immediately from the fact that  $\phi$  is a 3-cocycle: substitute (2.11) to the equation  $\delta\phi = 0$  to obtain  $\delta_3\alpha = 0$ .  $\square$

The following statement characterizes polynomials of degree  $\leq 3$ .

**Lemma 5.3.** *A function  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  is a polynomial of degree  $\leq 3$  if and only if  $\delta_3\alpha = 0$ .*

*Proof.* This is elementary, see also [28, 13].  $\square$

Proposition 3.1 is proved.

Let us now prove Corollary 3.2. If the map  $\phi$  is symmetric, then Theorem 1 implies the existence of the generating function  $\alpha$ . The map  $\phi$  is then given by (2.11). One checks by an elementary calculation that

$$\phi(x + y, z, t) + \phi(x, z, t) + \phi(y, z, t) = \delta_3\alpha(x, y, z, t).$$

By Lemma 5.2, one has  $\delta_3\alpha = 0$ . It follows that  $\phi$  is trilinear.

Furthermore, from (3.2), we deduce that  $\phi$  is alternate.

Corollary 3.2 is proved.

**5.3. Uniqueness of the generating function.** Let us show that there is a canonical way to choose a generating function.

**Lemma 5.4.** (i) *Given a  $(\mathbb{Z}_2)^n$ -graded quasialgebra  $\mathcal{A}$  with a generating function, one can choose the generating function in such a way that it satisfies*

$$(5.2) \quad \begin{cases} \alpha(0) = 0, \\ \alpha(x) = 1, \quad |x| = 1. \end{cases}$$

(ii) *The generating function normalized by (5.2) is unique.*

*Proof.* Part (i). Every generating function  $\alpha$  vanishes on the zero element  $0 = (0, \dots, 0)$ , cf. Section 2.6. Furthermore, a generating function corresponding to a given algebra  $\mathcal{A}$ , is defined up to a 1-cocycle on  $(\mathbb{Z}_2)^n$ . Indeed, the functions  $\beta = \delta\alpha$  and  $\phi = \delta_2\alpha$  that define the quasialgebra structure do not change if one adds a 1-cocycle to  $\alpha$ . Since every 1-cocycle is a linear function, we obtain

$$\alpha(x) \sim \alpha(x) + \sum_{1 \leq i \leq n} \lambda_i x_i.$$

One therefore can normalize  $\alpha$  in such a way that  $\alpha(x) = 1$  for all  $x$  such that  $|x| = 1$ .

Part (ii). The generating function normalized in this way is unique. Indeed, any other function, say  $\alpha'$ , satisfying (5.2) differs from  $\alpha$  by a polynomial of degree  $\geq 2$ , so that  $\alpha - \alpha'$  cannot be a 1-cocycle. Therefore,  $\beta' \neq \beta$  which means the quasialgebra structure is different.  $\square$

We will assume the normalization (5.2) in the sequel, whenever we speak of *the* generating function corresponding to a given algebra.

Let us now consider an algebra  $\mathcal{A}$  with  $n$  generators  $u_1, \dots, u_n$ . The group of permutations  $\mathfrak{S}_n$  acts on  $\mathcal{A}$  by permuting the generators.

**Corollary 5.5.** *If the group of permutations  $\mathfrak{S}_n$  acts on  $\mathcal{A}$  by automorphisms, then the corresponding generating function  $\alpha$  is  $\mathfrak{S}_n$ -invariant.*

*Proof.* Let  $\alpha$  be a generating function. Since the algebra  $\mathcal{A}$  is stable with respect to the  $\mathfrak{S}_n$ -action, the function  $\alpha \circ \sigma$  is again a generating function. If, moreover,  $\alpha$  satisfies (5.2), then  $\alpha \circ \sigma$  also satisfies this condition. The uniqueness Lemma 5.4 implies that  $\alpha \circ \sigma = \alpha$ .  $\square$

**5.4. From the generating function to the twisting function.** Given an arbitrary polynomial  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  of  $\deg \alpha \leq 3$  such that  $\alpha(0) = 0$ , we will give an explicit formula for a twisting function  $f$  such that  $(\mathbb{K}[(\mathbb{Z}_2)^n], f)$  admits  $\alpha$  as a generating function.

The procedure is linear, we associate to every monomial

$$(5.3) \quad \begin{aligned} x_i x_j x_k &\longmapsto x_i x_j y_k + x_i y_j x_k + y_i x_j x_k \\ x_i x_j &\longmapsto x_i y_j, \\ x_i &\longmapsto x_i y_i \end{aligned}$$

where  $i < j < k$ . The following statement is straightforward.

**Proposition 5.6.** *The formula (5.3) defines the unique twisting function  $f$  satisfying the property*

$$(5.4) \quad f(x, x) = \alpha(x).$$

## 6. PROOF OF THE SIMPLICITY CRITERION

In this section, we prove Theorem 2. We use the notation  $\mathcal{A}$  to refer to any of the algebras  $\mathbb{O}_n, \mathbb{O}_{p,q}$  and  $\mathbb{M}_n, \mathbb{M}_{p,q}$ .

**6.1. The idea of the proof.** Our proof of simplicity of a twisted group algebra  $\mathcal{A}$  will be based on the following lemma.

**Lemma 6.1.** *If for every homogeneous element  $u_x$  in  $\mathcal{A}$  there exists an element  $u_y$  in  $\mathcal{A}$  such that  $u_x$  and  $u_y$  anticommute, then  $\mathcal{A}$  is simple.*

*Proof.* Let us suppose that there exists a nonzero proper two-sided ideal  $\mathcal{I}$  in  $\mathcal{A}$ . Every element  $u$  in  $\mathcal{I}$  is a linear combinations of some homogeneous elements of  $\mathcal{A}$ . We write

$$u = \lambda_1 u_{x_1} + \cdots + \lambda_k u_{x_k}.$$

Among all the elements of  $\mathcal{I}$  we choose an element such that the number  $k$  of homogeneous components is the smallest possible. We can assume that  $k \geq 2$ , otherwise  $u$  is homogeneous and therefore  $u^2$  is non-zero and proportional to 1, so that  $\mathcal{I} = \mathcal{A}$ . In addition, up to multiplication by  $u_{x_1}$  and scalar normalization we can assume that

$$u = 1 + \lambda_2 u_{x_2} + \cdots + \lambda_k u_{x_k}.$$

If there exists an element  $u_y \in \mathcal{A}$  anticommuting with  $u_{x_2}$  then one obtains that  $u \cdot u_y - u_y \cdot u$  is a nonzero element in  $\mathcal{I}$  with a shorter decomposition into homogeneous components. This is a contradiction with the choice of  $u$ . Therefore,  $\mathcal{A}$  has no proper ideal.  $\square$

We now need to study central elements in  $\mathcal{A}$ , i.e., the elements commuting with every element of  $\mathcal{A}$ .

**6.2. Central elements.** In this section we study the center  $\mathcal{Z}(\mathcal{A})$  of  $\mathcal{A}$ , i.e.,

$$\mathcal{Z}(\mathcal{A}) = \{w \in \mathcal{A} \mid w \cdot a = a \cdot w, \text{ for all } a \in \mathcal{A}\}.$$

The unit 1 of  $\mathcal{A}$  is obviously an element of the center. We say that  $\mathcal{A}$  has a trivial center if  $\mathcal{Z}(\mathcal{A}) = \mathbb{K}1$ .

Consider the following particular element

$$z = (1, \dots, 1)$$

in  $(\mathbb{Z}_2)^n$ , with all the components equal to 1, and the associated homogeneous element  $u_z$  in  $\mathcal{A}$ .

**Lemma 6.2.** *The element  $u_z$  in  $\mathcal{A}$  is central if and only if*

- (1)  $n = 4m$  in the cases  $\mathcal{A} = \mathbb{O}_n, \mathbb{O}_{p,q}$ ;
- (2)  $n = 4m + 2$  in the cases  $\mathcal{A} = \mathbb{M}_n, \mathbb{M}_{p,q}$ .

*Proof.* The element  $u_z$  in  $\mathcal{A}$  is central if and only if for all  $y \in (\mathbb{Z}_2)^n$  one has  $\beta(y, z) = 0$ . We use the generating function  $\alpha$ . Recall that  $\beta(y, z) = \alpha(y+z) + \alpha(y) + \alpha(z)$ . The value  $\alpha(x)$  depends only on the weight  $|x|$ , see Table (3.7). For every  $y$  in  $\mathbb{Z}_2^n$ , one has

$$|z + y| = |z| - |y|.$$

Case (1). According to the Table (3.7), one has  $\alpha(x) = 0$  if and only if  $|x|$  is a multiple of 4.

Assume  $n = 4m$ . One gets  $\alpha(z) = 0$  and for every  $y$  one has  $\alpha(y) = 0$  if and only if  $\alpha(y + z) = 0$ . So, in that case, one always has  $\alpha(y) = \alpha(y + z)$  and therefore  $\beta(y, z) = 0$ .

Assume  $n = 4m + r$ ,  $r = 1, 2$  or  $3$ . We can always choose an element  $y$  such that  $|y| = |r - 2| + 1$ . We get

$$\alpha(z) = \alpha(y) = \alpha(y + z) = 1.$$

Hence,  $\beta(y, z) = 1$ . This implies that  $u_z$  is not central.

Case (2). According to the Table (3.7), one has  $\alpha(x) = 0$  if and only if  $|x|$  is not equal to  $1 \pmod 4$ .

Assume  $n = 4m + 2$ . One gets  $\alpha(z) = 0$  and for every  $y$  one has  $|y| = 1 \pmod 4$  if and only if  $|y + z| = 1 \pmod 4$ . So, in that case, one always has  $\alpha(y) = \alpha(y + z)$  and therefore  $\beta(y, z) = 0$ .

Assume  $n = 4m + r$ ,  $r = 0, 1$  or  $3$ . We choose the element  $y = (1, 0, \dots, 0)$ , if  $r = 0, 3$ , or  $y = (1, 1, 0, \dots, 0)$ , if  $r = 1$ . We easily compute  $\beta(y, z) = 1$ . This implies that  $u_z$  is not central.  $\square$

Let us consider the case where  $u_z$  is not central.

**Lemma 6.3.** *If  $u_z$  is not central, then  $\mathcal{A}$  has a trivial center.*

*Proof.* It suffices to prove that for every homogeneous element  $u_x$  in  $\mathcal{A}$ , that is not proportional to 1, there exists an element  $u_y$  in  $\mathcal{A}$ , such that  $u_x$  and  $u_y$  anticommute. Indeed, if  $u$  is central, then each homogeneous component of  $u$  is central.

Let us fix  $x \in (\mathbb{Z}_2)^n$  and the corresponding homogeneous element  $u_x \in \mathcal{A}$ , such that  $x$  is neither 0, nor  $z$ . We want to find an element  $y \in (\mathbb{Z}_2)^n$  such that  $\beta(x, y) = 1$  or equivalently  $u_x$  anticommutes with  $u_y$ . Using the invariance of the functions  $\alpha$  and  $\beta$  under permutations of the coordinates, we can assume that  $x$  is of the form

$$x = (1, \dots, 1, 0, \dots, 0),$$

where first  $|x|$  entries are equal to 1 and the last entries are equal to 0. We assume  $0 < |x| < n$ , so that,  $x$  starts with 1 and ends by 0.

**Case  $\mathcal{A} = \mathbb{O}_n$  or  $\mathbb{O}_{p,q}$ .** If  $|x| \neq 4\ell$ , then we use exactly the same arguments as in the proof of Lemma 6.2 in order to find a suitable  $y$  (one can also take one of the elements  $y = (1, 0, \dots, 0)$  or  $y = (0, \dots, 0, 1)$ ). Assume  $|x| = 4\ell$ . Consider the element

$$y = (0, 1, \dots, 1, 0, \dots, 0),$$

with  $|y| = |x|$ . One has  $\alpha(x) = \alpha(y) = 0$  and  $\alpha(x + y) = 1$ . So we also have  $\beta(x, y) = 1$  and deduce  $u_x$  anticommutes with  $u_y$ .

**Case  $\mathcal{A} = \mathbb{M}_n$  or  $\mathbb{M}_{p,q}$ .** Similarly to the proof of Lemma 6.2, if  $k \neq 4\ell + 2$  then we can find a  $y$  such that  $u_y$  anticommutes with  $u_x$ . If  $k = 4\ell + 2$  then  $\alpha(x) = 0$ . The element  $y = (0, \dots, 0, 1)$  satisfies  $\alpha(y) = 1$  and  $\alpha(x + y) = 0$ .  $\square$

Consider now the case where  $u_z$  is a central element. There are two different possibilities:  $u_z^2 = 1$ , or  $u_z^2 = -1$ .

**Lemma 6.4.** *If  $u_z \in \mathcal{A}$  is a central element and if  $u_z^2 = 1$ , then the algebra splits into a direct sum of two subalgebras:*

$$\mathcal{A} = \mathcal{A}^+ \oplus \mathcal{A}^-,$$

where  $\mathcal{A}^+ := \mathcal{A} \cdot (1 + u_z)$  and  $\mathcal{A}^- := \mathcal{A} \cdot (1 - u_z)$ .

*Proof.* Using  $u_z^2 = 1$ , one immediately obtains

$$(6.1) \quad \begin{aligned} (1 \pm u_z)^2 &= 2(1 \pm u_z), \\ (1 + u_z) \cdot (1 - u_z) &= 0. \end{aligned}$$

In addition, using the expression of  $\phi$  in terms of  $\beta$  given in (3.2) and the fact that  $\beta(\cdot, z) = 0$ , one deduces that  $\phi(\cdot, \cdot, z) = 0$  and thus  $a \cdot (b \cdot u_z) = (a \cdot b) \cdot u_z$  for all  $a, b \in \mathcal{A}$ . It follows that

$$(6.2) \quad (a \cdot (1 \pm u_z)) \cdot (b \cdot (1 \pm u_z)) = (a \cdot b) \cdot ((1 \pm u_z) \cdot (1 \pm u_z))$$

for all  $a, b \in \mathcal{A}$ . This expression, together with the above computations (6.1), shows that  $\mathcal{A}^+$  and  $\mathcal{A}^-$  are, indeed, two subalgebras of  $\mathcal{A}$  and that they satisfy  $\mathcal{A}^+ \cdot \mathcal{A}^- = \mathcal{A}^- \cdot \mathcal{A}^+ = 0$ . Moreover, for any  $a \in \mathcal{A}$ , one can write

$$a = \frac{1}{2} a \cdot (1 + u_z) + \frac{1}{2} a \cdot (1 - u_z).$$

This implies the direct sum decomposition  $\mathcal{A} = \mathcal{A}^+ \oplus \mathcal{A}^-$ .  $\square$

Notice that the elements  $\frac{1}{2}(1 + u_z)$  and  $\frac{1}{2}(1 - u_z)$  are the units of  $\mathcal{A}^+$  and  $\mathcal{A}^-$ , respectively.

**6.3. Proof of Theorem 2, part (i).** If  $n \neq 4m$ , then by Lemma 6.1 and Lemma 6.3 we immediately deduce that  $\mathbb{O}_n$  is simple.

If  $n = 4m$ , then  $u_z$  is central and, in the complex case, one has  $u_z^2 = 1$ . By Lemma 6.2 and Lemma 6.4, we immediately deduce that  $\mathbb{O}_n$  is not simple and one has

$$\mathbb{O}_{4m} = \mathbb{O}_{4m} \cdot (1 + u_z) \oplus \mathbb{O}_{4m} \cdot (1 - u_z),$$

where  $z = (1, \dots, 1) \in (\mathbb{Z}_2)^n$ . It remains to show that the algebras  $\mathbb{O}_{4m-1}$  and  $\mathbb{O}_{4m} \cdot (1 \pm u_z)$  are isomorphic. Indeed, using the computations (6.1) and (6.2), one checks that the map

$$u_x \mapsto \frac{1}{2} u_{(x,0)} \cdot (1 \pm u_z),$$

where  $x \in (\mathbb{Z}_2)^{n-1}$ , is the required isomorphism.

The proof in the case of  $\mathbb{M}_n$  is completely similar.

**6.4. Proof of Theorem 2, part (ii).** The algebras  $\mathbb{O}_{p,q}$  with  $p+q \neq 4m$  and the algebras  $\mathbb{M}_{p,q}$  with  $p+q \neq 4m+2$  are simple because their complexifications are.

If now  $u_z$  is central, then the property  $u_z^2 = 1$  or  $-1$  becomes crucial. Using the expressions for  $f_{\mathbb{O}}$  or  $f_{\mathbb{M}}$ , one computes

$$\begin{aligned} f_{\mathbb{O}_{p,q}}(z, z) &= \sum_{i < j < k} z_i z_j z_k + \sum_{i \leq j} z_i z_j + \sum_{1 \leq i \leq p} z_i \\ &= \frac{n(n-1)(n-2)}{6} + \frac{n(n+1)}{2} + p \\ &= p, \quad \text{mod } 2. \end{aligned}$$

And similarly, one obtains  $f_{\mathbb{M}_{p,q}}(z, z) = p$ . It follows that  $u_z^2 = (-1)^p$ .

If  $p$  is even, then Lemma 6.2 just applied guarantees that  $\mathcal{A}$  is not simple.

Finally, if  $u_z$  is central and  $p$  is odd, then  $u_z^2 = -1$ .

**Lemma 6.5.** *If  $u_z$  is central and  $p$  is odd, then*

$$\mathbb{O}_{p,q} \cong \mathbb{O}_{p,q-1} \otimes \mathbb{C}, \quad \mathbb{M}_{p,q} \cong \mathbb{M}_{p,q-1} \otimes \mathbb{C}.$$

*Proof.* We construct an explicit isomorphism from  $\mathbb{O}_{p,q-1} \otimes \mathbb{C}$  to  $\mathbb{O}_{p,q}$  as follows.

$$\begin{aligned} u_x \otimes 1 &\longmapsto u_{(x,0)} \\ u_x \otimes \sqrt{-1} &\longmapsto u_{(x,0)} \cdot u_z, \end{aligned}$$

for all  $x \in (\mathbb{Z}_2)^{n-1}$ . We check that the above map is indeed an isomorphism of algebras by noticing that  $f_{\mathbb{O}_{p,q}}((x,0), (y,0)) = f_{\mathbb{O}_{p,q-1}}(x, y)$ .  $\square$

Let us show that Lemma 6.5 implies that the (real) algebras  $\mathbb{O}_{p,q}$  with  $p+q = 4m$  and  $p$  odd and the algebras  $\mathbb{M}_{p,q}$  with  $p+q = 4m+2$  and  $p$  odd are simple. Indeed,

$$\mathbb{O}_{p,q-1} \otimes \mathbb{C} \cong \mathbb{O}_{p+q-1}^{\mathbb{R}}$$

viewed as a real algebras. We then use the following well-known fact. A simple unital complex algebra viewed as a real algebra remains simple.

The proof of Theorem 2 is complete.

Lemma 6.5 also implies Corollary 4.2.

## 7. HURWITZ-RADON SQUARE IDENTITIES

In this section, we use the algebras  $\mathbb{O}_n$  (and, in the real case,  $\mathbb{O}_{0,n}$ ) to give explicit formulæ for solutions of a classical problem of products of squares. Recall, that the octonion algebra is related to the 8-square identity. In an arbitrary commutative ring, the product  $(a_1^2 + \cdots + a_8^2)(b_1^2 + \cdots + b_8^2)$  is again a sum of 8 squares  $c_1^2 + \cdots + c_8^2$ , where  $c_k$  are explicitly given by bilinear forms in  $a_i$  and  $b_j$  with coefficients  $\pm 1$ , see, e.g., [11]. This identity is equivalent to the fact that  $\mathbb{O}$  is a composition algebra, that is, for any  $a, b \in \mathbb{O}$ , the norm of the product is equal to the product of the norms:

$$(7.1) \quad \mathcal{N}(a \cdot b) = \mathcal{N}(a)\mathcal{N}(b).$$

Hurwitz proved that there is no similar  $N$ -square identity for  $N > 8$ , as there is no composition algebra in higher dimensions.

The celebrated Hurwitz-Radon Theorem [18, 24] establishes the maximal number  $r$ , as function of  $N$ , such that there exists an identity

$$(7.2) \quad (a_1^2 + \cdots + a_N^2)(b_1^2 + \cdots + b_r^2) = (c_1^2 + \cdots + c_N^2),$$

where  $c_k$  are bilinear forms in  $a_i$  and  $b_j$ . The theorem states that  $r = \rho(N)$  is the maximal number, where  $\rho(N)$  is the Hurwitz-Radon function defined as follows. Write  $N$  in the form  $N = 2^{4m+\ell} N'$ , where  $N'$  is odd and  $\ell = 0, 1, 2$  or  $3$ , then

$$\rho(N) := 8m + 2^\ell.$$

It was proved by Gabel [15] that the bilinear forms  $c_k$  can be chosen with coefficients  $\pm 1$ . Note that the only interesting case is  $N = 2^n$  since the general case is an immediate corollary of this particular one. We refer to [25, 27] for the history, results and references.

In this section, we give explicit formulæ for the solution to the Hurwitz-Radon equation.

**7.1. The explicit solution.** We give explicit solution for Hurwitz-Radon equation (7.2) for any  $N = 2^n$  with  $n$  not a multiple of 4.

We label the  $a$ -variables and the  $c$ -variables by elements of  $(\mathbb{Z}_2)^n$ . In order to describe the labeling of the  $b$ -variables, we consider the following particular elements of  $(\mathbb{Z}_2)^n$ :

$$\begin{aligned} e_0 &:= (0, 0, \dots, 0), \\ \bar{e}_0 &:= (1, 1, \dots, 1), \\ e_i &:= (0, \dots, 0, 1, 0, \dots, 0), \text{ where } 1 \text{ occurs at the } i\text{-th position,} \\ \bar{e}_i &:= (1, \dots, 1, 0, 1, \dots, 1), \text{ where } 0 \text{ occurs at the } i\text{-th position,} \end{aligned}$$

for all  $1 \leq i \leq n$  and  $1 < j \leq n$ . We then introduce the following subset  $H_n$  of  $(\mathbb{Z}_2)^n$ :

$$(7.3) \quad \begin{aligned} H_n &= \{e_i, \bar{e}_i, 1 \leq i \leq n\}, \text{ for } n \equiv 1 \pmod{4}, \\ H_n &= \{e_i, e_1 + e_j, 0 \leq i \leq n, 1 < j \leq n\}, \text{ for } n \equiv 2 \pmod{4}, \\ H_n &= \{e_i, \bar{e}_i, 0 \leq i \leq n\}, \text{ for } n \equiv 3 \pmod{4}. \end{aligned}$$

In each case, the subset  $H_n$  contains exactly  $\rho(2^n)$  elements.

We write the Hurwitz-Radon identity in the form

$$\left( \sum_{x \in (\mathbb{Z}_2)^n} a_x^2 \right) \left( \sum_{x \in H_n} b_x^2 \right) = \sum_{x \in (\mathbb{Z}_2)^n} c_x^2.$$

We will establish the following.

**Theorem 3.** *The bilinear forms*

$$(7.4) \quad c_x = \sum_{y \in H_n} (-1)^{f_{\mathbb{O}}(x+y, y)} a_{x+y} b_y,$$

where  $f_{\mathbb{O}}$  is the twisting function of the algebra  $\mathbb{O}_n$  defined in (1.3), are a solution to the Hurwitz-Radon identity.

In order to prove Theorem 3 we will need to define the natural norm on  $\mathbb{O}_n$ .

**7.2. The Euclidean norm.** Assume that a twisted group algebra  $\mathcal{A} = (\mathbb{K}[(\mathbb{Z}_2)^n], f)$  is equipped with a generating function  $\alpha$ . Assume furthermore that the twisting function satisfies  $f(x, x) = \alpha(x)$ , as in (5.4).

The anti-involution on  $\mathcal{A}$  is defined for every  $a = \sum_{x \in (\mathbb{Z}_2)^n} a_x u_x$ , where  $a_x \in \mathbb{C}$  (or in  $\mathbb{R}$ ) are scalar coefficients and  $u_x$  are the basis elements, by the formula

$$\bar{a} = \sum_{x \in (\mathbb{Z}_2)^n} (-1)^{\alpha(x)} a_x u_x.$$

We then define the following norm of an element  $a \in \mathcal{A}$ :

$$\mathcal{N}(a) := (a \cdot \bar{a})_0.$$

**Proposition 7.1.** *The above norm is nothing but the Euclidean norm in the standard basis:*

$$(7.5) \quad \mathcal{N}(a) = \sum_{x \in (\mathbb{Z}_2)^n} a_x^2.$$

*Proof.* One has:

$$\mathcal{N}(a) = \sum (-1)^{\alpha(x)} a_x^2 u_x \cdot u_x = \sum (-1)^{\alpha(x)+f(x,x)} a_x^2.$$

The result then follows from the assumption  $f(x, x) = \alpha(x)$ .  $\square$

The following statement is a general criterion for  $a, b \in \mathcal{A}$  to satisfy the composition equation (7.1). This criterion will be crucial for us to establish the square identities.

**Proposition 7.2.** *Elements  $a, b \in \mathcal{A}$  satisfy (7.1), if and only if for all  $x, y, z, t \in (\mathbb{Z}_2)^2$  such that*

$$x + y + z + t = 0, \quad (x, y) \neq (z, t), \quad a_x b_y a_z b_t \neq 0,$$

*one has  $\alpha(x + z) = \alpha(y + t) = 1$ .*

*Proof.* Calculating the left-hand-side of (7.1), we obtain

$$\mathcal{N}(a \cdot b) = \sum_{x+y+z+t=0} (-1)^{f(x,y)+f(z,t)} a_x b_y a_z b_t$$

According to (7.5), the product of the norm in the right-hand-side is:

$$\mathcal{N}(a)\mathcal{N}(b) = \sum_{x,y} a_x^2 b_y^2.$$

It follows that the condition (7.1) is satisfied if and only if

$$f(x, y) + f(z, t) + f(x, t) + f(z, y) = 1,$$

whenever  $(x, y) \neq (z, t)$  and  $a_x b_y a_z b_t \neq 0$ .

Taking into account the linearity of the function (5.3) and substituting  $t = x + y + z$ , one finally gets (after cancellation):

$$f(z, x) + f(x, z) + f(x, x) + f(z, z) = 1.$$

In terms of the function  $\alpha$  this is exactly the condition  $\alpha(x + z) = 1$ . Hence the result.  $\square$

**7.3. Proof of Theorem 3.** Let us apply Proposition 7.2 to the case of the algebra  $\mathbb{O}_n$ .

Given the variables  $(a_x)_{x \in (\mathbb{Z}_2)^n}$  and  $(b_x)_{x \in H_n}$ , where  $H_n$  is the subset defined in (7.3), form the following vectors in  $\mathbb{O}_n$ ,

$$a = \sum_{x \in (\mathbb{Z}_2)^n} a_x u_x, \quad b = \sum_{y \in H_n} b_y u_y.$$

Taking two distinct elements  $y, t \in H_n$  one always has  $\alpha_{\mathbb{O}}(y + t) = 1$ . Therefore, from Proposition 7.2 one deduces that  $\mathcal{N}(a)\mathcal{N}(b) = \mathcal{N}(a \cdot b)$ . Writing this equality in terms of coordinates of the three elements  $a, b$  and  $c = a \cdot b$ , one obtains the result.

Theorem 3 is proved.

**Example 7.3.** The most obvious choice of two elements  $a, b \in \mathbb{O}_n$  that satisfy the condition (7.1) is:  $a = a_0 u_0 + \sum a_i u_i$  and  $b = b_0 u_0 + \sum b_i u_i$ . One immediately obtains in this case the following elementary but elegant identity:

$$(a_0^2 + \cdots + a_n^2)(b_0^2 + \cdots + b_n^2) = (a_0 b_0 + \cdots + a_n b_n)^2 + \sum_{0 \leq i < j \leq n} (a_i b_j - b_j a_i)^2,$$

for an arbitrary  $n$ , known as the Lagrange identity.



**An explicit formula for the Parker loop.** Let us now give the generating function of the Parker loop. We identify the Golay code with the space  $(\mathbb{Z}_2)^{12}$ , in such a way that the  $i$ -th row of the matrix  $G$  is identified with the  $i$ -th standard basic vector  $(0, \dots, 0, 1, 0, \dots, 0)$  of  $(\mathbb{Z}_2)^{12}$ . The coordinates of a vector  $x \in (\mathbb{Z}_2)^{12}$  are denoted by  $(x_1, \dots, x_{11}, x_{12})$ , note that the last basis vector  $u_{12}$  is special.

**Proposition 8.2.** *The Parker loop is given by the following generating function  $\alpha$  from  $(\mathbb{Z}_2)^{12}$  to  $\mathbb{Z}_2$ .*

$$\begin{aligned} \alpha_G(x) = & \sum_{1 \leq i \leq 11} x_i x_{i+1} (x_{i+5} + x_{i+8} + x_{i+9}) + x_i x_{i+2} (x_{i+6} + x_{i+8}) \\ & + x_{12} \left( \sum_{1 \leq i \leq 11} x_i + \sum_{1 \leq i < j \leq 11} x_i x_j \right), \end{aligned}$$

where the indices of  $x_{i+k}$  are understood modulo 11.

*Proof.* The ternary function  $\delta f(x, y, z)$  is obviously symmetric in  $x, y, z$ . Theorem 1 then implies the existence of a generating function. This function is a 3-rd order polynomial, by Proposition 3.1. It follows that the function  $\alpha_G$  is completely determined by the second- and third-order relations between the generators that are very easy to check.  $\square$

The explicit formula for the factor set  $f$  in coordinates on  $(\mathbb{Z}_2)^{12}$  is immediately obtained by (5.3). Note that the signature in this case is  $(11, 1)$ , so that we have to add  $x_{12}y_{12}$  to (5.3).

**Remark 8.3.** The difference between the loops generated by the basis elements of  $\mathbb{O}_n$  and  $\mathbb{M}_n$  and the Parker loop is that the function  $\alpha_G$  is not  $\mathfrak{S}_n$ -invariant. Our classification results cannot be applied in this case.

APPENDIX: LINEAR ALGEBRA AND DIFFERENTIAL CALCULUS OVER  $\mathbb{Z}_2$

This subject is an amazing mixture of linear algebra and elementary group cohomology theory. The purpose of this Appendix is to relate the algebraic problems we study to the general framework of linear algebra over  $\mathbb{Z}_2$  which is a classical domain.

**Automorphisms of  $(\mathbb{Z}_2)^n$  and linear equivalence.** All the algebraic structures on  $(\mathbb{Z}_2)^n$  we consider are invariant with respect to the action of the group automorphisms

$$\text{Aut}((\mathbb{Z}_2)^n) \cong \text{GL}(n, \mathbb{Z}_2).$$

For instance, the generating function  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$ , as well as  $\beta$  and  $\phi$ , are considered up to the  $\text{Aut}((\mathbb{Z}_2)^n)$ -equivalence (called “congruence” in the classic literature [2]).

**Quadratic forms.** Obviously but remarkably, the notion of quadratic form on  $(\mathbb{Z}_2)^n$  with coefficients in  $\mathbb{Z}_2$  coincides with the notion of bilinear skew-symmetric form (often called alternate, cf. [2]). Indeed, for an arbitrary quadratic form  $\alpha$ , the corresponding polar bilinear form  $\beta = \delta\alpha$  satisfies  $\beta(x, x) = 0$ , while symmetry and skew-symmetry over  $\mathbb{Z}_2$  is the same condition.

It is very easy to prove the following statement which is a version of the (linear) Darboux theorem (see [2] and also [26] for the details). *There exists only one (up to equivalence) non-degenerate quadratic form on  $(\mathbb{Z}_2)^{2m}$ :*

$$(8.1) \quad \alpha(x) = x_1x_{m+1} + \cdots + x_mx_{2m}.$$

*There are no non-degenerate quadratic forms on  $(\mathbb{Z}_2)^{2m+1}$ .*

The generating function for a Clifford algebra with  $n$  generators, see formula (2.13), is nothing but the same quadratic form written in a different basis. The two quadratic forms are therefore equivalent, but each of the formulæ, (2.13) and (8.1), has its merits. The formula (8.1) is very simple and written in the standard Darboux basis; the formula (2.13) is invariant with respect to the action of the symmetric group  $\mathfrak{S}_n$ .

The normal form (8.1) has various algebraic corollaries. Let us give one example: we immediately obtain the well-known factorization of the complex Clifford algebras:

$$Cl_{2m} \cong Cl_2^{\otimes m} \cong \mathbb{C}[2^m],$$

where  $\mathbb{C}[2^m]$  are  $(2^m \times 2^m)$ -matrices. Indeed, the function (8.1) is nothing but the sum of  $m$  generating functions of  $Cl_2$ . The other classical symmetry and periodicity theorems for the Clifford algebras can also be deduced in this way.

Let us mention that bilinear forms over  $\mathbb{Z}_2$  is still a subject for nice research development [19].

**Cubic polynomials.** In this paper, we were led to consider polynomials  $\alpha : (\mathbb{Z}_2)^n \rightarrow \mathbb{Z}_2$  of degree 3:

$$\alpha(x) = \sum_{i < j < k} \alpha_{ijk}^3 x_i x_j x_k + \sum_{i < j} \alpha_{ij}^2 x_i x_j,$$

where  $\alpha_{ijk}^3$  and  $\alpha_{ij}^2$  are arbitrary coefficients (equal to 1 or 0). It turns out that it is far of being obvious to understand what means  $\alpha$  is “non-degenerate”.

To every polynomial  $\alpha$ , we associate a binary function  $\beta = \delta\alpha$  and a trilinear form  $\phi = \delta_2\alpha$ , see formula (2.11), which is of course just the polarization (or linearization) of  $\alpha$ . The form  $\phi$  is alternate:  $\phi(x, x, \cdot) = \phi(x, \cdot, x) = \phi(\cdot, x, x) = 0$  and depends only on the homogeneous part of degree 3 of  $\alpha$ , i.e., only on  $\alpha_{ijk}^3$ . There are three different ways to understand the notion of non-degeneracy.

(1) The most naive way:  $\alpha$  (and  $\phi$ ) is non-degenerate if for all linearly independent  $x, y \in (\mathbb{Z}_2)^n$ , the linear function  $\phi(x, y, \cdot) \neq 0$ . One can show that, with this definition, *there are no non-degenerate cubic forms on  $(\mathbb{Z}_2)^n$  for  $n \geq 3$* . This is of course not the way we take.

(2) The second way to understand non-degeneracy is as follows. The trilinear map  $\phi$  itself defines an  $n$ -dimensional algebra. Indeed, identifying  $(\mathbb{Z}_2)^n$  with its dual space, the trilinear function  $\phi$  defines a product  $(x, y) \mapsto \phi(x, y, \cdot)$ . One can say that  $\phi$  (and  $\alpha$ ) is non-degenerate if this algebra is simple. This second way is much more interesting and is

related to many different subjects. For instance, classification of simple Lie (super)algebras over  $\mathbb{Z}_2$  is still an open problem, see [8] and references therein. This definition also depends only on the homogeneous part of of degree 3 of  $\alpha$ .

(3) We understand non-degeneracy yet in a different way. We say that  $\alpha$  is non-degenerate if for all linearly independent  $x, y$  there exists  $z$  such that

$$\beta(x, z) \neq 0, \quad \beta(y, z) = 0,$$

where  $\beta = \delta\alpha$ . This is equivalent to the fact that the algebra with the generated function  $\alpha$  is simple, cf. Section 6.

We believe that every non-degenerate (in the above sense) polynomial of degree 3 on  $(\mathbb{Z}_2)^n$  is equivalent to one of the two forms (3.5) and (3.6). Note that a positive answer would imply the uniqueness results of Section 3.3 without the  $\mathfrak{S}_n$ -invariance assumption.

**Higher differentials.** Cohomology of abelian groups with coefficients in  $\mathbb{Z}_2$  is a well-known and quite elementary theory explained in many textbooks. Yet, it can offer some surprises.

Throughout this work, we encountered and extensively used the linear operators  $\delta_k$ , for  $k = 1, 2, 3$ , cf. (2.11) and (5.1), that associate a  $k$ -cochain on  $(\mathbb{Z}_2)^n$  to a function. These operators were defined in [28], and used in the Moufang loops theory, [17, 13, 23]. Operations of this type are called *natural* or *invariant* since they commute with the action of  $\text{Aut}((\mathbb{Z}_2)^n)$ . The operator  $\delta_k$  fits to the usual understanding of “higher derivation” since the condition  $\delta_k\alpha = 0$  is equivalent to the fact that  $\alpha$  is a polynomial of degree  $\leq k$ .

## REFERENCES

- [1] A. Adem, R. Milgram, Cohomology of finite groups, Fundamental Principles of Math. Sci., **309**, Springer-Verlag, Berlin, 2004.
- [2] A.A. Albert, *Symmetric and alternate matrices in an arbitrary field. I*, Trans. Amer. Math. Soc. **43** (1938), 386–436.
- [3] H. Albuquerque, A. Elduque, J.M. Pérez-Izquierdo, *Alternative quasialgebras*, Bull. Austral. Math. Soc. **63** (2001), 257–268.
- [4] H. Albuquerque, S. Majid, *Quasialgebra structure of the octonions*, J. Algebra **220** (1999), 188–224.
- [5] H. Albuquerque, S. Majid, *Clifford algebras obtained by twisting of group algebras*, J. Pure Appl. Algebra **171** (2002), 133–148.
- [6] J. Baez, *The octonions*, Bull. Amer. Math. Soc. (N.S.) **39** (2002), 145–205
- [7] Ya.G. Berkovich, E.M. Zhmud’, Characters of finite groups, Part 1. Translations of Mathematical Monographs, **172**, AMS, Providence, RI, 1998.
- [8] S. Bouarroudj, P. Grozman, D. Leites, *Classification of finite dimensional modular Lie superalgebras with indecomposable Cartan matrix*, SIGMA, **5** (2009), 63pp.
- [9] S.B. Conlon, *Twisted group algebras and their representations*, J. Austr. Math. Soc. **4** (1964), 152–173.
- [10] J.H. Conway, N.J.A. Sloane, Sphere packings, lattices and groups. Third edition. Springer-Verlag, New York, 1999.
- [11] J.H. Conway, D.A. Smith, On quaternions and octonions: their geometry, arithmetic, and symmetry, A K Peters, Ltd., Natick, MA, 2003.
- [12] H. S. M. Coxeter, *Integral Cayley numbers*, Duke Math. J. **13** (1946). 561–578.
- [13] A. Drápal, P. Vojtechovský, *Symmetric multilinear forms and polarization of polynomials*, Linear Algebra Appl. **431** (2009), no. 5-7, 998–1012.
- [14] A. Elduque, *Gradings on octonions*, J. Alg. **207** (1998), 342–354.
- [15] M.R. Gabel, *Generic orthogonal stably free projectives*, J. Algebra **29** (1974), 477–488.

- [16] E. Goodaire, E. Jespers, M. Polcino, *Alternative loop rings*, North-Holland Publ., Amsterdam, 1996.
- [17] R.L. Griess Jr., *Code Loops*, *J. Alg.* **100** (1986), 224–234.
- [18] A. Hurwitz, *Über die Komposition der quadratischen Formen*, *Math. Ann.* 88 (1923) 1–25.
- [19] A. Lebedev, *Non-degenerate bilinear forms in characteristic 2, related contact forms, simple Lie algebras and superalgebras*, arXiv:0601536v2.
- [20] V. Lychagin, *Colour calculus and colour quantizations*, *Acta Appl. Math.* **41** (1995), 193–226.
- [21] S. Morier-Genoud, V. Ovsienko, *Well, Papa, can you multiply triplets?*, *Math. Intell.* **31** (2009), 1–2.
- [22] S. Morier-Genoud, V. Ovsienko, *Simple graded commutative algebras*, *J. Alg.* **323** (2010), 1649–1664.
- [23] G. Nagy, P. Vojtechovský, *The Moufang loops of order 64 and 81*, *J. Symb. Comp.* **42** (2007), 871–883.
- [24] J. Radon, *Lineare scharen orthogonale Matrizen*, *Abh. Math. Sem. Univ. Hamburg* 1 (1922) 1–14.
- [25] A.R. Rajwade, *Squares*. London Mathematical Society Lecture Note Series, 171. Cambridge
- [26] C.H Sah, *Symmetric bilinear forms and quadratic forms*, *J. Alg.* **20** (1972), 144–160.
- [27] D. Shapiro, *Compositions of quadratic forms*, Walter de Gruyter & Co., Berlin, 2000.
- [28] H. Ward, *Combinatorial polarization*, *Discrete Math.* **26** (2) (1979) 185–197.

SOPHIE MORIER-GENOUD, UNIVERSITÉ PARIS 6, IMJ, EQUIPE ANALYSE ALGÈBRIQUE, UFR 929 DE MATHÉMATIQUES, 175 RUE DU CHEVALERET, 75013 PARIS, FRANCE

VALENTIN OVSIENKO, CNRS, INSTITUT CAMILLE JORDAN, UNIVERSITÉ CLAUDE BERNARD LYON 1, 43 BOULEVARD DU 11 NOVEMBRE 1918, 69622 VILLEURBANNE CEDEX, FRANCE  
*E-mail address:* `sophiemg@math.jussieu.fr`, `ovsienko@math.univ-lyon1.fr`