

Oberwolfach Preprints



OWP 2013 - 28

GERT-MARTIN GREUEL AND NGUYEN HONG DUC

Right Simple Singularities in Positive
Characteristic

Mathematisches Forschungsinstitut Oberwolfach gGmbH
Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

Starting in 2007, the MFO publishes a preprint series which mainly contains research results related to a longer stay in Oberwolfach. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 30 free hard copies (DIN A4, black and white copy) by surface mail.

Of course, the full copy right is left to the authors. The MFO only needs the right to publish it on its website *www.mfo.de* as a documentation of the research work done at the MFO, which you are accepting by sending us your file.

In case of interest, please send a **pdf file** of your preprint by email to *rip@mfo.de* or *owlf@mfo.de*, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany

Tel +49 7834 979 50
Fax +49 7834 979 55
Email admin@mfo.de
URL www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.
Copyright of the content is held by the authors.

RIGHT SIMPLE SINGULARITIES IN POSITIVE CHARACTERISTIC

GERT-MARTIN GREUEL AND NGUYEN HONG DUC

ABSTRACT. We classify isolated singularities $f \in K[[x_1, \dots, x_n]]$, which are simple, i.e. have no moduli, w.r.t. right equivalence, where K is an algebraically closed field of characteristic $p > 0$. For $K = \mathbb{R}$ or \mathbb{C} this classification was initiated by Arnol'd, resulting in the famous ADE-series. The classification w.r.t. contact equivalence for $p > 0$ was done by Greuel and Kröning with a result similar to Arnol'd's. It is surprising that w.r.t. right equivalence and any given $p > 0$ we have only finitely many simple singularities, i.e. there are only finitely many k such that A_k and D_k are right simple, all the others have moduli. We conjecture a similar finiteness result for singularities with an arbitrary number of moduli. A major point of this paper is the generalization of the notion of modality to the algebraic setting, its behaviour under morphisms, and its relations to formal deformation theory. As an application we show that the modality is semicontinuous in any characteristic.

1. INTRODUCTION

We classify isolated singularities $f \in K[[x_1, \dots, x_n]]$, K an algebraically closed field of characteristic $p > 0$, which have no moduli (modality 0) w.r.t. right equivalence, meaning that there are only finitely many right equivalence classes (see Definition 2.3), where f and g are right equivalent, if they differ by a change of coordinates, see Section A.2. These singularities are called right simple, following Arnol'd, who classified right simple singularities for $K = \mathbb{R}$ and \mathbb{C} (cf. [Arn72]). He showed that the simple singularities are exactly the ADE-singularities, i.e. the two infinite series $A_k, k \geq 1$, $D_k, k \geq 4$, and the three exceptional singularities E_6, E_7, E_8 . It turned out later that the ADE-singularities of Arnol'd are also exactly those of modality 0 for contact equivalence. In the late eighties, Greuel and Kröning showed in [GK90] that the contact simple singularities over a field of positive characteristic are again exactly the ADE-singularities but with a few more normal forms in small characteristic.

A classification w.r.t. right equivalence in positive characteristic however, was never considered so far. A surprising fact of our classification is that for any fixed $p > 0$ there exist only finitely many right simple singularities. We conjecture that this is a general fact for right equivalence in positive characteristic (cf. Conjecture 3.5). For example, if $p = 2$ and n is even, there is just one right simple hypersurface,

$$x_1x_2 + x_3x_4 + \dots + x_{n-1}x_n,$$

while for n odd no right simple singularity exist. A table with normal forms for any $n \geq 1$ and any $p > 0$ is given in section 3 (Theorems 3.1 - 3.3). The problem is even interesting for univariate power series ($n = 1$) (see Section 3.1).

Date: December 11, 2013.

In section 2 we give a precise definition of the number of moduli (modality) for families of power series parametrized by an algebraic variety. In fact, we give two definitions of G -modality, both related to the action of an algebraic group G on a variety X and show that they coincide (Propositions A.2 and A.7), a result which is valid in any characteristic. This unifies the arguments used in the classification, avoiding a lot of similar calculations for different cases.

Moreover, we prove that the G -modality is upper semicontinuous for G the right resp. the contact group (Proposition 2.7).

We introduce the notion of G -completeness (Definition 2.9) which suffices to determine the modality, and we generalize the Kas-Schlessinger theorem [KaS72] to deformations (unfoldings) of formal power series over algebraic varieties. The semiuniversal deformation with section of an isolated hypersurface singularity is G -complete for G the right resp. the contact group (see Proposition 2.14). However, in contrast to the complex analytic case, the usual semiuniversal deformation is not sufficient to determine the modality and hence is not G -complete; we have to consider versal deformations with section (cf. Example 2.13).

Acknowledgement. We would like to thank the referees for their careful reading of the manuscript and helpful comments which improved the presentation of this paper. The second author was partly supported by DAAD (Germany), NAFOSTED (Vietnam), and the Oberwolfach Leibniz Fellows programme of the Mathematisches Forschungsinstitut Oberwolfach (Germany).

2. MODALITY

In the sixties V. I. Arnol'd introduced the notion of modality into singularity theory for real and complex hypersurfaces (cf. [AGV85, Part II]), related to Riemann's idea of moduli for Riemann surfaces. The purpose of this section is to make the notion of modality precise in the case of hypersurface singularities over an algebraically closed field K of arbitrary characteristic and relate it to deformation theory. We investigate right (resp. contact) unfoldings of a formal power series over algebraic varieties and define the modality w.r.t. unfoldings. We introduce the notion of right (resp. contact) complete unfoldings, which can be used to give an alternative definition of right (resp. contact) modality. We use étale neighbourhoods in order to show that an algebraic representation of the semiuniversal deformation is complete, see Proposition 2.14. The results of this section are used for the classification in Section 3.

2.1. G -modality. We use a Rosenlicht stratification of a variety¹ X under the action of an algebraic group G to define modality. By Rosenlicht [Ros56, Thm.2] (see also [Ros63]) there exists an open dense subset $X_1 \subset X$, which is invariant under G s.t. X_1/G is a geometric quotient (cf. [MFK82, §1]), in particular, the orbit space X_1/G is an algebraic variety and the projection $p_1: X_1 \rightarrow X_1/G$, $x \mapsto [x]$, is a surjective morphism.

¹By an algebraic variety we mean a separated scheme of finite type over an algebraically closed field K , see [Har77], which is fixed through this paper. By a point we mean a closed point.

Since $X \setminus X_1$ is a variety of lower dimension, which is invariant under G , we can apply the theorem of Rosenlicht to $X \setminus X_1$ and get an invariant open dense subset $X_2 \subset X \setminus X_1$ s.t. X_2/G is a geometric quotient. Continuing in this way with $X_3 \subset (X \setminus X_1) \setminus X_2$, we can finally write X as finite disjoint union of G -invariant locally closed algebraic subvarieties $X_i, i = 1, \dots, s$, such that X_i/G is a geometric quotient with quotient morphism $p_i: X_i \rightarrow X_i/G$. We call $\{X_i, i = 1, \dots, s\}$ a *Rosenlicht stratification of X under G* . Note that a Rosenlicht stratification is by no means unique and that the proof of Rosenlicht, which works for arbitrary G , is not constructive.

Definition 2.1. Let $\{X_i, i = 1, \dots, s\}$ be a Rosenlicht stratification of the algebraic variety X under the action of an algebraic group G with quotient morphisms $p_i: X_i \rightarrow X_i/G$, and let U be an open neighbourhood of $x \in X$. We define

$$G\text{-mod}(U) := \max_{1 \leq i \leq s} \{\dim(p_i(U \cap X_i))\},$$

and call

$$G\text{-mod}(x) := \min\{G\text{-mod}(U) \mid U \text{ a neighbourhood of } x\}$$

the *G-modality* of x (in X).

Note that here and later, wherever we write $\dim S$, the set S is *constructible*, i.e. it is a finite union of locally closed subsets of a variety, so that $\dim S$ is defined. By Corollary A.3 $G\text{-mod}(U)$ and $G\text{-mod}(x)$ are independent of the Rosenlicht stratification.

Remark 2.2. Let $\{X_i\}$ be a Rosenlicht stratification of X under G . In [Wal83] Wall introduced the *r-modal set* $M_r(X)$ to be the closure of the union $\cup\{X_i \mid i \in I_r\}$ with $I_r := \{i \mid \dim X_i/G \geq r\}$, which satisfies

$$X = M_0 \supset M_1 \supset \dots \supset M_{\dim X}.$$

Using our definition of $G\text{-mod}(x)$ one can show that

$$M_r(X) = \{x \in X \mid G\text{-mod}(x) \geq r\}.$$

Moreover, we have $G\text{-mod}(x) = r \Leftrightarrow x \in M_r(X) \setminus M_{r+1}(X)$ for $x \in X$, and for any open subset $U \subset X$, $G\text{-mod}(U) = \max\{G\text{-mod}(x) \mid x \in U\}$.

Now we will give the definition of modality for isolated singularities. We recall two important invariants of singularities. For $f \in K[[\mathbf{x}]]$, let $\mu(f) := \dim K[[\mathbf{x}]]/j(f)$ be the *Milnor number* and $\tau(f) := \dim K[[\mathbf{x}]]/(\langle f \rangle + j(f))$ the *Tjurina number* of f , where $j(f)$ is the jacobian ideal, i.e. the ideal in $K[[\mathbf{x}]]$ generated by all partials of f . Note that f (resp. $K[[\mathbf{x}]]/\langle f \rangle$) has an *isolated singularity* if $\mu(f) < \infty$ (resp. $\tau(f) < \infty$).

Let $f \in \mathfrak{m} \subset K[[\mathbf{x}]]$, with $\mathfrak{m} = \langle \mathbf{x} \rangle$ the maximal ideal, be such that $\mu(f) < \infty$ (resp. $\tau(f) < \infty$) and let $G = \mathcal{R}$ be right group (resp. $G = \mathcal{K}$ be the contact group). In order to define the modality of a power series f by using algebraic groups, we have to consider its k -jet

$$j^k f := j^k(f) := \text{image of } f \text{ in } J_k := K[[\mathbf{x}]]/\mathfrak{m}^{k+1}$$

for sufficiently large k . Let $G_k = \mathcal{R}_k$ resp. $G_k = \mathcal{K}_k$ the k -jet of G , see Section A.2.

Definition 2.3. We define the G -modality of f , $G\text{-mod}(f)$, to be the G_k -modality of $j^k(f)$ in J_k (in the sense of Definition 2.1) for sufficiently large integer k . We call f *right* (resp. *contact*) *simple*, *unimodal*, *bimodal* and *r-modal* if $\mathcal{R}\text{-mod}(f)$ (resp. $\mathcal{K}\text{-mod}(f)$) equals to 0, 1, 2 and r respectively.

Here, an integer k is *sufficiently large* for f w.r.t. G if there exists a neighbourhood U of $j^k(f)$ in J_k s.t. every $g \in \mathfrak{m}$ with $j^k g \in U$ is k -determined w.r.t. G . This means that each $h \in \mathfrak{m}$ s.t. $j^k(h) = j^k(g)$, is G -equivalent to g . Combining Propositions 2.6 and 2.12 below we obtain that $G\text{-mod}(f)$ is independent of the sufficiently large k . The existence of a sufficiently large integer k for f w.r.t G will be shown by the following proposition.

Proposition 2.4. *Let $f \in \mathfrak{m}^2$ be such that $\mu(f) < \infty$ (resp. $\tau(f) < \infty$) and let $G = \mathcal{R}$ (resp. $G = \mathcal{K}$). Then every $k \geq 2 \cdot \mu(f)$ (resp. $k \geq 2 \cdot \tau(f)$) is sufficiently large for f w.r.t. \mathcal{R} (resp. w.r.t. \mathcal{K}). For $f \in \mathfrak{m} \setminus \mathfrak{m}^2$, $k = 1$ is sufficiently large for f w.r.t. G .*

Proof. By the upper semi-continuity of μ, τ (Lemma A.13), the subsets

$$U_\mu := \{g \in K[[\mathbf{x}]] \mid \mu(g) \leq \mu(f)\} \text{ and } U_\tau := \{g \in K[[\mathbf{x}]] \mid \tau(g) \leq \tau(f)\}$$

are open² in $K[[\mathbf{x}]]$. It follows from [BGM12, Cor. 1] that g is k -determined w.r.t. G for all $g \in U_\mu$ (resp. U_τ) and all $k \geq 2 \cdot \mu(f)$ (resp. $k \geq 2 \cdot \tau(f)$). This means that k is sufficiently large for f w.r.t. G . For $f \in \mathfrak{m} \setminus \mathfrak{m}^2$, f is non-singular and the result follows from the implicit function theorem. \square

2.2. Complete unfoldings. Let T be an (affine) variety over K with structure sheaf \mathcal{O}_T and its algebra of global section $\mathcal{O}(T) = \mathcal{O}_T(T)$. If $F = \sum a_\alpha \mathbf{x}^\alpha \in \mathcal{O}(T)[[\mathbf{x}]]$, $a_\alpha \in \mathcal{O}(T)$, then for each $t \in T$, $a_\alpha(t) \in K$ denotes the image of a_α in $\mathcal{O}_{T,t}/\mathfrak{m}_t = K$, with \mathfrak{m}_t the maximal ideal of the stalk $\mathcal{O}_{T,t}$. Therefore

$$f_t(\mathbf{x}) := F(\mathbf{x}, t) = \sum a_\alpha(t) \mathbf{x}^\alpha \in K[[\mathbf{x}]], t \in T,$$

defines a family of power series f_t parametrized by $t \in T$. In the following we often write $f_t(\mathbf{x})$ or $F(\mathbf{x}, t)$ instead of f_t or F , just to show the variables \mathbf{x} and the parameter $t \in T$. Moreover if $T = \mathbb{A}^l = \text{Spec}(K[t_1, \dots, t_l])$ then $F \in K[\mathbf{t}][[\mathbf{x}]] \subset K[[\mathbf{x}, \mathbf{t}]]$ with $\mathbf{t} := (t_1, \dots, t_l)$. Let $f \in \mathfrak{m}$ and let $t_0 \in T$. An element $F(\mathbf{x}, t) \in \mathcal{O}(T)[[\mathbf{x}]]$ is called an *unfolding* or *deformation with trivial section* of f at $t_0 \in T$ over T if $f_{t_0} = f$ and $f_t \in \mathfrak{m}$ for all $t \in T$.

Definition 2.5. Let $f \in \mathfrak{m}$ be such that $\mu(f) < \infty$ (resp. $\tau(f) < \infty$), and let $f_t(\mathbf{x})$ be an unfolding of f at t_0 over an affine variety T . Let $G = \mathcal{R}$ (resp. $G = \mathcal{K}$), let k be sufficiently large for f w.r.t. G and let Φ_k be the morphism of algebraic varieties from T to the k -jet space J_k defined by

$$\Phi_k : T \rightarrow J_k, t \mapsto j^k f_t(\mathbf{x}).$$

We define $G\text{-mod}_F(f) := G_k\text{-mod}_{\Phi_k}(t_0)$ and call it the G -modality of f w.r.t. the unfolding F . Note that G_k acts on J_k and that $G_k\text{-mod}_{\Phi_k}(t_0)$ is understood in the sense of Definition A.5.

² $V \subset K[[\mathbf{x}]]$ is open iff $j^l(V)$ is open in J_l for all l .

Proposition 2.6. *For any unfolding F of f at t_0 , the number $G\text{-mod}_F(f)$ is independent of the sufficient large integer k for f w.r.t. G .*

Proof. Let U_G denote the open neighbourhood U_μ resp. U_τ of f in $K[[\mathbf{x}]]$, defined as in the proof of Proposition 2.4. It is easy to see that the map

$$\Phi : T \longrightarrow K[[\mathbf{x}]], \quad t \mapsto f_t(\mathbf{x})$$

is continuous. Then the pre-image $V_G = \Phi^{-1}(U_G)$ is an open neighbourhood of t_0 . For each k sufficient large for f w.r.t. G we consider the map

$$\varphi_k : V_G \xrightarrow{i} T \xrightarrow{\Phi} K[[\mathbf{x}]] \xrightarrow{j^k} J_k.$$

By Corollary A.10,

$$G\text{-mod}_{\Phi_k}(t_0) = G\text{-mod}_{\varphi_k}(t_0)$$

since $\Phi_k = j^k \circ \Phi$. If k_1, k_2 are both sufficient large for f w.r.t. G , then we can easily check that

$$\varphi_{k_1}^{-1}(G_{k_1} \cdot \varphi_{k_1}(t)) = \varphi_{k_2}^{-1}(G_{k_2} \cdot \varphi_{k_2}(t)), \quad \forall t \in V_G.$$

Corollary A.11 yields that

$$G_{k_1}\text{-mod}_{\varphi_{k_1}}(t_0) = G_{k_2}\text{-mod}_{\varphi_{k_2}}(t_0)$$

which proves the proposition. \square

Proposition 2.7 (Semicontinuity of modality). *Let $G = \mathcal{R}$ (resp. $G = \mathcal{K}$). Then the G -modality is upper semicontinuous, i.e. for all $i \in \mathbb{N}$, the sets*

$$U_i := \{f \in \mathfrak{m} \subset K[[\mathbf{x}]] \mid G\text{-mod}(f) \leq i\}$$

are open in $K[[\mathbf{x}]]$. Consequently, the G -modality is upper semicontinuous for unfoldings, i.e. for any unfolding $f_t(\mathbf{x})$ at t_0 over T of f with $\mu(f) < \infty$ (resp. $\tau(f) < \infty$) the set

$$\{t \in T \mid G\text{-mod}(f_t) \leq G\text{-mod}(f)\}$$

is open in T .

Proof. Let $f \in U_i$, and let $k \geq 2\mu(f)$ (resp. $k \geq 2\tau(f)$). By Proposition 2.4 k is sufficiently large for f w.r.t. G and hence $G\text{-mod}(f) = G_k\text{-mod}(j^k f)$. Take an open neighbourhood V of $j^k f$ in J_k such that $G_k\text{-mod}(V) = G_k\text{-mod}(j^k f)$. Set $U := (j^k)^{-1}(V) \cap \tilde{U}$, where

$$\tilde{U} := \{g \in K[[\mathbf{x}]] \mid \mu(g) \leq \mu(f)\}.$$

By Lemma A.13 \tilde{U} is open and hence U is an open neighbourhood of f in $K[[\mathbf{x}]]$. We now show that $U \subset U_i$. In fact, for any $g \in U$ one has $k \geq 2\mu(f) \geq 2\mu(g)$ and hence k is sufficiently large for g due to Proposition 2.4. Then

$$G\text{-mod}(g) = G_k\text{-mod}(j^k g) \leq G_k\text{-mod}(V) = G_k\text{-mod}(j^k f) = G\text{-mod}(f) \leq i.$$

This implies that $U \subset U_i$ and hence U_i is open in $K[[\mathbf{x}]]$. \square

So far we considered families of singularities parametrized by (affine) varieties, in particular by sufficiently high jet spaces. Now we want to use the semiuniversal deformation (with trivial section) of a singularity since its base space has much smaller dimension. However for moduli problems, the formal deformation theory is not sufficient. We have to pass to the étale topology and apply Artin's resp. Elkik's algebraization theorems.

Recall that an étale neighbourhood of a point s in a variety S consists of a variety U with a point $u \in U$ and an étale morphism $\varphi : U \rightarrow S$ with $\varphi(u) = s$ (see, [Mum88, Definition III.5.1]). φ is a morphism of pointed varieties, usually denoted by $\varphi : U, u \rightarrow S, s$. The connected étale neighbourhoods of s form a filtered system and the direct limit

$$\tilde{\mathcal{O}}_{S,s} := \varinjlim \mathcal{O}_{U,u} = \varinjlim \mathcal{O}(U)$$

is called the *Henselization* (see [Na53], [Ra70], [KPPRM78]) of $\mathcal{O}_{S,s}$. We have $\hat{\mathcal{O}}_{S,s} = \hat{\tilde{\mathcal{O}}}_{S,s} = \hat{\mathcal{O}}_{U,u}$ where $\hat{}$ denotes the completion w.r.t. the maximal ideal. The Henselization of $K[\mathbf{x}]_{\langle \mathbf{x} \rangle}$ is the ring $K\langle \mathbf{x} \rangle$ of algebraic power series in $\mathbf{x} = (x_1, \dots, x_n)$. $K[\mathbf{x}] \subset K\langle \mathbf{x} \rangle \subset K[[\mathbf{x}]]$ and $K\langle \mathbf{x} \rangle$ may be considered as the union of all $\mathcal{O}(U) \subset K[[\mathbf{x}]]$ or $\mathcal{O}_{U,u} \subset K[[\mathbf{x}]]$ for U, u an étale neighbourhood of 0 in \mathbb{A}^n . This implies

Remark 2.8. For each finitely generated subalgebra $A \subset K\langle \mathbf{x} \rangle$ containing $K[\mathbf{x}]$ there exist an étale neighbourhood $\varphi : V, v_0 \rightarrow \mathbb{A}^n, 0$ of $0 \in \mathbb{A}^n$ and an inclusion $A \hookrightarrow \mathcal{O}(V)$, $a(\mathbf{x}) \mapsto a_\varphi \in \mathcal{O}(V)$. If $a \in K[\mathbf{x}]$ then $a_\varphi(v) = a(\varphi(v))$ for all $v \in V$, where $a_\varphi(v) \in K$ is the image of a_φ in $\mathcal{O}_{V,v}/\mathfrak{m}_v$.

Definition 2.9. Let $F(\mathbf{x}, t)$ be an unfolding of f at t_0 over an affine variety T . Let $G = \mathcal{R}$ or $G = \mathcal{K}$.

- (a) An unfolding $H(\mathbf{x}, s)$ of f at s_0 over an affine variety S is called a *pullback* or an *induced unfolding* of F if there exists a morphism $\psi : S, s_0 \rightarrow T, t_0$ s.t. $H(\mathbf{x}, s) = F(\mathbf{x}, \psi(s))$.
- (b) An unfolding $H(\mathbf{x}, s)$ of f at s_0 over an affine variety S is called an *étale G -pullback* of F if there exist an étale neighbourhood $\varphi : W, w_0 \rightarrow S, s_0$ and a morphism $\psi : W, w_0 \rightarrow T, t_0$ such that $H(\mathbf{x}, \varphi(w))$ is G -equivalent to $F(\mathbf{x}, \psi(w))$ for all $w \in W$.
- (c) The unfolding $F(\mathbf{x}, t)$ is called *right* (resp. *contact*) *complete* (or *G -complete*) if any unfolding of f is an étale G -pullback of F .

The following lemma is an immediate consequence of the definition.

Lemma 2.10. *Let H be an étale G -pullback of the unfolding F . If H is G -complete, then so is F .*

Proposition 2.11. *Any singularity f with $\mu(f) < \infty$ (resp. $\tau(f) < \infty$) has a right (resp. contact) complete unfolding given by a sufficiently large jet space. More precisely, if k is sufficiently large for f w.r.t. \mathcal{R} (resp. w.r.t. \mathcal{K}), then the unfolding of f at 0 over $J_k = \mathbb{A}^N$ (with the identification: $\sum_{|\alpha| \leq k} c_\alpha \mathbf{x}^\alpha \mapsto c = (c_\alpha)_{|\alpha| \leq k}$, $\mathcal{O}(J_k) = K[(c_\alpha)_{|\alpha| \leq k}]$)*

$$f_c(\mathbf{x}) = F(\mathbf{x}, c) = f(\mathbf{x}) + \sum_{|\alpha| \leq k} c_\alpha \mathbf{x}^\alpha \in K[(c_\alpha)_{|\alpha| \leq k}][[\mathbf{x}]]$$

is right (resp. contact) complete.

Proof. Since k is sufficiently large for f w.r.t. right (resp. contact) equivalence, there exists a neighbourhood $U \subset J_k$ of $j^k f$ such that each $g \in U$ is right (resp. contact) k -determined. Let $h_s(\mathbf{x}) := H(\mathbf{x}, s)$ be an arbitrary unfolding of f at s_0 over S and let $W := \psi^{-1}(U)$ be the pre-image of U by the morphism

$$\psi : S \longrightarrow \mathbb{A}^N, s \mapsto j^k h_s(\mathbf{x}) - j^k f(\mathbf{x}).$$

Then $H(\mathbf{x}, s)$ is right (resp. contact) equivalent to $F(\mathbf{x}, \psi(s))$ for all $s \in W$ since

$$j^k H(\mathbf{x}, s) = j^k F(\mathbf{x}, \psi(s)) \in U$$

and hence H is a pullback of F . Since every pullback is an étale G -pullback, this proves the proposition. \square

Proposition 2.12. *Let $f \in K[[\mathbf{x}]]$ be such that $\mu(f) < \infty$ (resp. $\tau(f) < \infty$) and let $G = \mathcal{R}$ (resp. $G = \mathcal{K}$). Let $F(\mathbf{x}, t)$ be an unfolding of f at t_0 over T .*

- (i) *If the unfolding $H(\mathbf{x}, s)$ of f at s_0 over S is an étale G -pullback of F , then $G\text{-mod}_H(f) \leq G\text{-mod}_F(f)$.*
- (ii) *We always have $G\text{-mod}(f) \geq G\text{-mod}_F(f)$. Equality holds if $F(\mathbf{x}, t)$ is G -complete.*

Proof. (i) Since H is an étale G -pullback of F , there exist an étale neighbourhood $\varphi : W, w_0 \rightarrow S, s_0$ and a morphism $\psi : W, w_0 \rightarrow T, t_0$ such that $G \cdot H(\mathbf{x}, \varphi(w)) = G \cdot F(\mathbf{x}, \psi(w))$ for all $w \in W$.

Let k be sufficiently large for f and let Φ_k and Ψ_k be the morphisms defined by

$$\Phi_k : T \rightarrow J_k, t \mapsto j^k f_t(\mathbf{x}) \text{ and } \Psi_k : S \rightarrow J_k, s \mapsto j^k h_s(\mathbf{x}).$$

Then $\Phi^{-1}(G \cdot \Phi(w)) = \Psi^{-1}(G \cdot \Psi(w))$ for all $w \in W$ with $\Phi := \Phi_k \circ \psi$ and $\Psi := \Psi_k \circ \varphi$. Combining Corollary A.10 and A.11 we obtain

$$G\text{-mod}_F(f) = G_k\text{-mod}_{\Phi_k}(t_0) \geq G_k\text{-mod}_{\Phi}(w_0) = G_k\text{-mod}_{\Psi}(w_0) = G_k\text{-mod}_{\Psi_k}(s_0) = G\text{-mod}_H(f).$$

- (ii) Since k is sufficiently large for f w.r.t. G ,

$$G\text{-mod}(f) = G_k\text{-mod}(j^k f) \text{ and } G\text{-mod}_F(f) = G_k\text{-mod}_{\Phi_k}(t_0).$$

It follows from Corollary A.9 that $G_k\text{-mod}(j^k f) \geq G_k\text{-mod}_{\Phi_k}(t_0)$. Hence

$$G\text{-mod}(f) \geq G\text{-mod}_F(f).$$

We identify J_k with \mathbb{A}^N via the map $\sum_{|\alpha| \leq k} c_\alpha \mathbf{x}^\alpha \mapsto c := (c_\alpha)_{|\alpha| \leq k}$ and consider the unfolding

$$h_c(\mathbf{x}) := H(\mathbf{x}, c) := f(\mathbf{x}) + \sum_{|\alpha| \leq k} c_\alpha \mathbf{x}^\alpha$$

of f at 0 over \mathbb{A}^N . Since the translation

$$\begin{aligned} \psi : \mathbb{A}^N &\longrightarrow J_k &= \mathbb{A}^N \\ c &\mapsto j^k h_c(\mathbf{x}) &= j^k f + c \end{aligned}$$

is an isomorphism, it follows from Corollary A.9 that

$$G_k\text{-mod}(j^k f) = G_k\text{-mod}_\psi(0)$$

and hence $G\text{-mod}(f) = G\text{-mod}_H(f)$.

Now, if the unfolding F is G -complete, then H is an étale G -pullback of F . By (i), $G\text{-mod}_H(f) \leq G\text{-mod}_F(f)$ and hence $G\text{-mod}(f) = G\text{-mod}_F(f)$. \square

Example 2.13. (a) The unfolding $F(x, t) = x^{p+1} + t_1x + \dots + t_px^p$ of the univariate polynomial $f = x^{p+1}$ over $T = \mathbb{A}^p$ is right complete. Here (t_1, \dots, t_p) are the coordinates of $t \in T$.

Indeed, for any unfolding $H(x, s)$ of f at s_0 over some variety S we write

$$H(x, s) = a_1(s)x + \dots + a_p(s)x^p + a_{p+1}(s)x^{p+1} + \dots$$

with $a_i(s) \in \mathcal{O}(S)$. Then $a_i(s_0) = 0 \ \forall i \leq p$ and $a_{p+1}(s_0) \neq 0$. Consider the morphism $\varphi : S \rightarrow T$, $s \mapsto (a_1(s), \dots, a_p(s))$ and the open neighbourhood $W := \{s \in S \mid a_{p+1}(s) \neq 0\}$ of s_0 in S . Then it follows from [BGM12, Thm. 2.1] that

$$F(x, \varphi(s)) \sim_r H(x, s), \text{ for each } s \in W.$$

Note that $\{x, \dots, x^p\}$ is a basis of $\mathfrak{m}/\mathfrak{m}j(f)$ and that F is a semiuniversal deformation of f with trivial section by Proposition 2.14.

(b) The right semi-universal deformation (without section) of $f = x^{p+1} \in K[[x]]$ with $\text{char}(K) = p > 2$ is given by $H(x, t) = x^{p+1} + t_1x + \dots + t_{p-1}x^{p-1}$. This unfolding of f over \mathbb{A}^{p-1} is not right complete. In fact, it is not difficult to see that $H(x, t)$ is equivalent to one of $\{x, \dots, x^{p-1}, x^{p+1}\}$ for $t \in \mathbb{A}^p$. Corollary 2.17 yields that $\mathcal{R}\text{-mod}_H(f) = 0$, while $\mathcal{R}\text{-mod}(f) > 0$ by Theorem 3.1 and hence H is not right complete due to Proposition 2.7.

To see this directly, consider the family $x^{p+1} + sx^p$ in characteristic $p > 0$ over \mathbb{A}^1 which, as an unfolding with trivial section, cannot be induced by a morphism $\varphi : \mathbb{A}^1 \rightarrow \mathbb{A}^{p-1}$: Since $H(x, \varphi(s))$ has multiplicity $\neq p$ in $K[[x]]$ for all $s \neq 0$, it cannot be right equivalent to $x^{p+1} + sx^p$ which has multiplicity p for $s \neq 0$.

This is of course not a contradiction to F being versal as deformation without section, which means that the family $x^{p+1} + sx^p \in K[[x, s]]$ can be induced from H (up to isomorphism in $K[[x, s]]$ over $K[[s]]$) by a morphism $\varphi : K[[t_1, \dots, t_{p-1}]] \rightarrow K[[s]]$. In fact, define φ by $t_1 \mapsto -s^p, t_i \mapsto 0$ for $i > 1$, then, if $\text{char}(K) = p$,

$$H(x, \varphi(s)) = -s^p x + x^{p+1} = (x - s)^p x \sim_r x^{p+1} + sx^p,$$

via the isomorphism $\Phi : K[[x, s]] \rightarrow K[[x, s]]$ over $K[[s]]$, given by $x \mapsto x - s, s \mapsto s$. However, Φ does not respect the trivial section.

If $\text{char}(K) \neq p$, we can use the Tschirnhaus transformation $x \mapsto x - \frac{s}{p}$ to eliminate sx^p from $x^{p+1} + sx^p$ and to show that $x^{p+1} + sx^p$ can be induced from H .

The following proposition is stronger than Proposition 2.11 because it reduces the number of parameters of a G -complete unfolding considerably. For the proof we need the nested Artin approximation theorem.

Proposition 2.14. *Let $f \in \mathfrak{m} \subset K[[\mathbf{x}]]$ be such that $\mu(f) < \infty$ (resp. $\tau(f) < \infty$). Let $g_1, \dots, g_l \in K[\mathbf{x}]$ be a system of K -generators of $\mathfrak{m}/\mathfrak{m} \cdot j(f)$ (resp. of $\mathfrak{m}/\langle f \rangle + \mathfrak{m} \cdot j(f)$), with $j(f)$ the jacobian ideal of f . Then the unfolding (with trivial section) of f over \mathbb{A}^l ,*

$$F(\mathbf{x}, t) = f(\mathbf{x}) + \sum_{i=1}^l t_i g_i(\mathbf{x}) \in K[\mathbf{t}][[\mathbf{x}]]$$

is (an algebraic representative of) a formally versal deformation of f with trivial section with respect to right (resp. contact) equivalence, which is semi-universal if the system $\{g_i\}$ is a basis. This unfolding is right (resp. contact) complete.

Proof. We first show that if $\{g_i\}$ is a basis, then F represents a formally semiuniversal deformation of f with trivial section with respect to right (resp. contact) equivalence. Indeed we may consider F as a deformation of f over $K[[\mathbf{t}]] := K[[t_1, \dots, t_l]]$. It is shown in [BGM12, Prop. 2.7] that the tangent space to the base space of the semiuniversal deformation with trivial section is $\mathfrak{m}/\mathfrak{m} \cdot j(f)$ (resp. $\mathfrak{m}/\langle f \rangle + \mathfrak{m} \cdot j(f)$). The proof of the existence of a semiuniversal deformation in [KaS72] or [GLS06, Thm. II.1.16] can be easily adapted to deformations with section, showing the versality of F and hence proving the first claim.

Let $G = \mathcal{R}$ (resp. $G = \mathcal{K}$) and let k be sufficiently large for f w.r.t. G . Let j^k denote the projections $K[\mathbf{t}][[\mathbf{x}]] \rightarrow K[\mathbf{t}][[\mathbf{x}]]/\langle \mathbf{x} \rangle^{k+1}$ and $K[[\mathbf{x}]] \rightarrow K[[\mathbf{x}]]/\langle \mathbf{x} \rangle^{k+1}$. Then we may replace F resp. f by $j^k F$ resp. $j^k f$ and assume that $F = j^k F \in K[\mathbf{x}, \mathbf{t}]$ and $f = j^k f \in K[\mathbf{x}]$ by the following facts:

- (a) $\{j^k g_1, \dots, j^k g_l\}$ is a system of generators of $\mathfrak{m}/\mathfrak{m} \cdot j(j^k f)$ (resp. of $\mathfrak{m}/\langle f \rangle + \mathfrak{m} \cdot j(j^k f)$) with $j(j^k f)$ the jacobian ideal of $j^k f$.
- (b) F is G -complete if and only if $j^k F(\mathbf{x}, t)$ is a G -complete unfolding of $j^k f(\mathbf{x})$ at t_0 over T (see [Ng13, Prop. 3.4.12]).

Consider the complete unfolding of f over $\mathbb{A}^N = J_k$ at 0

$$h_c(\mathbf{x}) := H(\mathbf{x}, c) = f(\mathbf{x}) + \sum_{|\alpha| \leq k} c_\alpha \mathbf{x}^\alpha$$

as in Proposition 2.11. By Lemma 2.10, the proof is completed by showing that H is an étale G -pullback of F .

In fact, consider H as a deformation with trivial section of f over $K[[\mathbf{c}]] := K[[c_\alpha]_{|\alpha| \leq k}]$. Although F and H are polynomials, the versality of F implies only that H can be induced formally from F . This means, there exist tuples of formal power series $\hat{\Phi} = (\hat{\Phi}_1, \dots, \hat{\Phi}_n) \in (\langle \mathbf{x} \rangle K[[\mathbf{x}, \mathbf{c}]])^n$, $\hat{\varphi} = (\hat{\varphi}_1, \dots, \hat{\varphi}_l) \in (\langle \mathbf{c} \rangle K[[\mathbf{c}]])^l$ and a unit $\hat{u}(\mathbf{x}, \mathbf{c}) \in K[[\mathbf{x}, \mathbf{c}]]$ (with $\hat{u} = 1$ for $G = \mathcal{R}$) with

$$(2.1) \quad \hat{u}(0, 0) \neq 0 \text{ and } \left(\det \left(\frac{\partial \hat{\Phi}_i}{\partial x_j} \right) \right)_{(0,0)} \neq 0,$$

such that

$$H(\hat{\Phi}(\mathbf{x}, \mathbf{c}), \mathbf{c}) = \hat{u}(\mathbf{x}, \mathbf{c}) \cdot F(\mathbf{x}, \hat{\varphi}(\mathbf{c})).$$

Let $\mathbf{y} = (y_1, \dots, y_{n+l+1})$ be new indeterminates (omitting y_{n+l+1} if $G = \mathcal{R}$) and let

$$P(\mathbf{x}, \mathbf{c}, \mathbf{y}) = H(y_1, \dots, y_n, \mathbf{c}) - y_{n+l+1} \cdot F(\mathbf{x}, y_{n+1}, \dots, y_{n+l}) \in K[\mathbf{x}, \mathbf{c}, \mathbf{y}].$$

The formal versality of F implies that $P = 0$ has a formal solution $\hat{\mathbf{y}} = (\hat{y}_1, \dots, \hat{y}_{n+l+1})$ with

$$\begin{aligned} \hat{y}_i &:= \hat{\Phi}_i \in K[[\mathbf{x}, \mathbf{c}]], 1 \leq i \leq n, \\ \hat{y}_{n+j} &:= \hat{\varphi}_j \in K[[\mathbf{c}]], 1 \leq j \leq l, \\ \hat{y}_{n+l+1} &:= \hat{u} \in K[[\mathbf{x}, \mathbf{c}]]. \end{aligned}$$

By the nested Artin Approximation Theorem ([Po86, Theorem 1.4]), there exists an algebraic solution $\tilde{\mathbf{y}} = (\tilde{y}_1, \dots, \tilde{y}_{n+l+1})$ of $P = 0$ such that

$$\begin{aligned} \tilde{y}_i &\in K\langle \mathbf{x}, \mathbf{c} \rangle, 1 \leq i \leq n, \\ \tilde{y}_{n+j} &\in K\langle \mathbf{c} \rangle, 1 \leq j \leq l, \\ \tilde{y}_{n+l+1} &\in K\langle \mathbf{x}, \mathbf{c} \rangle \end{aligned}$$

and

$$(2.2) \quad \tilde{\mathbf{y}}(\mathbf{c}, \mathbf{x}) - \hat{\mathbf{y}}(\mathbf{c}, \mathbf{x}) \in \langle \mathbf{c}, \mathbf{x} \rangle^2.$$

Passing to the k -jet spaces by the projection $j^k : K[[\mathbf{x}, \mathbf{c}]] \rightarrow K[[\mathbf{x}, \mathbf{c}]]/\langle \mathbf{x} \rangle^{k+1}$ we get

$$j^k(P(\mathbf{x}, \mathbf{c}, j^k(\tilde{\mathbf{y}}))) = j^k(H(j^k(\tilde{y}_1), \dots, j^k(\tilde{y}_n), \mathbf{c})) - j^k(\tilde{y}_{n+l+1}) \cdot j^k(F(\mathbf{x}, j^k(\tilde{y}_{n+1}), \dots, j^k(\tilde{y}_{n+l})))) = 0.$$

Let $A \subset K[[\mathbf{c}]]$ be the subalgebra generated by $c_\alpha, |\alpha| \leq k$, and all the coefficients of $\mathbf{x}^\alpha, |\alpha| \leq k$, which appear in all $j^k(\tilde{y}_i)(\mathbf{x}, \mathbf{c}), i = 1, \dots, n + l + 1$. Then $A \subset K\langle \mathbf{c} \rangle$ since $K\langle \mathbf{c}, \mathbf{x} \rangle \subset K\langle \mathbf{c} \rangle[[\mathbf{x}]]$ (cf. [Ng13, Lemma 3.4.9]). It follows from Remark 2.8 that there exists an étale neighbourhood $\varphi : V, v_0 \rightarrow \mathbb{A}^N, 0$ and an inclusion $A \hookrightarrow \mathcal{O}(V), a \mapsto a_\varphi$ such that if $a \in K[\mathbf{c}]$ the $a_\varphi(v) = a(\varphi(v))$ for all $v \in V$. This implies that, denoting by $\iota : A[\mathbf{x}] \hookrightarrow \mathcal{O}(V)[\mathbf{x}]$ the induced inclusion,

$$j^k(P(\mathbf{x}, \varphi(v), \Phi(\mathbf{x}, v))) = 0 \text{ in } \mathcal{O}(V)[\mathbf{x}],$$

where $\Phi(\mathbf{x}, v) = (\Phi_1(\mathbf{x}, v), \dots, \Phi_n(\mathbf{x}, v))$ with $\Phi_i((\mathbf{x}, v)) = \iota(j^k(\tilde{y}_i)(\mathbf{x}, \mathbf{c}))$. Combining (2.1), (2.2) we obtain that there exists an open neighbourhood $W' \subset V$ of v_0 such that for all $v \in W'$,

$$\Phi(\mathbf{x}, v) \in \text{Aut}(K[[\mathbf{x}]]) \text{ and } u(\mathbf{x}, v) := \iota(j^k \tilde{y}_{n+l+1}(\mathbf{x}, \mathbf{c})) \in K[[\mathbf{x}]]^*.$$

This implies $j^k H(\Phi(\mathbf{x}, v), \varphi(v))$ is G -equivalent to $F(\mathbf{x}, \psi(v))$, where

$$\psi : V, v_0 \rightarrow \mathbb{A}^l, \psi(v) = (\psi_1(v), \dots, \psi_l(v))$$

with $\psi_j(v) = \iota(j^k(\tilde{y}_{n+j})(\mathbf{c}))$. Moreover since k is sufficiently large for f w.r.t. G , there exists an open neighbourhood $W \subset W'$ of v_0 such that

$$H(\Phi(\mathbf{x}, v), \varphi(v)) \sim_G j^k H(\Phi(\mathbf{x}, v), \varphi(v)), \forall v \in W.$$

Hence

$$H(\Phi(\mathbf{x}, v), \varphi(v)) \sim_G j^k H(\Phi(\mathbf{x}, v), \varphi(v)) \sim_G F(\mathbf{x}, \psi(v)), \forall u \in W,$$

which proves the claim. \square

The following proposition will be used to show how the modality of f is related to the number of parameters in families of normal forms. To make this precise we introduce G -modular families. By a G -modular family over a variety S we mean a family $h_s(\mathbf{x}) = H(\mathbf{x}, s) \in \mathcal{O}(S)[[\mathbf{x}]]$ such that for each $s \in S$ there are only finitely many $s' \in S$ such that h_s is G -equivalent to $h_{s'}$.

Proposition 2.15. *Let $f \in K[[\mathbf{x}]]$ be such that $\mu(f) < \infty$ (resp. $\tau(f) < \infty$) and let $G = \mathcal{R}$ (resp. $G = \mathcal{K}$). Let $f_t(\mathbf{x}) = F(\mathbf{x}, t)$ be any unfolding of f at t_0 over T . Assume that there exist an open neighbourhood $W \subset T$ of t_0 and families $h_{s_i}^{(i)}(\mathbf{x}) = H^{(i)}(\mathbf{x}, s_i) \in \mathcal{O}(S_i)[[\mathbf{x}]]$ over affine varieties $S_i, i = 1, \dots, q$, such that for all $t \in W$ there exists $s_i \in S_i$ such that $f_t \sim_G h_{s_i}^{(i)}$.*

(i) *We have*

$$G\text{-mod}_F(f) \leq \max_{i=1, \dots, q} \{\dim S_i\}.$$

(ii) *Assume moreover that each family $h_{s_i}^{(i)}(\mathbf{x}), i = 1, \dots, q$, is G -modular and that for each open neighbourhood V of t_0 in W and for all $s_i \in S_i$ there exists a $t \in V$ such that $f_t \sim_G h_{s_i}^{(i)}$. Then*

$$G\text{-mod}_F(f) = \max_{i=1, \dots, q} \{\dim S_i\}.$$

Proof. (i) Let k be sufficiently large for f w.r.t. G . Considering the morphisms

$$\Phi_k : T \rightarrow J_k, t \mapsto j^k f_t(\mathbf{x}) \text{ and } \Phi_k^{(i)} : S_i \rightarrow J_k, s \mapsto j^k h_s^{(i)}(\mathbf{x}), i = 1, \dots, q,$$

and applying Corollary A.12(i) we obtain that

$$G_k\text{-mod}_{\Phi_k}(W) \leq \max_{i=1, \dots, q} \{G_k\text{-mod}_{\Phi_k^{(i)}}(S_i)\}.$$

Hence

$$G\text{-mod}_F(f) = G_k\text{-mod}_{\Phi_k}(0) \leq G_k\text{-mod}_{\Phi_k}(W) \leq \max_{i=1, \dots, q} \{G_k\text{-mod}_{\Phi_k^{(i)}}(S_i)\} \leq \max_{i=1, \dots, q} \{\dim S_i\}.$$

(ii) Let V be an open neighbourhood of 0 such that $G\text{-mod}_F(f) = G_k\text{-mod}_{\Phi_k}(0) = G_k\text{-mod}_{\Phi_k}(V)$. It follows from Corollary A.12(ii) that

$$G_k\text{-mod}_{\Phi_k}(V) = \max_{i=1, \dots, q} \{G_k\text{-mod}_{\Phi_k^{(i)}}(S_i)\}.$$

Moreover since the $h_{s_i}^{(i)}$ are modular, we can see, with the notations of Definition A.6, that for each $s_i \in S_i$ the set $V_{G_k, \Phi_k^{(i)}}(s_i)$ is finite and hence $G_k\text{-mod}_{\Phi_k^{(i)}}(S_i) = G_k\text{-par}_{\Phi_k^{(i)}}(S_i) = \dim S_i$. This completes the proof. \square

Combining Propositions 2.12 and 2.15 we get

Corollary 2.16. *Let $f \in K[[\mathbf{x}]]$ be such that $\mu(f) < \infty$ (resp. $\tau(f) < \infty$) and let $G = \mathcal{R}$ (resp. $G = \mathcal{K}$). Let $f_t(\mathbf{x}) = F(\mathbf{x}, t)$ be a G -complete unfolding of f at t_0 over T (e.g. F an algebraic representative of a G -versal deformation with trivial section of f as in Proposition 2.14). Assume that there are finitely many G -modular families $h_{s_i}^{(i)}(\mathbf{x}) = H^{(i)}(\mathbf{x}, s_i) \in \mathcal{O}(S_i)[[\mathbf{x}]]$ and a neighbourhood $W \subset T$ of t_0 such that for each open neighbourhood $V \subset W$ of t_0 we have*

$$\bigcup_{t \in V} G \cdot f_t = \bigcup_i \bigcup_{s_i \in S_i} G \cdot h_{s_i}^{(i)}.$$

Then $G\text{-mod}(f) = \max\{\dim S_i\}$.

Proof. By Proposition 2.14 an algebraic representative of any G -versal deformation (with section) of f is G -complete. By Proposition 2.12 $G\text{-mod}(f) = G\text{-mod}_F(f)$. The rest follows from Proposition 2.15(ii). \square

Note that in his classification of right simple, unimodal and bimodal singularities Arnol'd constructed parametrized *normal forms*, being actually \mathcal{R} -modular families of dimension 0, 1 and 2.

The above corollary makes precise (and proves) the statement by Wall [Wal83] for complex analytic singularities saying: “if the set of germs f_t (t small) at points x near 0 falls into finitely many families of right (resp. contact) equivalence classes, each depending on r parameters (at most) then f is right (resp. contact) r -modal (at most).”

Corollary 2.17. *Let $f \in K[[\mathbf{x}]]$ be such that $\mu(f) < \infty$ resp. $\tau(f) < \infty$. f is G -simple iff it is of finite G -unfolding type, i.e. there exists a finite set \mathcal{F} of G -classes of singularities satisfying: for any (or, equivalently, for one G -complete) unfolding $F(\mathbf{x}, t)$ of f at t_0 over an affine variety T , there exists a Zariski open neighbourhood V of $t_0 \in T$, such that the set of G -classes of singularities of $F(\mathbf{x}, t)$, $t \in V$, belongs to the set \mathcal{F} .*

Proposition 2.18. *Let $f \in K[[\mathbf{x}]] = K[[x_1, \dots, x_n]]$ be such that $\mu(f) < \infty$ (resp. $\tau(f) < \infty$) and let $G = \mathcal{R}$ (resp. $G = \mathcal{K}$). Let $m = mt(f)$ be the multiplicity of f . Then*

$$G\text{-mod}(f) \geq \binom{n+m-1}{m} - n^2.$$

In particular,

- (i) if $n = 2$, then $G\text{-mod}(f) \geq m - 3$;
- (ii) if $n \geq 3$ and $m \geq 3$, then $G\text{-mod}(f) \geq \frac{m^2 + 3m - 16}{2}$.

Proof. Let k be sufficiently large for f w.r.t. G . Then $k \geq m$ and $G\text{-mod}(f) = G_k\text{-mod}(j^k f)$. Put $X := \mathfrak{m}^m / \mathfrak{m}^{k+1} \subset J_k$. It follows from Proposition A.4(iii) that

$$G\text{-mod}(f) \text{ in } J_k \geq G\text{-mod}(f) \text{ in } X.$$

Let the linear group $G' := GL(n, K)$ act on $X' := \mathfrak{m}^m / \mathfrak{m}^{m+1}$ by $G' \times X' \rightarrow X'$, $(A, g(\mathbf{x})) \mapsto g(A\mathbf{x})$. Consider the projection $p : X \rightarrow X'$. It is easy to see that p is open and $G \cdot g \subset p^{-1}(G' \cdot p(g))$ for

all $g \in X$. Then Proposition A.4(iv) yields

$$G\text{-mod}(g) \geq G'\text{-mod}(p(g)), \forall g \in X.$$

In order to prove the proposition, it is sufficient to show that

$$G'\text{-mod}(p(g)) \geq \binom{n+m-1}{m} - n^2, \forall g \in X.$$

Indeed, it is easy to see that

$$\dim X' = \binom{n+m-1}{m} \text{ and } \dim GL(n, K) = n^2.$$

Hence, by Proposition A.4(i),

$$\begin{aligned} GL(n, K)\text{-mod}(p(g)) &\geq \dim X' - \dim GL(n, K) \\ &= \binom{n+m-1}{m} - n^2, \end{aligned}$$

which completes the proof.

(i) and (ii) follow from explicit calculations. □

3. CLASSIFICATION OF RIGHT SIMPLE SINGULARITIES

In this section we classify the right simple singularities $f \in \mathfrak{m} \subset K[[x_1, \dots, x_n]]$ for K an algebraically closed field of characteristic $p > 0$. The classification of contact simple singularities was done in [GK90]. In contrast to $\text{char}(K) = 0$, where the classification of right simple and contact simple singularities coincides, the classification is very different in positive characteristic. For example, for every $p > 0$, there are only finitely many classes of right simple singularities and for $p = 2$ only the A_1 -singularity in an even number of variables is right simple. The classification of right simple singularities is summarized in Theorems 3.1, 3.2 and 3.3. Note that $f \in \mathfrak{m} \setminus \mathfrak{m}^2 \Leftrightarrow \mu(f) = 0$ and then $f \sim_r x_1$ (hence right simple) by the implicit function theorem. We may therefore assume in the following that $f \in \mathfrak{m}^2$.

Theorem 3.1. *Let $\text{char}K = p > 0$. Let $f \in \mathfrak{m}^2 \subset K[[x]]$ be a univariate power series such that its Milnor number $\mu := \mu(f)$ is finite. Then*

$$\mathcal{R}\text{-mod}(f) = [\mu/p]$$

is the integer part of μ/p . In particular, f is right simple if and only if $\mu < p$, and then $f \sim_r x^{\mu+1}$.

Theorem 3.2. *Let $p = \text{char}(K) > 2$.*

- (i) *A plane curve singularity $f \in \mathfrak{m}^2 \subset K[[x, y]]$ is right simple if and only if it is right equivalent to one of the following normal forms*

Name	Normal form
A_k	$x^2 + y^{k+1} \quad 1 \leq k \leq p - 2$
D_k	$x^2y + y^{k-1} \quad 4 \leq k < p$
E_6	$x^3 + y^4 \quad 3 < p$
E_7	$x^3 + xy^3 \quad 3 < p$
E_8	$x^3 + y^5 \quad 5 < p$

Table 3.2 (i)

(ii) A hypersurface singularity $f \in \mathfrak{m}^2 \subset K[[x_1, \dots, x_n]]$, $n \geq 3$, is right simple if and only if it is right equivalent to one of the following normal forms

Normal form
$g(x_1, x_2) + x_3^2 + \dots + x_n^2 \mid g$ is one of the singularities in Table 3.2 (i)

Table 3.2 (ii)

Theorem 3.3. Let $p = \text{char}(K) = 2$. A hypersurface singularity $f \in \mathfrak{m}^2 \subset K[[x_1, \dots, x_n]]$ with $n \geq 2$, is right simple if and only if n is even and if it is right equivalent to

$$A_1 : x_1x_2 + x_3x_4 + \dots + x_{n-1}x_n.$$

The following interesting corollary follows immediately from the classification of right simple singularities.

Corollary 3.4. For any $p = \text{char}(K) > 0$ there are only finitely many right simple singularities $f \in \mathfrak{m}^2 \subset K[[x_1, \dots, x_n]]$ (up to right equivalence). For $p = 2$, either no or exactly one right simple singularity exists.

The corollary also shows that if $f_k, k \geq 1$, is any sequences of simple singularities in positive characteristic then the sequence of Milnor numbers $\mu(f_k), k \geq 1$, is bounded. Note that this is wrong in characteristic zero since the $A_k, D_k, k \geq 1$, with Milnor number k , are all simple. We like to pose the following conjecture:

Conjecture 3.5. Let $f_k \in K[[x_1, \dots, x_n]]$, $\text{char}(K) = p > 0$, be a sequence of isolated singularities with Milnor number going to infinity if $k \rightarrow \infty$. Then the right modality of f_k goes to infinity.

3.1. Univariate singularities in positive characteristic. Let $K[[x]]$ be the ring of univariate formal power series. It is obvious that $G\text{-mod}(f) = 0$ for all $f \in K[[x]]$ if either $p := \text{char}(K) = 0$ or $G = \mathcal{K}$. For the complete classification of univariate singularities with any modality and for the proof of Theorem 3.1 we refer to [Ng13, Thm. 4.2.8], [Ng12, Thm. 3.1]. Here we prove only the second part, i.e. f is right simple iff $\mu < p$ and then $f \sim_r x^{\mu+1}$. The “if”-statement follows easily from the upper semi-continuity of the Milnor number and the following fact: If $p \nmid mt(f)$ (in particular, if $\mu < p$) then $f \sim_r x^{mt(f)}$ (cf. [BGM12, Cor. 1]).

It suffices to show that if $\mu \geq p$ then $\mathcal{R}\text{-mod}(f) \geq 1$. Indeed, since $\mu \geq p$, $m := mt(f) \geq p$.

If $m = p$ then we may assume that $f = x^p + a_{p+1}x^{p+1} + \dots \in K[[x]]$. Consider the unfolding

$$f_t = f + tx^{p+1} = x^p + (t + a_{p+1})x^{p+1} + \dots$$

of f . We show that $f_t \sim_r f_{t'}$ implies $t = t'$. If $\varphi(x) = u_1x + u_2x^2 + \dots$ is in \mathcal{R} then $u_1 \neq 0$ and

$$\varphi(f_t) = u_1^p x^p + u_2^p x^{2p} + \dots + (t + a_{p+1})u_1^{p+1} x^{p+1} + \dots$$

If $\varphi(f_t) = f_{t'}$ then $u_1^p = 1$, hence $u_1 = 1$, and $t = t'$. This implies that $\mathcal{R}\text{-mod}(f) \geq 1$ by Corollary 2.17.

Now, assume that $m > p$ and consider the unfolding $g_t := G(x, t) := f + t \cdot x^p$ of f at 0 over \mathbb{A}^1 . By Proposition 2.7, there exists an open neighbourhood V of 0 in \mathbb{A}^1 such that $\mathcal{R}\text{-mod}(g_t) \leq \mathcal{R}\text{-mod}(f)$ for all $t \in V$. Take a $t_0 \in V \setminus \{0\}$, then the above case with $mt = p$ yields that $\mathcal{R}\text{-mod}(g_{t_0}) \geq 1$ since $mt(g_{t_0}) = p$, and hence

$$\mathcal{R}\text{-mod}(f) \geq \mathcal{R}\text{-mod}(g_{t_0}) \geq 1.$$

3.2. Right simple plane curve singularities in characteristic > 2 . Here and in the next section let $f \in K[[x, y]]$, $mt(f)$ its multiplicity, and $\mu = \mu(f)$ its Milnor number, which we assume to be finite. Let $p = \text{char}(K)$.

Proposition 3.6. *Let $mt(f) = 2$ and $p > 2$.*

- (i) *If $\mu < p - 1$, then $f \sim_r A_\mu$ and f is right simple.*
- (ii) *If $\mu \geq p - 1$, then f is not right simple.*

Proof. Since $mt(f) = 2$ and $p > 2$, it follows from the right splitting lemma (Lemma 3.9) that f is right equivalent to $x^2 + g(y)$ (with $g(y) = y^2$ if $\text{crk}(f) = 0$ (case A_1) and $mt(g) \geq 3$ if $\text{crk}(f) = 1$). Here $\text{crk}(f)$ denotes the corank of the Hessian of f , see Section 3.3.

(i) If $\mu < p - 1$ then $mt(g) < p$. By Theorem 3.1, $g \sim_r y^{mt(g)}$ and hence $f \sim_r A_\mu$. Moreover, Theorem 3.1 yields that g is right simple and so is f by Lemma 3.11(iii).

(ii) If $\mu \geq p - 1$, then $mt(g) \geq p$. Combining Theorem 3.1 and Lemma 3.11(iii) we get that f is not right simple. \square

Proposition 3.7. *Let $p > 3$, let $mt(f) = 3$ and f_3 be the tangent cone (i.e. the homogeneous component of degree 3) of f . Let $r(f_3)$ be the number of linear factors of f_3 .*

- (i) *If $r(f_3) \geq 2$ then $f \sim_r x^2y + g(y)$ with $mt(g) = \mu - 1 \geq 3$. If additionally $4 \leq \mu < p$, then $f \sim_r D_\mu$ and f is right simple.*
- (ii) *If $r(f_3) = 1$, $p = 5$ and $6 \leq \mu \leq 7$ then $f \sim_r E_\mu$ and f is right simple.*
- (iii) *If $r(f_3) = 1$, $p > 5$ and $6 \leq \mu \leq 8$ then $f \sim_r E_\mu$ and f is right simple.*

Proof. This may be proved in much the same way as [GLS06, Thm. I.2.51, Cor. I.2.52, Thm. I.2.53, Cor. I.2.54], by applying the finite determinacy theorem in positive characteristic [BGM12, Thm. 2.1]. For details we refer to [Ng13, Prop. 4.3.5]. \square

Proposition 3.8. *Let $mt(f) = 3$. Let $r(f_3)$ be the number of linear factors of f_3 . Then f is not right simple if*

- (i) either $p = 3$;
- (ii) or $p > 3$, $r(f_3) \geq 2$ and $\mu \geq p$;
- (iii) or $p > 5$, $r(f_3) = 1$ and $\mu > 8$;
- (iv) or $p = 5$, $r(f_3) = 1$ and $\mu \geq 8$.

Proof. (i) We consider the unfolding $F(x, y, t) = f + t \cdot x^2$ of f at 0 over \mathbb{A}^1 . Since $mt(f) = 3$ and since $p = 3$, it is easy to see that $\mu(f_t) > 2$ for all $t \neq 0$. Proposition 3.6(ii) yields that $f_t, t \neq 0$, is not right simple and hence neither is f by Proposition 2.7.

(ii) By Proposition 3.7(i), $f \sim_r x^2y + g(y)$ with $mt(g) = \mu - 1$. It suffices to show that $h := x^2y + g(y)$ is not right simple. We write

$$g(y) = a \cdot y^{\mu-1} + \text{higher terms}, a \neq 0$$

and consider the unfolding

$$h_t := H(x, y, t) = \begin{cases} x^2y + g(y) + tx^2 + ty^p & \text{if } \mu > p \\ x^2y + g(y) + at^2x^2 + 2atxy^{(p-1)/2} & \text{if } \mu = p \end{cases}$$

of h at 0 over \mathbb{A}^1 . It is easy to see that $\mu(h_t) \geq p$ for all $t \neq 0$ (in fact, $\mu(h_t) = p$ for almost all t). It follows from Proposition 3.6 that $h_t, t \neq 0$, is not right simple and hence neither is h due to Proposition 2.7.

(iii) This is done by the same argument as in [GLS06, Thm. I.2.55(2)(ii)].

(iv) Since $r(f_3) = 1$ and $\mu \geq 8$, using the same argument as in [GLS06, Thm. I.2.53], we get

$$f \sim_r g := x^2y + \alpha y^5 + \beta xy^4 + h(x, y)$$

with $\alpha, \beta \in K$ and $h \in \mathfrak{m}^6$. Consider the unfolding $g_t := G(x, y, t) = g(x, y) + t \cdot xy^4$ of g at 0 over \mathbb{A}^1 and assume that $g_t \sim_r g_{t'}$, i.e. there exists an automorphism

$$\begin{aligned} \Phi : K[[x, y]] &\longrightarrow K[[x, y]] \\ x &\mapsto \varphi = \sum a_{ij} x^i y^j \\ y &\mapsto \psi = \sum b_{ij} x^i y^j \end{aligned}$$

such that $g_t(x, y) = g_{t'}(\varphi, \psi)$. By a simple calculation we conclude that $(\beta + t)^3 = (\beta + t')^3$ and hence, for fixed t , $g_t \sim_r g_{t'}$ for at most three values of t' . It follows from Corollary 2.17 that g is not right simple and hence neither is f . \square

Proof of Theorem 3.2(i). It follows from Propositions 2.18, 3.6, 3.7 and 3.8. \square

3.3. Right simple hypersurface singularities in characteristic > 2 . Our aim is to prove Theorem 3.2(ii). Let $f \in K[[\mathbf{x}]] = K[[x_1, \dots, x_n]]$. We denote by

$$H(f) := \left(\frac{\partial^2 f}{\partial x_i \partial x_j} (0) \right)_{i,j=1,\dots,n} \in \text{Mat}(n \times n, K)$$

the *Hessian (matrix)* of f and by $\text{crk}(f) := n - \text{rank}(H(f))$ the *corank* of f .

Lemma 3.9 (Right splitting lemma in characteristic different from 2). *If $f \in \mathfrak{m}^2 \subset K[[\mathbf{x}]]$, $\text{char}(K) > 2$, has corank $\text{crk}(f) = k \geq 0$, then*

$$f \sim_r g(x_1, \dots, x_k) + x_{k+1}^2 + \dots + x_n^2$$

with $g \in \mathfrak{m}^3$. g is called the residual part of f , it is uniquely determined up to right equivalence.

Proof. cf. [GLS06, Thm. I.2.47]. The proof in [GLS06] is given for $K = \mathbb{C}$ but works in characteristic different from 2. \square

Lemma 3.10. *Let $p = \text{char}(K) > 2$ and let*

$$f_i(x_1, \dots, x_n) = x_n^2 + f'_i(x_1, \dots, x_{n-1}) \in \mathfrak{m}^2 \subset K[[x_1, \dots, x_n]], \quad i = 1, 2.$$

Then $f_1 \sim_r f_2$ if and only if $f'_1 \sim_r f'_2$.

Proof. The direction, $f'_1 \sim_r f'_2 \Rightarrow f_1 \sim_r f_2$ is obvious. We now assume that $f_1 \sim_r f_2$. Then $\text{crk}(f_1) = \text{crk}(f_2) := k$ and therefore $\text{crk}(f'_1) = \text{crk}(f'_2) = k$. It follows from Lemma 3.9 that

$$f'_i \sim_r g_i(x_1, \dots, x_k) + x_{k+1}^2 + \dots + x_{n-1}^2, \quad i = 1, 2.$$

and hence

$$f_i \sim_r g_i(x_1, \dots, x_k) + x_{k+1}^2 + \dots + x_{n-1}^2 + x_n^2.$$

This implies

$$g_1(x_1, \dots, x_k) + x_{k+1}^2 + \dots + x_n^2 \sim_r g_2(x_1, \dots, x_k) + x_{k+1}^2 + \dots + x_n^2$$

since $f_1 \sim_r f_2$. The uniqueness of g_i shows that $g_1 \sim_r g_2$, i.e. there exists an automorphism $\Phi' \in \text{Aut}_K(K[[x_1, \dots, x_k]])$ such that $\Phi'(g_1) = g_2$. The automorphism

$$\begin{aligned} \Phi : K[[x_1, \dots, x_{n-1}]] &\longrightarrow K[[x_1, \dots, x_{n-1}]] \\ x_i &\mapsto \Phi'(x_i), \quad i = 1, \dots, k \\ x_j &\mapsto x_j, \quad j = k+1, \dots, n-1 \end{aligned}$$

yields that

$$g_1(x_1, \dots, x_k) + x_{k+1}^2 + \dots + x_{n-1}^2 \sim_r g_2(x_1, \dots, x_k) + x_{k+1}^2 + \dots + x_{n-1}^2$$

and hence $f'_1 \sim_r f'_2$. This completes the proof. \square

Lemma 3.11. *Let $p = \text{char}(K) > 2, n \geq 2$, and let*

$$f(x_1, \dots, x_n) = x_n^2 + f'(x_1, \dots, x_{n-1}) \in \mathfrak{m}^2 \subset K[[x_1, \dots, x_n]]$$

be such that $\mu(f) < \infty$.

- (i) *Let $F'(\mathbf{x}', t) \in \langle x_1, \dots, x_{n-1} \rangle \subset \mathcal{O}(T)[[x_1, \dots, x_{n-1}]]$ be an unfolding of f' at t_0 over an affine variety T and let $F(\mathbf{x}, t) = x_n^2 + F'(\mathbf{x}', t)$. Then*

$$\mathcal{R}\text{-mod}_F(f) = \mathcal{R}\text{-mod}_{F'}(f').$$

(ii) *We have*

$$\mathcal{R}\text{-mod}(f) \text{ in } K[[x_1, \dots, x_n]] = \mathcal{R}\text{-mod}(f') \text{ in } K[[x_1, \dots, x_{n-1}]].$$

Proof. Let k be sufficiently large for f and for f' w.r.t. \mathcal{R} . Let \mathfrak{m}' be the maximal ideal in $K[[\mathbf{x}']]$ and let $J'_k := K[[\mathbf{x}']]/\mathfrak{m}'^{k+1}$.

(i) Consider the morphisms

$$\begin{array}{ccc} h_k : T & \longrightarrow & J_k & \text{and} & h'_k : T & \longrightarrow & J'_k \\ t & \mapsto & j^k f_t & & t & \mapsto & j^k f'_t. \end{array}$$

and $p : J_k \rightarrow J'_k$ the natural projection. Then $h'_k = p \circ h_k$ and

$$h_k^{-1}(\mathcal{R} \cdot h_k(t)) = h_k'^{-1}(\mathcal{R} \cdot h'_k(t))$$

by Lemma 3.10. It follows from Corollary A.11 that $\mathcal{R}\text{-mod}_{h_k}(t_0) = \mathcal{R}\text{-mod}_{h'_k}(t_0)$, and hence $\mathcal{R}\text{-mod}_F(f) = \mathcal{R}\text{-mod}_{F'}(f')$.

(ii) Let $\{g'_1(\mathbf{x}'), \dots, g'_l(\mathbf{x}')\}$ be a system of generators of $\mathfrak{m}'/\mathfrak{m}' \cdot j(f')$. Then $\{x_n, g_1(\mathbf{x}), \dots, g_l(\mathbf{x})\}$ with $g_i(\mathbf{x}) = g'_i(\mathbf{x}')$, is a system of generators of $\mathfrak{m}/\mathfrak{m} \cdot j(f)$. Proposition 2.14 yields that

$$F'(\mathbf{x}', t) = f' + \sum_{i=1}^l t_i g'_i(\mathbf{x}') \quad (\text{resp. } F_1(\mathbf{x}, t) = f + \sum_{i=1}^l t_i g_i(\mathbf{x}) + t_{l+1} x_n)$$

is a right complete unfoldings of f' (resp. of f) over \mathbb{A}^l (resp. \mathbb{A}^{l+1}), i.e.

$$\mathcal{R}\text{-mod}(f) = \mathcal{R}\text{-mod}_{F_1}(f) \quad (\text{resp. } \mathcal{R}\text{-mod}(f') \text{ in } K[[\mathbf{x}']] = \mathcal{R}\text{-mod}_{F'}(f'))$$

due to Proposition 2.12. Note that $F_1(\mathbf{x}, t) \sim_r x_1$ and therefore $\mathcal{R}\text{-mod}(F_1(\mathbf{x}, t)) = 0$ for all $t = (t_1, \dots, t_{l+1}) \in \mathbb{A}^{l+1}$ with $t_{l+1} \neq 0$.

Consider the inclusion $\mathbb{A}^l \subset \mathbb{A}^{l+1}$, $t = (t_1, \dots, t_l) \mapsto (t_1, \dots, t_l, 0)$ and the unfolding

$$F = f + \sum_{i=1}^l t_i g_i(\mathbf{x})$$

of f at 0 over \mathbb{A}^l . Since $\mathcal{R}\text{-mod}(F_1(\mathbf{x}, t)) = 0$ for all $t \in \mathbb{A}^{l+1} \setminus \mathbb{A}^l$, using the same argument as in the proof of Proposition A.4(iii) (see also [Ng13, Prop. 3.2.6]) we obtain that $\mathcal{R}\text{-mod}_{F_1}(f) = \mathcal{R}\text{-mod}_F(f)$. This implies, by (i), that

$$\mathcal{R}\text{-mod}_{F_1}(f) = \mathcal{R}\text{-mod}_F(f) = \mathcal{R}\text{-mod}_{F'}(f')$$

and hence $\mathcal{R}\text{-mod}(f) = \mathcal{R}\text{-mod}(f')$. □

Proof of Theorem 3.2(ii). The “if”-statement follows from Theorem 3.2(i) and Lemma 3.11. We now consider any simple singularity $f \in \mathfrak{m}^2 \subset K[[\mathbf{x}]]$. Then, by the splitting lemma,

$$f \sim_r f'(x_1, \dots, x_k) + x_{k+1}^2 + \dots + x_n^2$$

with $f' \in \langle x_1, \dots, x_k \rangle^3$ and $k = \text{crk}(f)$. Again by Lemma 3.11,

$$\mathcal{R}\text{-mod}(f') = \mathcal{R}\text{-mod}(f) = 0.$$

It follows from Proposition 2.18 that

$$0 = \mathcal{R}\text{-mod}(f') \geq \binom{m+k-1}{m} - k^2,$$

where $m = mt(f') \geq 3$. This implies that $k \leq 2$, i.e.

$$f \sim_r g(x_1, x_2) + x_3^2 + \dots + x_n^2,$$

for some simple singularity $g \in K[[x_1, x_2]]$. The proof thus follows from Theorem 3.1, 3.2(i) and Lemma 3.10. \square

3.4. Right simple hypersurface singularities in characteristic 2. Let $p = \text{char}(K) = 2$ and let $n \geq 2$.

Lemma 3.12 (Right splitting lemma in characteristic 2). *Let $f \in \mathfrak{m}^2 \subset K[[x_1, \dots, x_n]]$, $n \geq 2$. Then there exists an l , $0 \leq 2l \leq n$ such that*

$$f \sim_r x_1x_2 + x_3x_4 + \dots + x_{2l-1}x_{2l} + g(x_{2l+1}, \dots, x_n)$$

with $g \in \langle x_{2l+1}, \dots, x_n \rangle^3$ or $g \in x_{2l+1}^2 + \langle x_{2l+1}, \dots, x_n \rangle^3$ if $2l < n$. g is called the residual part of f , it is uniquely determined up to right equivalence.

Proof. It follows easily from [GK90, Lemmas 1 and 2]. \square

Lemma 3.13. *Let $\mu(f) < \infty$ and*

$$f(x_1, \dots, x_n) = x_{n-1}x_n + f'(x_1, \dots, x_{n-2}) \in \mathfrak{m}^2 \subset K[[x_1, \dots, x_n]].$$

Then

$$\mathcal{R}\text{-mod}(f) \text{ in } K[[x_1, \dots, x_n]] = \mathcal{R}\text{-mod}(f') \text{ in } K[[x_1, \dots, x_{n-2}]].$$

Proof. By using the same arguments as in the proof of Lemma 3.11. \square

Remark 3.14. Since $\mu(x_1x_2) = 1$, $x_1x_2 \in K[[x_1, x_2]]$ is right 2-determined and any unfolding of x_1x_2 is either right equivalent to itself or smooth. Hence x_1x_2 is right simple.

Proposition 3.15. *Let $f \in \mathfrak{m}^2 \subset K[[\mathbf{x}]]$ with $\mu(f) < \infty$. Then f is not right simple if*

- (i) either $f = x_1^2 + g(x_1, \dots, x_n) \in K[[\mathbf{x}]]$ with $g \in \mathfrak{m}^3$,
- (ii) or $f \in \mathfrak{m}^3$.

Proof. (i) Let $k \geq 3$ be sufficiently large for f w.r.t. \mathcal{R} and let $X := \mathfrak{m}^2/\mathfrak{m}^{k+1}$. Then $\mathcal{R}\text{-mod}(f) = \mathcal{R}_k\text{-mod}(j^k f)$. Let

$$Y := x_1^2 + \mathfrak{m}^3/\mathfrak{m}^{k+1} \subset J_k, \quad Y' := x_1^2 + \mathfrak{m}^3/\mathfrak{m}^4$$

and let

$$\mathcal{H} := \{\Phi \in \mathcal{R}_k \mid \Phi(x_1) = x_1\}, \quad \mathcal{H}' := \{\Phi \in \mathcal{R}_1 \mid \Phi(x_1) = x_1\}.$$

Then \mathcal{H} (resp. \mathcal{H}') acts on Y (resp. Y') by $(\Phi, y) \mapsto \Phi(y)$ and we have

$$i^{-1}(\mathcal{R}_k \cdot i(y)) = \mathcal{H} \cdot y \subset p^{-1}(\mathcal{H}' \cdot p(y)) \quad \forall y \in Y'$$

with the inclusion $i : Y \hookrightarrow X$ and the projection $p : Y \twoheadrightarrow Y'$. It follows from Proposition A.4(iv) that

$$\mathcal{R}_k\text{-mod}(y) \geq \mathcal{H}\text{-mod}(y) \geq \mathcal{H}'\text{-mod}(p(y)), \forall y \in Y.$$

Moreover, Proposition A.4(i) yields that

$$\mathcal{H}'\text{-mod}(p(y)) \geq \dim Y' - \dim \mathcal{H}' = \binom{n+2}{3} - n(n-1) \geq 1.$$

This implies that $\mathcal{R}_k\text{-mod}(y) \geq 1$ for all $y \in Y$ and hence $\mathcal{R}\text{-mod}(f) \geq 1$.

(ii) By (i), f_t is not right simple for all $t \neq 0$, where $f_t(\mathbf{x}) := f(\mathbf{x}) + tx_1^2$ is an unfolding of f at 0 over \mathbb{A}^1 . Hence Proposition 2.7 yields that f is not right simple. \square

Proof of Theorem 3.3. The “if”-statement is obvious. Now, take a right simple singularity $f \in \mathfrak{m}^2 \subset K[[\mathbf{x}]]$. Then $mt(f) = 2$ by Proposition 2.18. The splitting lemma (Lemma 3.12) yields that f is right equivalent to

$$x_1x_2 + x_3x_4 + \dots + x_{2l-1}x_{2l} + g(x_{2l+1}, \dots, x_n)$$

with $g \in \langle x_{2l+1}, \dots, x_n \rangle^3$ or $g \in x_{2l+1}^2 + \langle x_{2l+1}, \dots, x_n \rangle^3$ if $2l < n$. Combining Lemma 3.13 and Proposition 3.15 we obtain that $2l = n$, which proves the theorem. \square

APPENDIX A.

A.1. Modality for algebraic group actions. Let an algebraic group G act on the variety X . We define the notion of number of G -parameters and show that it coincides with the G -modality, which proves the independence of modality of the Rosenlicht stratification. Moreover if X' is any variety (without G -action) and if $h : X' \rightarrow X$ is a morphism, we generalize these notions to the equivalence relation induced by h on X' . This allows us to use deformation theory. As consequences, we give interesting properties of modality, which are used in Section 2.

Definition A.1. Let $U \subset X$ be an open neighbourhood of $x \in X$ and W be constructible in X . We introduce

$$\begin{aligned} \dim_x W &:= \max\{\dim Z \mid Z \text{ an irreducible component of } W \text{ containing } x\}, \\ U(i) &:= U_G(i) := \{y \in U \mid \dim_y(U \cap G \cdot y) = i\}, i \geq 0, \\ G\text{-par}(U) &:= \max_{i \geq 0}\{\dim U(i) - i\}, \end{aligned}$$

and call

$$G\text{-par}(x) := \min\{G\text{-par}(U) \mid U \text{ a neighbourhood of } x\}$$

the *number of G -parameters* of x (in X).

The following proposition is a special case of Proposition A.7 (with $h = \text{id}$), which is proven below.

Proposition A.2. *We have $G\text{-par}(U) = G\text{-mod}(U)$ and therefore $G\text{-par}(x) = G\text{-mod}(x)$ for all $x \in X$.*

Corollary A.3. $G\text{-mod}(U)$ and $G\text{-mod}(x)$ are independent of the Rosenlicht stratification of X .

We call $x, y \in X$ G -equivalent, denoted by $x \sim_G y$, if they lie in the same G -orbit.

Proposition A.4. Let the algebraic group G act on the variety X .

- (i) If G and X are both irreducible then $G\text{-mod}(x) \geq \dim X - \dim G$.
- (ii) For any $y \sim_G x$, $G\text{-mod}(x) = G\text{-mod}(y)$.
- (iii) If the subvariety $X' \subset X$ is invariant under G and if $x \in X'$ then

$$G\text{-mod}(x) \text{ in } X \geq G\text{-mod}(x) \text{ in } X'.$$

Equality holds if $G\text{-mod}(x) \text{ in } X' \geq G\text{-mod}(y)$, $\forall y \in X \setminus X'$.

- (iv) Let the algebraic group G' act on the variety X' and let $p : X \rightarrow X'$ be a morphism of varieties.

- (1) If p is open and if

$$G \cdot x \subset p^{-1}(G' \cdot p(x)), \quad \forall x \in X,$$

then

$$G\text{-mod}(x) \geq G'\text{-mod}(p(x)), \quad \forall x \in X.$$

- (2) If p is equivariant (i.e. $G \cdot x = p^{-1}(G' \cdot p(x))$, $\forall x \in X$), then

$$G\text{-mod}(x) \leq G'\text{-mod}(p(x)), \quad \forall x \in X.$$

In particular, if p is open and equivariant then $G\text{-mod}(x) = G'\text{-mod}(p(x))$, $\forall x \in X$.

For the elementary but not so short proof, using $G\text{-mod} = G\text{-par}$, we refer to [Ng13, Prop. 3.2.4–3.2.7].

In order to relate the notion of modality to deformation theory we introduce the notion of G -modality w.r.t. a morphism from an arbitrary variety to the G -variety X .

Definition A.5. Let $\{X_j, j = 1, \dots, s\}$ be a Rosenlicht stratification of X under G with projections $p_j : X_j \rightarrow X_j/G$.

Let $h : X' \rightarrow X$ be a morphism of algebraic varieties, and let U' be an open neighbourhood of $x' \in X'$. Set $X'_j := h^{-1}(X_j)$, $U'_j := U' \cap X'_j$. We define

$$G\text{-mod}_h(U') := \max_{j=1, \dots, s} \{\dim(p_j(h(U'_j)))\},$$

and call

$$G\text{-mod}_h(x') := \min\{G\text{-mod}(U') \mid U' \text{ a neighbourhood of } x' \text{ in } X'\}$$

the G -modality of x' w.r.t. h .

Definition A.6. Let the algebraic group G act on the variety X , let $h : X' \rightarrow X$ be a morphism of algebraic varieties and let U' be an open neighbourhood of $x' \in X'$. For $u' \in U'$ and each $i \geq 0$

we define

$$\begin{aligned} V_{G,h}(u') &:= \{v' \in X' \mid G \cdot h(v') = G \cdot h(u')\} = h^{-1}(G \cdot h(u')), \\ U'(i) &:= U'_{G,h}(i) := \{u' \in U' \mid \dim_{u'}(U' \cap V_{G,h}(u')) = i\}, \\ G\text{-par}_h(U') &:= \max_{i \geq 0} \{\dim U'_{G,h}(i) - i\}, \end{aligned}$$

and call

$$G\text{-par}_h(x') := \min\{G\text{-par}_h(U') \mid U' \text{ a neighbourhood of } x' \in X'\}$$

the number of G -parameters of x' w.r.t. h .

If we call $x', y' \in X'$ equivalent if $h(x')$ and $h(y')$ ly in the same orbit in X , we get an equivalence relation on X' with equivalence class of x' equal to

$$V_{G,h}(x') = h^{-1}(G \cdot h(x')).$$

It follows in particular that each equivalence class is locally closed in X' .

Proposition A.7. *We have $G\text{-par}_h(U') = G\text{-mod}_h(U')$ and therefore $G\text{-par}_h(x') = G\text{-mod}_h(x')$ for all $x' \in X'$.*

For the proof we need the following properties of fibers of a morphism ([Mum88], [Bor91]). Let $f : X \rightarrow Y$ be a morphism of (not necessarily irreducible) algebraic varieties. First, Chevalley's theorem says that if $W \subset X$ is constructible in X , then $f(W)$ is constructible in Y . Secondly, the function $x \mapsto e(x) := \dim_x f^{-1}(f(x))$ is *upper semi-continuous*, i.e. for all integers n the set $\{x \in X \mid e(x) \geq n\}$ is closed. For the proof of these statements and the following lemma some well known results in [Mum88, I.8] for irreducible varieties, are used. For details see [Ng13, Cor. 3.1.7 and 3.1.8].

Lemma A.8. *Let $f : X \rightarrow Y$ be a morphism algebraic varieties and let $e : X \rightarrow \mathbb{N}$ be the function defined by $x \mapsto e(x) := \dim_x f^{-1}(f(x))$.*

(i) *If $e(x)$ is constant, say $e(x) = i$ for all $x \in X$, then*

$$\dim X = i + \dim f(X).$$

(ii) *We have*

$$\max_{i \geq 0} \{\dim(e^{-1}(i)) - i\} \geq \dim f(X).$$

Proof of Proposition A.7. We use the notations of Definition A.6 and consider the composition

$$h_j : U'_j \xrightarrow{h} X_j \xrightarrow{p_j} X_j/G.$$

Note that

$$h_j^{-1}(h_j(x')) = U'_j \cap h^{-1}((G \cdot h(x'))) = U'_j \cap V_{G,h}(x'), \quad \forall x' \in U'_j.$$

By the upper semi-continuity of the functions $e_j : U'_j \rightarrow \mathbb{N}, x' \mapsto \dim_{x'} h_j^{-1}(h_j(x'))$, the sets $e_j^{-1}(i)$ are locally closed and then

$$U'(i) = \bigcup_{j=1}^s e_j^{-1}(i)$$

is constructible in X' for all $i \geq 0$.

Taking an i such that $G\text{-par}_h(U') = \dim U'(i) - i$ and applying Lemma A.8(i) we deduce

$$G\text{-par}_h(U') = \max_j \{\dim e_j^{-1}(i) - i\} = \max_j \{\dim (h_j(e_j^{-1}(i)))\} \leq G\text{-mod}_h(U').$$

Let $j \in \{1, \dots, s\}$ be such that

$$G\text{-mod}_h(U') = \dim (p_j(h(U'_j))) = \dim h_j(U'_j).$$

Then

$$G\text{-par}_h(U') = \max_i \{\dim U'(i) - i\} \geq \max_i \{\dim(e_j^{-1}(i)) - i\} \geq \dim h_j(U'_j) = G\text{-mod}_h(U'),$$

where the second inequality follows from Lemma A.8(ii). Hence $G\text{-par}_h(U') = G\text{-mod}_h(U')$. \square

Let the algebraic group G act on the variety X . The proofs of the following corollaries follow easily from Definition A.6, A.5 and Proposition A.7 (for details we refer to [Ng13, Cor. 3.3.4-3.3.7]).

Corollary A.9. *Let $h : X' \rightarrow X$ be a morphism of algebraic varieties. Then*

$$G\text{-mod}_h(x') \leq G\text{-mod}(h(x')), \quad \forall x' \in X'.$$

Equality holds if for every open neighbourhood U' of x' , there exists an open neighbourhood U of $h(x')$ in X s.t. $U \subset h(U')$. In particular, equality holds if h is open.

Corollary A.10. *Let*

$$g : X'' \xrightarrow{h'} X' \xrightarrow{h} X$$

be morphisms of algebraic varieties. Then

$$G\text{-mod}_g(x'') \leq G\text{-mod}_h(h'(x'')), \quad \forall x'' \in X''.$$

Equality holds if for every open neighbourhood U'' of x'' , there exists an open neighbourhood U' of $h'(x'')$ in X' s.t. $U' \subset h'(U'')$. In particular, equality holds if h' is open.

Corollary A.11. *Let the algebraic groups G resp. G' act on the varieties X resp. X' . Let*

$$h : Y \rightarrow X \text{ and } h' : Y \rightarrow X'$$

be two morphisms of varieties such that

$$h^{-1}(G \cdot h(y)) = h'^{-1}(G' \cdot h'(y)), \quad \forall y \in Y.$$

Then for any open subset $V \subset Y$ we have $G\text{-mod}_h(V) = G'\text{-mod}_{h'}(V)$. Consequently

$$G\text{-mod}_h(y) = G'\text{-mod}_{h'}(y), \quad \forall y \in Y.$$

Corollary A.12. *Let $h : X' \rightarrow X$ and $h_i : Y_i \rightarrow X, i = 1, \dots, k$, be morphisms of varieties and let U' be open in X' satisfying, that for all $x' \in U'$ there exists $y_i \in Y_i$ such that $h(x') \sim_G h_i(y_i)$. Then*

- (i) $G\text{-mod}_h(U') \leq \max\{G\text{-mod}_{h_i}(Y_i) | i = 1, \dots, k\}$.
- (ii) *Assume moreover that for all $y_i \in Y_i$ there exists $x' \in X'$ such that $h(x') \sim_G h_i(y_i)$ then*

$$G\text{-mod}_h(U') = \max\{G\text{-mod}_{h_i}(Y_i) | i = 1, \dots, k\}.$$

A.2. Right and contact groups. The group $\mathcal{R} := \text{Aut}_K(K[[\mathbf{x}]])$ of automorphisms of the local analytic K -algebra $K[[\mathbf{x}]] = K[[x_1, \dots, x_n]]$ is called the *right group*. The *contact group* \mathcal{K} is the semi-direct product of $\text{Aut}_K(K[[\mathbf{x}]])$ with the group $(K[[\mathbf{x}]])^*$ of units in $K[[\mathbf{x}]]$. These groups act on $K[[\mathbf{x}]]$ by

$$\begin{array}{ccc} \mathcal{R} \times K[[\mathbf{x}]] & \longrightarrow & K[[\mathbf{x}]] \quad \text{and} \quad \mathcal{K} \times K[[\mathbf{x}]] \longrightarrow K[[\mathbf{x}]] \\ (\Phi, f) & \mapsto & \Phi(f) \quad \quad \quad ((\Phi, u), f) \mapsto u \cdot \Phi(f). \end{array}$$

Two elements $f, g \in K[[\mathbf{x}]]$ are called *right* (\sim_r) resp. *contact* (\sim_c) *equivalent* iff they belong to the same \mathcal{R} - resp. \mathcal{K} -orbit.

Note that neither \mathcal{R} nor \mathcal{K} are algebraic groups, as they are infinite dimensional. In order to be able to apply the results from the previous section, we have to pass to the jet spaces.

An element Φ in the right group \mathcal{R} is uniquely determined by n power series

$$\varphi_i := \Phi(x_i) = \sum_{j=1}^n a_i^j x_j + \text{terms of higher order}$$

such that $\det(a_i^j) \neq 0$. For each integer k we define the k -jet of Φ ,

$$\Phi_k := (j^k(\varphi_1), \dots, j^k(\varphi_n)).$$

Here

$$j^k : K[[\mathbf{x}]] \rightarrow J_k := K[[\mathbf{x}]]/\mathfrak{m}^{k+1}$$

denotes the canonical projection to the k -jet space J_k and $\mathfrak{m} = \langle \mathbf{x} \rangle$ the maximal ideal of $K[[\mathbf{x}]]$. For $f \in K[[\mathbf{x}]]$, $j^k f := j^k(f)$ is represented in $K[[\mathbf{x}]]$ by the power series expansion of f up to order k . We call

$$\mathcal{R}_k := \{\Phi_k \mid \Phi \in \mathcal{R}\} \text{ respectively } \mathcal{K}_k := \{(\Phi_k, j^k u) \mid (\Phi, u) \in \mathcal{K}\}$$

the k -jet of \mathcal{R} respectively of \mathcal{K} . Note that J_k is an affine space and $\mathcal{R}_k, \mathcal{K}_k$ are affine algebraic groups, all equipped with the Zariski topology. These groups act algebraically on J_k by

$$\begin{array}{ccc} \mathcal{K}_k \times J_k & \longrightarrow & J_k \quad \text{and} \quad \mathcal{K}_k \times J_k \longrightarrow J_k \\ (\Phi_k, j^k f) & \mapsto & j^k(\Phi_k(j^k f)) \quad \quad \quad ((\Phi_k, j^k u), j^k f) \mapsto j^k(j^k u \cdot \Phi_k(j^k f)). \end{array}$$

The first statement (i) of the following lemma says that the Milnor number μ and the Tjurina number τ are semi-continuous w.r.t. unfoldings. Its proof can be adapted from the construction in [GLS06, Thm. I.2.6] by applying [Har77, Thm. 12.8]. The second statement follows from (i).

In order to shorten notations we define a *topology* on $K[[\mathbf{x}]]$ to be the coarsest topology such that all projections $j^k : K[[\mathbf{x}]] \rightarrow J_k$ are continuous, i.e. $V \subset K[[\mathbf{x}]]$ is open iff $j^k(V)$ is open in J_k for all k .

Lemma A.13 (Semi-continuity of μ and τ). *Let $f \in K[[\mathbf{x}]]$ with $\mu(f) < \infty$ resp. $\tau(f) < \infty$.*

- (i) *Let $f_t(\mathbf{x}) = F(\mathbf{x}, t)$ be an unfolding of $f = f_{t_0}$ at t_0 over an affine variety T . Then there exists an open neighbourhood $U \subset T$ of t_0 such that*

$$\mu(f_t) \leq \mu(f), \text{ resp. } \tau(f_t) \leq \tau(f), \forall t \in U.$$

More general, for all $i \in \mathbb{N}$ the sets

$$\{t \in T \mid \mu(f_t) \leq i\} \text{ resp. } \{t \in T \mid \tau(f_t) \leq i\}$$

are open in T .

- (ii) *The functions μ and τ are upper semi-continuous on $K[[\mathbf{x}]]$, i.e. for all $i \in \mathbb{N}$, the sets*

$$U_{\mu,i} := \{f \in K[[\mathbf{x}]] \mid \mu(f) \leq i\} \text{ and } U_{\tau,i} := \{f \in K[[\mathbf{x}]] \mid \tau(f) \leq i\}$$

are open in $K[[\mathbf{x}]]$.

REFERENCES

- [AGV85] Arnol'd, V. I.; Gusein-Zade, S. M.; Varchenko, A. N., *Singularities of differentiable maps*, Vol I. Birkhäuser (1985).
- [Arn72] Arnol'd, V. I., *Normal forms for functions near degenerate critical points, the Weyl groups of A_k, D_k, E_k and Lagrangian singularities*, Functional Anal. Appl. 6 (1972), 254-272.
- [BGM12] Boubakri Y.; Greuel G.-M.; Markwig T., *Invariants of hypersurface singularities in positive characteristic*, Rev. Mat. Complut. 25 (2012), 61-85.
- [Bor91] Borel, A., *Linear Algebraic Groups*, 2nd edition, Graduate Texts in Mathematics 126, Springer (1991).
- [GK90] Greuel G.-M.; Kröning H., *Simple singularities in positive characteristic*, Math. Z. 203 (1990), 339-354.
- [GLS06] Greuel G.-M.; Lossen C.; Shustin E., *Introduction to Singularities and Deformations*, Math. Monographs, Springer-Verlag (2006).
- [Har77] Hartshorne R., *Algebraic Geometry*, Springer (1977).
- [KaS72] Kas, A.; Schlessinger, M., *On the versal deformation of a complex space with an isolated singularity*, Math. Ann. 196 (1972), 23-29.
- [KPPRM78] Kurke H.; Pfister G.; Popescu D.; Roczen M.; Mostowski T., *Die Approximationseigenschaft lokaler Ringe*, Lecture Notes in Math., 634 Springer-Verlag, Berlin-NewYork (1978).
- [MFK82] Mumford, D.; Fogarty, J.; Kirwan, F., *Geometric Invariant Theory*, Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete, 34. Springer-Verlag, Berlin (1982).
- [Mum88] Mumford, D., *The red book of varieties and schemes*, Lecture Notes in Math., 1358, Springer (1988).
- [Na53] Nagata M., *On the theory of Henselian rings*, Nagoya Math. J. 5 (1953), 45-57.
- [Ng12] Nguyen H. D., *The right classification of univariate power series in positive characteristic*, (2013) <http://arxiv.org/abs/1210.2868>
- [Ng13] Nguyen H. D., *Classification of singularities in positive characteristic*, Ph.D. thesis, TU Kaiserslautern, Dr. Hut-Verlag (2013). <http://www.dr.hut-verlag.de/9783843911030.html>
- [Po86] Popescu D., *General Néron desingularization and approximation*, Nagoya Math. J. 104 (1986), 85-115.

- [Ra70] Raynaud M., *Anneaux locaux Henséliens*, Lecture Notes in Math., 169, Springer, Berlin (1970).
- [Ros56] Rosenlicht, M., *Some basic theorems on algebraic groups*, Amer. J. Math. 78 (1956), 401-443.
- [Ros63] Rosenlicht, M., *A remark on quotient spaces*, An. Acad. Brasil. Ci. 35 (1963), 487-489.
- [Wal83] Wall, C. T. C., *Classification of unimodal isolated singularities of complete intersections*, pp 625-640 in Proc. Symp. in Pure Math. 40ii (Singularities) (ed. P. Orlik) Amer. Math. Soc., 1983.

GERT-MARTIN GREUEL
UNIVERSITÄT KAISERSLAUTERN, FACHBEREICH MATHEMATIK, ERWIN-SCHRÖDINGER-STRASSE,
67663 KAISERSLAUTERN
E-mail address: greuel@mathematik.uni-kl.de

NGUYEN HONG DUC
INSTITUTE OF MATHEMATICS, 18 HOANG QUOC VIET ROAD, CAU GIAY DISTRICT
10307, HANOI.
E-mail address: nhduc@math.ac.vn

UNIVERSITÄT KAISERSLAUTERN, FACHBEREICH MATHEMATIK, ERWIN-SCHRÖDINGER-STRASSE,
67663 KAISERSLAUTERN
E-mail address: dnguyen@mathematik.uni-kl.de