

MATHEMATISCHES FORSCHUNGSINSTITUT OBERWOLFACH

Report No. 51/2015

DOI: 10.4171/OWR/2015/51

Mini-Workshop: Singular Curves on $K3$ Surfaces and Hyperkähler Manifolds

Organised by
Concettina Galati, Calabria
Andreas Leopold Knutsen, Bergen
Alessandra Sarti, Poitiers

8 November – 14 November 2015

ABSTRACT. The workshop focused on Severi varieties on $K3$ surfaces, hyperkähler manifolds and their automorphisms. The main aim was to bring researchers in deformation theory of curves and singularities together with researchers studying hyperkähler manifolds for mutual learning and interaction, and to discuss recent developments and open problems.

Mathematics Subject Classification (2010): Primary: 14H10, 14H20, 14H51, 14C20, 14J28, 14J50; Secondary: 14B07, 14E30.

Introduction by the Organisers

The workshop was attended by 15 participants with broad geographic and thematic representation. Its main aim was to bring together researchers in deformation theory of curves and singularities, especially working on Severi varieties of singular curves on $K3$ surfaces, together with researchers studying hyperkähler manifolds and their automorphisms.

Severi varieties take their name from the mathematician who introduced them at the beginning of last century. Let S be a smooth complex projective surface and $|D|$ a linear system on S containing smooth irreducible curves. The *Severi variety of δ -nodal curves* $V_{|D|,\delta}^S \subseteq |D|$ is defined as the locally closed subset of $|D|$ parametrizing irreducible curves with only δ nodes as singularities. Curves on smooth surfaces, their moduli and their enumerative geometry have been fundamental topics of algebraic geometry from the beginning of the previous century until today, thanks to the contribution of Severi, Segre, Zeuthen, Albanese, Enriques, Castelnuovo, Zariski, Arbarello, Cornalba, Harris, Shustin, Greuel and

many others. An important breakthrough was made by Harris [18], who proved that Severi varieties of nodal plane curves are irreducible, as stated by Severi. Some years later, Kontsevich and Manin [23], by using Gromov-Witten theory, computed the degree of the Severi variety of rational plane curves. Their formulas were generalized by Caporaso and Harris [10], who found a recursive formula for the degree of Severi varieties of nodal plane curves of any genus, using only classical techniques. Later on, great progress was made in the study of the enumerative geometry of $V_{|D|,\delta}^S$, by among others Pandharipande, Vakil, Ran, Göttsche, Yau, Zaslow, Vainsencher, Tzeng and Thomas. Although a lot of work has been made on Severi varieties, many interesting problems remain open, especially in the case of $K3$ surfaces, as explained in the abstracts of **Ciliberto–Flamini** and **Dedieu**.

At the same time, the Brill-Noether theory of smooth curves on $K3$ surfaces has received a lot of attention in the last couple of decades, from the seminal papers of Lazarsfeld and Green [24, 17] to the more recent works on the Green conjecture and divisors on the moduli space of curves of Voisin, Farkas, Popa and Aprodu [26, 25, 14, 1]. Very recently, two conjectures about syzygies of curves, the *Green-Lazarsfeld secant conjecture* and the *Prym-Green conjecture* were (essentially) solved by Farkas and Kemeny in [12, 13] using curves on $K3$ surfaces, and an account of this is given in **Kemeny**'s abstract. Similarly, two outstanding conjectures by Wahl were established in [2], where it is proved that a Brill-Noether-Petri curve of genus ≥ 12 lies on a polarised $K3$ surface or on a limit of such if and only if the Wahl map for C is not surjective. An account of related open problems is made in **Sernesi**'s abstract.

The recent paper [11] starts the study of Brill-Noether theory of singular curves on a $K3$ surface S . Besides its intrinsic interest, the study is related to Mori theory of hyperkähler manifolds: indeed, curves on S with normalizations carrying pencils of degree k define rational curves on the Hilbert scheme $S^{[k]}$ of k points on the surface, one of the few examples known (together with its deformations) of hyperkähler manifolds. The other known examples are Albanese fibers of Hilbert schemes of points on abelian surfaces, called *generalized Kummer varieties*, (and their deformations), as well as two examples of O'Grady in dimensions 6 and 10. We recall that a (compact) *hyperkähler manifold* (or *irreducible holomorphic symplectic manifold*) is a simply-connected compact complex Kähler manifold X such that $H^0(X, \Omega_X^2)$ is spanned by a nowhere degenerate two-form. The interest in hyperkähler manifolds stems from Bogomolov's decomposition theorem for compact, complex Kähler manifolds with trivial canonical bundle in the 70s: up to finite étale cover they all decompose into products of Calabi-Yau, hyperkähler manifolds and tori. The birational geometry of hyperkähler manifolds is determined by their rational curves; in particular, rational curves determine their nef and ample cones, just like for $K3$ s. Many years of research on this topic, passing in particular through several works and conjectures of Hassett and Tschinkel, culminated recently in the work of Bayer and Macrì [5] using *Bridgeland stability*, which determines (up to numerical computations) the extremal rays of the Mori cone of the Hilbert schemes of points on a $K3$ surface.

Despite recent advances by different methods, the study of curves on $K3$ or abelian surfaces with normalizations carrying special pencils still seems to be the most efficient way of concretely producing rational curves on hyperkähler manifolds. The results in [11] were recently extended to abelian surfaces in [21]. Some consequences of the results in [11, 21] on the birational geometry of the associated hyperkähler manifolds are obtained in [22] and the results and some open problems are given in **Knutsen's** abstract.

Many of the recent results on singular curves on $K3$ (and abelian) surfaces have been proved by degenerating the surfaces. It is therefore natural to ask whether one can find similar degenerations of hyperkähler manifolds, as is done in **Galati's** abstract, which also gives a brief account on the $K3$ case.

Another way of producing rational curves on $S^{[k]}$ is through automorphisms, as in e.g. [15]: the idea is to start with a special $K3$ surface such that $S^{[k]}$ contains a family of rational curves not present on the general projective deformation of it, use an automorphism of $S^{[k]}$ to produce another family of rational curves, and prove that the latter can be preserved under deformation. This is an interesting point of view, but one needs automorphisms of $S^{[k]}$ *not* coming from automorphisms of S , i.e. non-natural, and at the moment only one such example is known: the involution of Beauville on $S^{[2]}$ when S is a quartic. Thus one is in need of new such constructions. But the construction of new non-natural automorphisms on $S^{[k]}$ and more generally on other hyperkähler manifolds is an interesting and very active research topic on its own. The interest in automorphisms of hyperkähler manifolds has grown tremendously the last years. The foundational work on $K3$ surfaces by Nikulin, Mukai and Morrison was followed by classification results of Sarti with coauthors [3, 4, 16] and the recent work of Huybrechts [20]. Finally, the study of non-symplectic automorphisms on $K3$ surfaces has found a recent application in the study of Chow groups of $K3$ surfaces in particular in relation to the study of rational curves and the Bloch-Beilinson conjecture [19, 20]. Very little is known in higher dimensions, again there are results of Sarti, Boissière and coauthors [6, 7, 8, 9]. The abstract of **Boissière** gives an overview of results on automorphisms of special hyperkähler manifolds; more precise results and some open problems are formulated in the abstracts of **Camere** and **Cattaneo**, concerning existence of automorphisms and moduli spaces.

The abstracts of **Lehn**, **Saccà** and **Markushevich** explain other fundamental topics related to hyperkähler manifolds such as the construction of new manifolds, computation of Hodge numbers and Lagrangian fibrations. Finally, the abstract of **Ohashi** explains results on the automorphism group of Enriques surfaces and curve configurations. The study of the automorphism group of Enriques surfaces is very natural when studying automorphisms of $K3$ surfaces.

To promote interaction, the participants were asked to focus their talks on background results and open problems. Most talks were given in the first two days of the workshop to have time to discuss the proposed problems. We present the abstracts in chronological order and end with a few lines about the discussed open questions.

REFERENCES

- [1] M. Aprodu, G. Farkas, *Green's conjecture for curves on arbitrary K3 surfaces*, Compos. Math. **147** (2011), 839–851.
- [2] E. Arbarello, A. Bruno, E. Sernesi, *On two conjectures by J. Wahl*, arXiv:1507.05002.
- [3] M. Artebani, A. Sarti, *Non-symplectic automorphisms of order three on K3 surfaces*, Math. Ann. **342** (2008), 903–921.
- [4] M. Artebani, A. Sarti, S. Taki, *K3 surfaces with non-symplectic automorphisms of prime order (with an appendix by S. Kondō)*, Math. Z. **268** (2011), 507–533.
- [5] A. Bayer and E. Macrì, *MMP for moduli of sheaves on K3s via wall crossing: nef and movable cones, lagrangian fibrations*, Invent. Math. **198** (2014), no. 3, 505–590.
- [6] S. Boissière, *Automorphismes naturels de l'espace de Douady de points sur une surface*, Canadian J. Math. **64** (2012), 3–23.
- [7] S. Boissière, M. Nieper-Wisskirchen, A. Sarti, *Higher dimensional Enriques varieties and automorphisms of generalized Kummer varieties*, J. Math. Pures Appl. **95** (2011), 553–563.
- [8] S. Boissière, M. Nieper-Wisskirchen, A. Sarti, *Smith theory and irreducible holomorphic symplectic manifolds*, J. Topology **6** (2013), 361–390.
- [9] S. Boissière, A. Sarti, *A note on automorphisms and birational transformations of holomorphic symplectic manifolds*, Proc. Amer. Math. Soc. **140** (2012), 4053–4062.
- [10] L. Caporaso, J. Harris, *Counting plane curves of any genus*, Invent. Math. **131** (1998), 345–392.
- [11] C. Ciliberto, A. L. Knutsen, *On k -gonal loci in Severi varieties on general K3 surfaces and rational curves on hyperkähler manifolds*, J. Math. Pures Appl. **101** (2014), 473–494.
- [12] G. Farkas, M. Kemeny, *The generic Green–Lazarsfeld Secant conjecture*, Invent. Math. (to appear).
- [13] G. Farkas, M. Kemeny, *The Prym–Green conjecture for torsion bundles of high order*, arXiv:1509.07162.
- [14] G. Farkas, M. Popa, *Effective divisors on \mathcal{M}_g , curves on K3 surfaces and the Slope Conjecture*, J. Alg. Geom. **14** (2005), 241–267.
- [15] F. Flamini, A. L. Knutsen, G. Pacienza, *On families of rational curves in the Hilbert square of a surface (with an appendix by E. Sernesi)*, Michigan Math. J. **58** (2009), 639–682.
- [16] A. Garbagnati, A. Sarti, *Elliptic fibrations and symplectic automorphisms on K3 surfaces*, Comm. in Algebra, **37** (2009), 3601–3631.
- [17] M. Green, R. Lazarsfeld, *Special divisors on curves on a K3 surface*, Invent. Math. **89** (1987), 357–370.
- [18] J. Harris, *On the Severi problem*, Invent. Math. **84** (1986), 445–461.
- [19] D. Huybrechts, *Curves and cycles on K3 surfaces, with an appendix by Claire Voisin*, Algebraic Geometry **1** (2014), 69–106.
- [20] D. Huybrechts, *Symplectic automorphisms of K3 surfaces of arbitrary order*, Math. Res. Lett. **19** (2012), 947–951.
- [21] A. L. Knutsen, M. Lelli-Chiesa, G. Mongardi, *Severi varieties and Brill-Noether theory of curves on abelian surfaces*, arXiv:1503.04465.
- [22] A. L. Knutsen, M. Lelli-Chiesa, G. Mongardi, *Wall divisors and algebraically coisotropic subvarieties of irreducible holomorphic symplectic manifolds*, arXiv:1507.06891.
- [23] M. Kontsevich, Yu. Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Comm. Math. Phys. **164** (1994), 525–562.
- [24] R. Lazarsfeld, *Brill-Noether-Petri without degenerations*, J. Diff. Geom. **23** (1986), 299–307.
- [25] C. Voisin *Green's canonical syzygy conjecture for generic curves of odd genus*, Compos. Math. **141** (2005), 1163–1190.
- [26] C. Voisin, *Green's conjecture for curves of even genus lying on a K3 surface*, J. Eur. Math. Soc. **4**, 363–404 (2002).

Acknowledgement: The MFO and the workshop organizers would like to thank the National Science Foundation for supporting the participation of junior researchers in the workshop by the grant DMS-1049268, “US Junior Oberwolfach Fellows”.

Mini-Workshop: Singular Curves on K3 Surfaces and Hyperkähler Manifolds

Table of Contents

Samuel Boissière	
<i>Recent progress on the classification of automorphisms of hyperkähler manifolds</i>	2947
Andrea Cattaneo (joint with S. Boissière, M. Nieper-Wisskirchen and A. Sarti)	
<i>Non-symplectic involutions on the Hilbert scheme of points on a K3 surface</i>	2949
Chiara Camere (joint with S. Boissière, A. Sarti)	
<i>Complex ball quotients from four-folds of $K3^{[2]}$-type</i>	2950
Hisanori Ohashi (joint with S. Mukai)	
<i>Curve configurations on Enriques surfaces and the automorphism groups</i>	2951
Andreas Leopold Knutsen (joint with C. Ciliberto; M. Lelli-Chiesa, G. Mongardi)	
<i>Rational curves in hyperkähler manifolds</i>	2952
Ciro Ciliberto, Flaminio Flamini	
<i>Nodal curves on K3 surfaces: state of the art and open problems</i>	2953
Thomas Dedieu (joint with E. Sernesi)	
<i>The problem of the density of nodal curves in equigeneric families</i>	2958
Concettina Galati	
<i>Moduli spaces of hyperkähler manifolds and compactification problems. What do we know?</i>	2959
Edoardo Sernesi	
<i>Problems related to “fake” K3 surfaces</i>	2960
Giulia Saccà (joint with G. Mongardi, A. Rapagnetta)	
<i>Geometry of O’Grady’s 6 dimensional example</i>	2960
Michael Kemeny (joint with G. Farkas)	
<i>Syzygies of curves and K3 surfaces</i>	2961
Manfred Lehn (joint with Ch. Lehn, Ch. Sorger, D. van Straten; N. Addington; and I. Dolgachev)	
<i>Twisted cubics on a cubic fourfold and in involution on the associated 8-dimensional symplectic manifold</i>	2962

Dimitri Markushevich

On the problem of compactification of Lagrangian fibrations2964

Short report on discussion sessions2965

Abstracts

Recent progress on the classification of automorphisms of hyperkähler manifolds

SAMUEL BOISSIÈRE

The group $\text{Aut}(X)$ of biholomorphisms of an irreducible holomorphic symplectic manifold X is a discrete complex Lie group. Let us focus on finite subgroups of $\text{Aut}(X)$ where X is a $K3$ surface or a deformation of the Hilbert scheme $\text{K3}^{[n]}$ of n points on a $K3$ surface.

The symplectic automorphisms are those acting trivially on the symplectic two-form. All finite groups of symplectic automorphisms of $K3$ surfaces have been classified by Nikulin [12] and Mukai [11]. A generalisation of this classification for deformations of $\text{K3}^{[2]}$ was obtained by Mongardi [10] and Höhn–Mason [9]. Of special interest is the solution of a conjecture of Camere [7] in [10], namely that every symplectic involution can be deformed to an involution of a Hilbert scheme of two points induced by a symplectic involution of the underlying $K3$ surface.

Let now G be any finite subgroup of $\text{Aut}(X)$. Looking at the action on the symplectic form, one gets an exact sequence

$$0 \longrightarrow G_0 \longrightarrow G \longrightarrow \mathbb{Z}/m\mathbb{Z} \longrightarrow 0$$

where G_0 contains only symplectic automorphisms and m is the non-symplectic index of G . If X is non-projective then $m = 1$ (see [3]). Otherwise, a bound for m is given by $\varphi(m) \leq b_2(X) - \rho(X)$, where φ is the Euler's totient function, $b_2(X)$ is the second Betti number of X and $\rho(X) \geq 1$ is its Picard number. The situation is particularly interesting when $m = p$ is a prime number: if X is a $K3$ surface, then $p \leq 19$; but if X is a deformation of $\text{K3}^{[n]}$ one has $p \leq 23$. Boissière–Camere–Mongardi–Sarti [4] have shown the existence of a unique variety in this deformation class that contains a non-symplectic automorphism of order 23. Non-symplectic involutions have been studied and classified by Beauville [2] and Ohashi–Wandel [13].

We give the main ingredients of the classification, obtained by Boissière–Camere–Sarti [5], of non-symplectic groups G of automorphisms of prime order p such that $3 \leq p \leq 19$, acting on a deformation of $\text{K3}^{[2]}$. The classification is governed by two primitive sublattices of the second cohomology lattice $H^2(X, \mathbb{Z})$, equipped with the Beauville–Bogomolov–Fujiki quadratic form: the sublattice $T(G)$ invariant by the automorphism, and its orthogonal complement $S(G)$. In order to get a classification of the pair of lattices $(T(G), S(G))$, we use on one side deep lattice theoretical results on existence of lattices with given signature and discriminant and of embedding of lattices with given orthogonal complement, and on the other side topological information on the fixed locus X^G which is closely related to some numerical invariants of the lattices $T(G)$ and $S(G)$: the Euler characteristics of X^G is computed by applying the Lefschetz topological fixed point formula, and the sum of the dimensions of the mod p cohomology of X^G is computed by using a

formula of Boissière–Nieper-Wißkirchen–Sarti [6] originated in Smith theory and making use of the classification theorem of finite-dimensional $\mathbb{F}_p[G]$ -modules of Diederichsen and Reiner. This classification reveals two interesting features:

- contrary to the classification of non-symplectic automorphisms on $K3$ surfaces [12, 1], in our situation the pair $(T(G), S(G))$ does not uniquely determine the geometry of the fixed locus;
- contrary to the situation for symplectic involutions on deformations of $K3^{[2]}$ recalled above, it is not true that non-symplectic automorphisms of prime order $p \geq 3$ are deformations of natural automorphisms on $K3^{[2]}$.

To illustrate these features we present geometric examples using Hilbert schemes of points on $K3$ surfaces and Fano varieties of lines on cubic fourfolds, endowed with a non-symplectic automorphism of order three.

REFERENCES

- [1] M. Artebani, A. Sarti, S. Taki, *K3 surfaces with non-symplectic automorphisms of prime order (with an appendix by S. Kondō)*, *Math. Z.*, **268** (2011), 507–533.
- [2] A. Beauville, *Antisymplectic involutions of holomorphic symplectic manifolds*, *J. Topology* **4** (2011), 300–304.
- [3] A. Beauville, *Some remarks on Kähler manifolds with $c_1 = 0$* , Classification of algebraic and analytic manifolds (Katata, 1982), *Progr. Math.*, vol. 39, Birkhäuser Boston, Boston, MA, 1983, pp. 1–26.
- [4] S. Boissière, C. Camere, G. Mongardi, A. Sarti, *Isometries of ideal lattices and hyperkähler manifolds*, *Int. Math. Res. Not.* (to appear)
- [5] S. Boissière, C. Camere, A. Sarti, *Classification of automorphisms on a deformation family of hyperkähler fourfolds by p -elementary lattices*, *Kyoto J. Math.* (to appear)
- [6] S. Boissière, M. Nieper-Wißkirchen, A. Sarti, *Smith theory and irreducible holomorphic symplectic manifolds*, *J. Topology* **6** (2013), 361–390.
- [7] C. Camere, *Symplectic involutions of holomorphic symplectic fourfolds*, *Bull. Lond. Math. Soc.* **44** (2012), 687–702.
- [8] D. Huybrechts, *A global Torelli theorem for hyperkähler manifolds [after M. Verbitsky]*, *Astérisque* (2012), no. 348, Séminaire Bourbaki: Vol. 2010/2011. Exposés 1027–1042.
- [9] G. Höhn, G. Mason, *Finite groups of symplectic automorphisms of hyperkähler manifolds of type $K3^{[2]}$* , arXiv:1409.6055.
- [10] G. Mongardi, *Symplectic involutions on deformations of $K3^{[2]}$* , *Cent. Eur. J. Math.* **10** (2012), 1472–1485.
- [11] S. Mukai, *Finite groups of automorphisms of $K3$ surfaces and the Mathieu group*, *Invent. Math.* **94** (1988), 183–221.
- [12] V. V. Nikulin, *Finite groups of automorphisms of Kählerian $K3$ surfaces*, *Trudy Moskov. Mat. Obshch.* **38** (1979), 75–137.
- [13] H. Ohashi, M. Wandel, *Non-natural non-symplectic involutions on symplectic manifolds of $K3^{[2]}$ -type*, arXiv:1305.6353.

Non-symplectic involutions on the Hilbert scheme of points on a K3 surface

ANDREA CATTANEO

(joint work with S. Boissière, M. Nieper-Wisskirchen and A. Sarti)

Let S be a generic projective K3 surface, whose Picard group is generated by a divisor H with $H^2 = 2t$, and consider the Hilbert scheme $S^{[2]}$ of 2 points on S . We describe the automorphism group $\text{Aut } S^{[2]}$ giving an account of results in [1].

We prove that $\text{Aut } S^{[2]}$ has at most one non-trivial element, which is not induced by an automorphism of S (except for the case $t = 1$) and corresponds to a non-symplectic involution. We give necessary and sufficient conditions on t for the existence of such involution. These conditions are related to the solutions of certain Pell's equations, and have a nice translation in the geometry of $S^{[2]}$: they reflect the presence of an ample class on $S^{[2]}$ of Beauville–Bogomolov degree 2.

Since any automorphism of $S^{[2]}$ induces an isometry of the Néron–Severi lattice $\text{NS}(S^{[2]})$ of $S^{[2]}$, the first step is to analyse the orthogonal group $O(\text{NS}(S^{[2]}))$ and then the homomorphism $\Psi : \text{Aut } S^{[2]} \rightarrow O(\text{NS}(S^{[2]}))$ associating to each automorphism its isometry. We show that Ψ is injective.

The main step in the study of $\text{Aut } S^{[2]}$ is then to relate the isometries in $O(\text{NS}(S^{[2]}))$ with the geometry of $S^{[2]}$, and in particular we show that the presence of non-trivial automorphisms depends on the generators of the ample cone of $S^{[2]}$.

We show under which conditions it is possible to extend an isometry of $\text{NS}(S^{[2]})$ to a Hodge isometry of $H^2(S^{[2]}, \mathbb{Z})$ and finally, using Hodge-theoretic Torelli theorems, how to understand if such extensions are induced by automorphisms.

The first case of a surface S such that $\text{Aut}(S^{[2]}) \simeq \mathbb{Z}/2\mathbb{Z}$ is for $H^2 = 4$ (i.e. $t = 2$), and was geometrically described by Beauville. Here S is a quartic surface in \mathbb{P}^3 , two points on S define a line that cuts S in two other points, and this sets up the non-symplectic involution on $S^{[2]}$.

The next case happens for $H^2 = 20$ (i.e. $t = 10$).

Question 1. *How can one find a geometric description of the corresponding involution on $S^{[2]}$?*

Up to now there is no such description, and there are several reasons to search for such a description. First of all because it will be the first *explicit* description after Beauville of an involution on $S^{[2]}$ and not on some of its deformations. Then because it provides a link with many other objects of their own interest: Grassmannians, the secant variety of S , moduli spaces of sheaves on S and on curves (in the linear system $|H|$) of genus eleven (this last link provided in [2]).

REFERENCES

- [1] S. Boissière, A. Cattaneo, M. Nieper-Wisskirchen, A. Sarti, *The automorphism group of the Hilbert scheme of two points on a generic projective K3 surface*, Proceedings of the Schiermonnikoog conference “K3 surfaces and their moduli”, 2015 (to appear).
- [2] S. Mukai, *Curves and K3 surfaces of genus eleven*, in Moduli of vector bundles (Sanda, 1994; Kyoto, 1994), Lecture Notes in Pure and Appl. Math. **179** (1996), 189–197.

Complex ball quotients from four-folds of $K3^{[2]}$ -type

CHIARA CAMERE

(joint work with S. Boissière, A. Sarti)

In [4] we study complex ball quotients arising from four-folds of $K3^{[2]}$ -type endowed with a non-symplectic automorphism of prime order p with given invariant sublattice T . Such automorphisms have order $3 \leq p \leq 23$ and have been completely classified in [3], [2] and [8].

Let (X, ι) be a T -polarized IHS manifold and $G = \langle \bar{\sigma} \rangle$ a cyclic group of prime order $p \geq 3$ acting non-symplectically on X . Let $\rho : G \rightarrow O(L)$ be a group homomorphism such that $T = L^\rho := \{x \in L \mid \rho(g)(x) = x, \forall g \in G\}$. We define a (ρ, T) -polarization of (X, ι) as a marking $\eta : L \rightarrow H^2(X, \mathbb{Z})$ such that $\eta|_T = \iota$ and $\bar{\sigma}^* = \eta \circ \rho(\bar{\sigma}) \circ \eta^{-1}$. Let $S(\xi)$ be the eigenspace relative to a primitive p -th root of the unity ξ inside $T^\perp \otimes \mathbb{C}$; the period of (X, η) belongs to the space $\Omega_T^{\rho, \xi} := \{x \in \mathbb{P}(S(\xi)) \mid q(x + \bar{x}) > 0\}$, which is a complex ball of dimension $\dim S(\xi) - 1$ if $\dim S(\xi) \geq 2$.

Amerik–Verbitsky’s [1] theory of algebraic MBM classes and their description of the Kähler cone \mathcal{K}_X of an IHS manifold X allow us to study the image and the locus of injectivity of the period map. Let $\Delta(L)$ be the set of $\delta \in L$ such that there exists a marked pair (X, ϕ) for which $\phi(\delta) \in \text{NS}(X)$ is MBM.

Theorem 2. *The period of a (ρ, T) -polarized pair (X, ϕ) belongs to $\Omega_T^{\rho, \xi} \setminus \Delta$, where*

$$\Delta := \bigcup_{\delta \in \Delta(L) \cap T^\perp} (H_\delta \cap \Omega_T^{\rho, \xi}).$$

Given a chamber $K(T)$ of the decomposition of the positive cone of $T \otimes \mathbb{R}$ given by walls in $\Delta(L) \cap T$, a (ρ, T) -polarized pair (X, ϕ) is $K(T)$ -general if $\phi(K(T)) = \mathcal{K}_X \cap (\phi(T) \otimes \mathbb{R})$. Let $\mathcal{M}_{K(T)}^{\rho, \xi}$ be the set of $K(T)$ -general (ρ, T) -polarized pairs.

Theorem 3. *The period map gives a bijection $\mathcal{P} : \mathcal{M}_{K(T)}^{\rho, \xi} \rightarrow \Omega := \Omega_T^{\rho, \xi} \setminus (\Delta \cup \Delta')$, where Δ' is a locally finite union of hyperplanes. Moreover, there exist arithmetic subgroups \mathcal{G} and Γ of $O(L)$ and $O(T^\perp)$ such that the complex ball quotient $\mathcal{M}_{K(T)}^{\rho, \xi} / \mathcal{G} \cong \Omega / \Gamma$ is a quasi-projective variety of dimension $\dim S(\xi) - 1$.*

Once the construction of an algebraic moduli space is established, one of the first natural questions that one can ask is:

Question 4. *What is its Kodaira dimension? In which cases is it rational?*

In the case of complex ball quotients arising from $K3$ surfaces, the pioneering works [6, 5] started the study of the ball quotients corresponding to some non-symplectic automorphisms of order three; later, [7] established rationality for many cases of order three. For natural automorphisms of order three corresponding to the cases studied in the cited papers, the question of rationality can hopefully be answered, since the complex ball quotients studied are the same. For higher

orders, rationality seems to be an open problem also in the case of the underlying K3 surfaces.

Some of the known non-natural automorphisms of order three appear on Fano varieties of special cubic four-folds. The question is how one can understand whether the corresponding ball quotients are rational.

REFERENCES

- [1] E. Amerik, M. Verbitsky, *Rational curves on hyperkähler manifolds*, Int. Math. Res. Notices (2015), doi: 10.1093/imrn/rnv133.
- [2] S. Boissière, C. Camere, G. Mongardi, A. Sarti, *Isometries of ideal lattices and hyperkähler manifolds*, Int. Math. Res. Not. (2015), doi: 10.1093/imrn/rnv137.
- [3] S. Boissière, C. Camere, A. Sarti, *Classification of automorphisms on a deformation family of hyperkähler fourfolds by p -elementary lattices*, Kyoto J. Math. (to appear).
- [4] S. Boissière, C. Camere, A. Sarti, *Complex ball quotients from four-folds of $K3^{[2]}$ -type*, arXiv:1512.02067.
- [5] I. Dolgachev, B. van Geemen, S. Kondō, *A complex ball uniformization of the moduli space of cubic surfaces via periods of K3 surfaces*, J. Reine Angew. Math. **588** (2005), 99–148.
- [6] S. Kondō, *The moduli space of curves of genus 4 and Deligne–Mostow’s complex reflection groups*, Algebraic geometry 2000, Azumino (Hotaka), Adv. Stud. Pure Math., vol. 36, Math. Soc. Japan, Tokyo, 2002, pp. 383–400.
- [7] S. Ma, H. Ohashi, S. Taki, *Rationality of the moduli spaces of Eisenstein K3 surfaces*, Trans. Amer. Math. Soc. **367** (2015), 8643–8679.
- [8] K. Tari, *Automorphismes des variétés de Kummer généralisées*, Ph.D.-thesis, Université de Poitiers, 2015.

Curve configurations on Enriques surfaces and the automorphism groups

HISANORI OHASHI

(joint work with S. Mukai)

For K3 and Enriques surfaces, curve configurations play a central role in studying their automorphism groups since the configurations determine a reflection group acting on the cohomology lattice that is a kind of complement to the automorphism group in the total orthogonal group of the hyperbolic lattice. They are described by the dual graph of (-2) -curves. Vinberg [5] developed an effective procedure to measure the size of the reflection group and Nikulin [2, 3, 4] and Kondō [1] classified K3 and Enriques surfaces with finite automorphism groups in detail.

We give a further example of Enriques surfaces with a good (-2) -configuration, which is used to determine their infinite automorphism groups precisely by the explicit generators and relations. The new feature here is that the configuration consists of classes of (-2) -curves and centers of numerically reflective involutions.

The resulting dual graph coincides with that of type V surfaces of Kondō [1].

Question 5. *Is there a method to classify the abstract good dual graphs in the Enriques lattice? Can we find surfaces with the (-2) -configurations?*

REFERENCES

- [1] S. Kondō, *Enriques surfaces with finite automorphism groups*, Japan. J. Math. **12** (1986), 191–282.
- [2] V. V. Nikulin, *Factor groups of groups of automorphisms of hyperbolic forms with respect to subgroups generated by 2-reflections (English)*, J. Soviet Math. **22** (1983), 1401–1476.
- [3] V. V. Nikulin, *On a description of the automorphism groups of Enriques surfaces*, Soviet Math. Dokl. **30** (1984), 282–285.
- [4] V. V. Nikulin, *Surfaces of type K3 with a finite automorphism group and a Picard group of rank three*, Proceedings of the Steklov Institute of Mathematics **3** (1985), 131–155.
- [5] E. B. Vinberg, *Some arithmetical discrete groups in Lobachevskii spaces*, in Discrete Subgroups of Lie Groups and Appl. to Moduli (Bombay 1973), Oxford Univ. Press, 1975, pp. 323–348.

Rational curves in hyperkähler manifolds

ANDREAS LEOPOLD KNUTSEN

(joint work with C. Ciliberto; M. Lelli-Chiesa, G. Mongardi)

Following notation in [4], we let $\varepsilon = 0$ (resp., $\varepsilon = 1$) when S is a K3 (resp., abelian) surface, and we denote by $S_\varepsilon^{[k]}$ the Hilbert scheme of k points on S when $\varepsilon = 0$ and the $2k$ -dimensional generalised Kummer variety on S when $\varepsilon = 1$. It is well-known that there is a canonical decomposition $N_1(S_\varepsilon^{[k]}) \cong N_1(S) \oplus \mathbb{Z}[\mathfrak{r}_k]$, where \mathfrak{r}_k is the class of a general rational curve contracted by the Hilbert-Chow morphism $S_\varepsilon^{[k]} \rightarrow \text{Sym}^{k+\varepsilon}(S)$, and the embedding $N_1(S) \subset N_1(S_\varepsilon^{[k]})$ is given by mapping an irreducible curve $C \subset S$ to the class of the curve $\{x \cup x_1 \cup \dots \cup x_{k+\varepsilon-1} \mid x \in C\}$ for fixed points $x_1, \dots, x_{k+\varepsilon-1} \in S \setminus C$. Rational curves in $S_\varepsilon^{[k]}$ correspond to (possibly reducible) curves on S with a partial normalization admitting a linear system of type $g_{k+\varepsilon}^1$. The main results concerning irreducible such curves and the rational curves they define in $S_\varepsilon^{[k]}$ are given in [2, 3]:

Theorem 6. *Let (S, L) be a general polarized K3 or abelian surface of genus $p := p_a(L)$. Let g and k be integers satisfying $2\varepsilon \leq g \leq p$ and $k + \varepsilon \geq 2$. Then the locus of irreducible curves of genus g of class $[L]$ carrying a $g_{k+\varepsilon}^1$ is nonempty if and only if $\rho(p, \alpha, (k + \varepsilon)\alpha + p - g) + \varepsilon\alpha(\alpha + 2) \geq 0$, where $\alpha = \left\lfloor \frac{g-\varepsilon}{2(k-1+2\varepsilon)} \right\rfloor$.*

Furthermore, whenever nonempty, this locus is equidimensional of dimension $\min\{g, 2(k-1+\varepsilon)\}$ and the normalization of a general element in each component carries a $\max\{0, \rho(g, 1, k + \varepsilon)\}$ -dimensional family of $g_{k+\varepsilon}^1$ s.

Theorem 7. *The family of rational curves in $S_\varepsilon^{[k]}$ obtained from curves on S with normalizations carrying a $g_{k+\varepsilon}^1$ is precisely $2(k-1)$ -dimensional and there is at least one irreducible component yielding rational curves of class $L - (g+k-1+\varepsilon)\mathfrak{r}_k$.*

We note that $2(k-1)$ is the expected dimension of any family of rational curves in a $2k$ -dimensional hyperkähler manifold, by a result of Ran [5]. As an interesting consequence, the family is preserved under any small deformation X_t of $X_0 = S_\varepsilon^{[k]}$ keeping their class algebraic. Another interesting consequence is, using a result of

Amerik and Verbitsky [1], that any irreducible components of the locus $W \subset S_\varepsilon^{[k]}$ of the family satisfies the property that the dimension of the fibers of the rational quotient of its desingularization equals the codimension of W in $S_\varepsilon^{[k]}$, and as such, is an *algebraically coisotropic subvariety*, cf. [6]. Also in view of recent conjectures of Voisin in [6], some interesting open questions are:

Question 8. *How can one compute $\dim W$ (in terms of properties of (S, L))?*

This question is answered in some particular cases in [4].

Question 9. *When deforming $X_0 = S_\varepsilon^{[k]}$ keeping the class of the rational curves algebraic, does (some component of) the locus they cover keep its dimension?*

REFERENCES

- [1] E. Amerik, M. Verbitsky, *Rational curves on hyperkähler manifolds*, Int. Math. Res. Notices (2015), doi: 10.1093/imrn/rnv133.
- [2] C. Ciliberto, A. L. Knutsen, *On k -gonal loci in Severi varieties on general K3 surfaces and rational curves on hyperkähler manifolds*, J. Math. Pures Appl. **101** (2014), 473–494.
- [3] A. L. Knutsen, M. Lelli-Chiesa, G. Mongardi, *Severi varieties and Brill-Noether theory of curves on abelian surfaces*, arXiv:1503.04465.
- [4] A. L. Knutsen, M. Lelli-Chiesa, G. Mongardi, *Wall divisors and algebraically coisotropic subvarieties of irreducible holomorphic symplectic manifolds*, arXiv:1507.06891.
- [5] Z. Ran, *Hodge theory and deformations of maps*, Compos. Math. **97** (1995), 309–328.
- [6] C. Voisin, *Remarks and questions on coisotropic subvarieties and 0-cycles of hyper-Kähler varieties*, arXiv:1501.02984.

Nodal curves on K3 surfaces: state of the art and open problems

CIRO CILIBERTO, FLAMINIO FLAMINI

The following questions came up from discussions with T. Dedieu, C. Galati, A. L. Knutsen and E. Sernesi.

Let \mathcal{K}_p be the moduli space of primitively polarised K3 surfaces (X, L) , with $L^2 = 2p - 2 > 0$ and let $\mathcal{V}_{p,n,\delta}$ be the (p, n, δ) -universal Severi variety parametrizing all triples (X, L, C) with $(X, L) \in \mathcal{K}_p$ and $C \in |nL|$ (with $n \geq 1$) irreducible and with only δ nodes as singularities, with

$$0 \leq \delta \leq \pi_a(p, n) := n^2(p - 1) + 1.$$

Consider the projection

$$\phi_{p,n,\delta} : \mathcal{V}_{p,n,\delta} \rightarrow \mathcal{K}_p$$

whose fiber over any $(X, L) \in \mathcal{K}_p$ is denoted by $V_{n,\delta}(X)$: if nonempty, this is the *Severi variety* of δ -nodal irreducible curves in $|nL|$ on X . The variety $V_{n,\delta}(X)$, if nonempty, is well-known to be smooth, pure of dimension $g_{p,n,\delta} := \pi_a(p, n) - \delta = n^2(p - 1) + 1 - \delta$, and this is the case if $(X, L) \in \mathcal{K}_p$ is general, so also $\mathcal{V}_{p,n,\delta}$ is smooth, pure and each irreducible components of it dominates \mathcal{K}_p . We often write g for $g_{p,n,\delta}$, which is the *geometric genus* of curves in $\mathcal{V}_{p,n,\delta}$.

Question 10. *Is $\mathcal{V}_{p,n,\delta}$ irreducible?*

Question 11. Let $(X, L) \in \mathcal{K}_p$ be general and $0 < \delta < \pi_a(p, n)$. Is $V_{n,\delta}(X)$ irreducible?

Question 12. Construct examples, if any, of $(X, L) \in \mathcal{K}_p$ and $\delta < \pi_a(p, n)$ such that $V_{n,\delta}(X)$ is empty or reducible.

Of course an affirmative answer to Question 11 implies an affirmative answer to Question 10 for $\delta < \pi_a(p, n)$. The state of the art, cf. [7], is that Question 10 has an affirmative answer for $n = 1$, $3 \leq p \leq 9$ or $p = 11$ and any (admissible) δ .

The following particular case of Questions 10 and 11 should be easier:

Question 13. Let $(X, L) \in \mathcal{K}_p$ be general and $\delta \leq \frac{\pi_a(p,n)}{3}$. Is $V_{n,\delta}(X)$ irreducible?

Partial results in this direction can be found in [18, 19].

Question 14. Let $(X, L) \in \mathcal{K}_p$ be general and $\delta \leq \pi_a(p, n)$. Compute the Hilbert polynomial of the closure of $V_{n,\delta}(X)$ in $|nL|$.

State of the art: the degree of $V_{1,\delta}(X)$ is known [5] and the *putative* degree of the 0-dimensional $V_{n,\pi_a(p,n)}(X)$ (i.e., the number of rational curves in $|nL|$) is known for any $n \geq 1$ (this is the *Yau–Zaslow formula* [3]). The case of $V_{n,1}(X)$, i.e., the *dual variety* of X embedded in projective space via $|nL|$, is trivial.

Next we want to propose questions concerning the *moduli map*

$$\psi_{p,n,\delta} : \mathcal{V}_{p,n,\delta} \longrightarrow \mathcal{M}_g,$$

where \mathcal{M}_g is the moduli space of smooth genus- g curves, sending a curve C to the class of its normalization. Of course, one can consider its restriction

$$\psi_{X,n,\delta} : V_{n,\delta}(X) \longrightarrow \mathcal{M}_g.$$

Naive dimension counts:

- ensure that no dominance is allowed for $\psi_{X,n,\delta}$, whereas
- suggest that $\psi_{p,n,\delta}$ should be

$$(1) \quad \begin{array}{ll} \text{dominant if} & g \leq 11, \\ \text{generically finite onto its image if} & g \geq 11. \end{array}$$

Question 15. Let $(X, L) \in \mathcal{K}_p$ be general and $\delta < \pi_a(p, n)$. Let V be an irreducible component of $V_{n,\delta}(X)$. Is $\psi_{X,n,\delta|V}$ generically finite or even birational (if $g > 1$) to its image?

State of the art: Let $(X, L) \in \mathcal{K}_p$. For any component $V \subseteq V_{n,\delta}(X)$ containing irreducible rational nodal curves in its Zariski closure, the restriction $\psi_{X,n,\delta|V}$ is generically finite [11]. In particular, for a general $(X, L) \in \mathcal{K}_p$, the moduli map is generically finite on at least one component of the Severi variety $V_{n,\delta}(X)$. The last statement follows from the fact that $V_{n,\pi_a(p,n)}(X)$ is nonempty [6]. In [17] one proves that if $\text{Pic}(X) = \mathbb{Z}\langle L \rangle$ and $C \in V_{n,\delta}(X)$ is such that either

$$n = 1, 2 \quad \text{and} \quad \delta \leq \frac{\pi_a(p,n)}{2} - 25, \quad \text{or} \quad n \geq 3 \quad \text{and} \quad \delta \leq 2(n-1)(p-1) - 25,$$

then the pull back of T_X to C' is stable, where C' is the normalization of C . This implies that $\psi_{X,n,\delta|V}$ generically finite onto its image in these cases.

Question 16. *Let \mathcal{V} be an irreducible component of $\mathcal{V}_{p,n,\delta}$ [resp. V be an irreducible component of $V_{n,\delta}(X)$ where $(X, L) \in \mathcal{K}_p$ is general]. Does the differential of $\psi_{p,n,\delta|_{\mathcal{V}}}$ [resp. $\psi_{X,n,\delta|_V}$] have maximal rank?*

State of the art: Results have been obtained recently by Kemeny [20] and in [8]. We quote the latter result, which is optimal for $n \geq 5$, but noting that Kemeny's results are stronger in case (B) for $n \leq 4$:

Theorem 17. (A) *For the following values of $p \geq 3$, n and $g = \pi_a(p, n) - \delta$ there is an irreducible component \mathcal{V} of $\mathcal{V}_{p,n,\delta}$, such that the moduli map $\mathcal{V} \rightarrow \overline{\mathcal{M}}_g$ is dominant:*

- $n = 1$ and $0 \leq g \leq 7$;
- $n = 2$, $p \geq g - 1$ and $0 \leq g \leq 8$;
- $n = 3$, $p \geq g - 2$ and $0 \leq g \leq 9$;
- $n = 4$, $p \geq g - 3$ and $0 \leq g \leq 10$;
- $n \geq 5$, $p \geq g - 4$ and $0 \leq g \leq 11$.

(B) *For the following values of p , n and $g = \pi_a(p, n) - \delta$ there is an irreducible component \mathcal{V} of $\mathcal{V}_{p,n,\delta}$, such that the moduli map $\mathcal{V} \rightarrow \overline{\mathcal{M}}_g$ is generically finite onto its image:*

- $n = 1$ and $p \geq g \geq 15$;
- $2 \leq n \leq 4$, $p \geq 15$ and $g \geq 16$;
- $n \geq 5$, $p \geq 7$ and $g \geq 11$.

Note that for $3 \leq p \leq 11$, $n = 1$ and $0 \leq \delta \leq p - 2$, the moduli map $\psi_{p,n,\delta}$ is dominant on any component $\mathcal{V} \subseteq \mathcal{V}_{p,n,\delta}$ by [16]. From [21, 22], the map $\psi_{p,1,0}$ is dominant for $p \leq 9$ and $p = 11$ but not for $p = 10$, the latter against expectation (1). Indeed, [13] proved that $\mathcal{S}_{10} := \text{Im}(\psi_{10,1,0})$ is a divisor in \mathcal{M}_{10} consisting of those curves of genus 10 lying in the primitive linear system of surfaces in \mathcal{K}_{10} : the exceptional behaviour is due to the fact that the general K3 surface of genus 10 is a codimension-3 linear section of a suitable 5-fold and the existence of Fano threefolds of that genus, as Mukai showed. Then [25, 14] gave another interesting realization of the divisor \mathcal{S}_{10} : it is the divisorial component of the locus in \mathcal{M}_{10} of curves for which $\text{SU}_C(2, K_C, 7) \neq \emptyset$, i.e. carrying a semistable, rank-two vector bundle with canonical determinant and at least 7 independent global sections.

On the other hand, in [23] it is proved that $\psi_{p,1,0}$ is generically finite for $p = 11$ and $p \geq 13$ but not for $p = 12$ (once again the latter against expectation (1)). More precisely, [12] proved that $\psi_{p,1,0}$ is birational for $p = 11$ and $p \geq 13$; then [24], for $p = 11$, and more recently [1], for any $p > 11$ congruent to 3 (mod 4), explicitly construct a rational inverse of $\psi_{p,1,0}$.

Question 18. *In the cases not covered by Theorem 17 (or Kemeny's result [20]), is it possible to find cases in which expectation (1) is contradicted, as happens for $\mathcal{V}_{p,1,0}$ for $p = 10$ and $p = 12$? If yes, what kind of geometric reasons are behind this fact? Is there any chance to get a divisorial image for $g \leq 11$?*

Question 19. *Is there any chance to describe some of the images $\text{Im}(\psi_{p,n,\delta})$ via higher-rank Brill-Noether theory as in [25, 14]?*

Question 20. Let \mathcal{V} be an irreducible component of $\mathcal{V}_{p,n,\delta}$ [resp. V be an irreducible component of $V_{n,\delta}(X)$ where $(X, L) \in \mathcal{K}_p$ is general] such that the differential $\psi_{p,n,\delta|_{\mathcal{V}}}$ [resp. $\psi_{X,n,\delta|_V}$] is of maximal rank. Is then $\psi_{p,n,\delta|_{\mathcal{V}}}$ [resp. $\psi_{X,n,\delta|_V}$] birational onto its image? What is the Kodaira dimension of this image? If $\psi_{p,n,\delta|_{\mathcal{V}}}$ is birational onto its image, is it possible to construct a rational inverse, as in [24, 1] (Torelli type theorem)?

As for the very last question, we would expect that, except maybe for very few cases that should be possible to classify, the image is of general type as soon as $3\delta > \pi(p, n)$. One may also consider the intersection of the images of the maps with Brill-Noether loci $\mathcal{M}_{g,d}^r$ in the moduli space \mathcal{M}_g of curves:

Question 21. What can one say about $\text{Im}(\psi_{X,n,\delta}) \cap \mathcal{M}_{g,d}^r$ and $\text{Im}(\psi_{p,n,\delta}) \cap \mathcal{M}_{g,d}^r$? In the first case, which consequences can one draw on the Mori theory of the Hilbert scheme of points $X^{[d]}$?

State of the art: For $\delta > 0$ and $n = r = 1$ rather complete results are found in [10] (extending preliminary results in [15]), see the abstract of Knutsen.

For a smooth, irreducible curve C one has the *Wahl map*

$$w_C : \wedge^2 H^0(C, \omega_C) \rightarrow H^0(C, \omega_C^{\otimes 3}).$$

If C sits on a $K3$ surface, then w_C is not surjective (see [4, 26]). The corank of w_C for C general in the image of $\psi_{p,n,0}$ has been studied in [12] and subsequent papers on the subject. Moreover, if C has general moduli of genus at least 11, but different from 12, then w_C is surjective (see [9]).

Wahl formulated in [27] the conjecture that if C is a smooth curve of genus $p \geq 8$ which is Brill–Noether general, then C is in the image of $\psi_{p,1,0}$ if and only if w_C is not surjective. This had been proved for $p = 10$ in [13]. In the recent paper [2] Wahl’s conjecture has been *essentially* proved, i.e.: if C is a smooth irreducible curve of genus $p \geq 19$ which is Brill–Noether–Petri general, then $C \in |L|$, with $(X, L) \in \mathcal{K}_p$ (i.e., C is in the image of $\psi_{p,1,0}$), or is a limit of such a curve (also see the abstract of Sernesi).

For nodal curves C one can consider suitable modifications of the Wahl map. Two different versions are given by Kemeny [20] and Halic [17]. We call them the *Kemeny–Wahl map* kw_C and the *Halic–Wahl map* hw_C . For both maps, what is *essentially* true is that, like the ordinary Wahl map, they are not surjective for curves on $K3$ surfaces but are surjective for a *general nodal curve*, cf. [20, 17].

Question 22. Compare kw_C and hw_C .

Question 23. Let C be general in a component of $\mathcal{V}_{p,n,\delta}$. What are the coranks of kw_C and hw_C ?

Question 24. Does it make sense to extend Wahl’s conjecture for nodal curves C using kw_C and/or hw_C , and to try to prove the analogue of Arbarello–Bruno–Sernesi’s theorem [2] in such a setting?

REFERENCES

- [1] E. Arbarello, A. Bruno, E. Sernesi, *Mukai's program for curves on a K3 surface*, Alg. Geom. **1** (2014), 532–557.
- [2] E. Arbarello, A. Bruno, E. Sernesi, *On two conjectures of J. Wahl*, arXiv:1507.05002.
- [3] A. Beauville, *Counting rational curves on K3 surfaces*, Duke Math. J. **97** (1999), 99–108.
- [4] A. Beauville, J.-Y. Merindol, *Sections hyperplanes des surfaces K3*, Duke Math. J. **55** (1987), 873–878.
- [5] J. Bryan, N. C. Leung, *The enumerative geometry of K3 surfaces and modular forms*, J. Amer. Math. Soc. **13** (2000), 371–410.
- [6] X. Chen, *Rational curves on K3 surfaces*, J. Alg. Geom. **8** (1999), 245–278.
- [7] C. Ciliberto, T. Dedieu, *On universal Severi varieties of low genus K3 surfaces*, Math. Z. **271** (2012), 953–960.
- [8] C. Ciliberto, F. Flamini, C. Galati, A.L. Knutsen, *Moduli of nodal curves on K3 surfaces*, arXiv:1502.07378.
- [9] C. Ciliberto, J. Harris, R. Miranda, *On the surjectivity of the Wahl map*, Duke Math. J. **57** (1988), 829–858.
- [10] C. Ciliberto, A. L. Knutsen, *On k -gonal loci in Severi varieties on general K3 surfaces and rational curves on hyperkähler manifolds*, J. Math. Pures Appl. **101** (2014), 473–494.
- [11] C. Ciliberto, A. L. Knutsen, *Errata corrige to [10]*, to appear.
- [12] C. Ciliberto, A. F. Lopez, R. Miranda, *Projective degenerations of K3 surfaces, Gaussian maps and Fano threefolds*, Invent. Math. **114** (1993), 641–667.
- [13] F. Cukierman, D. Ulmer, *Curves of genus 10 on K3 surfaces*, Comp. Math. **89** (1993), 81–90.
- [14] G. Farkas, M. Popa, *Effective divisors on $\overline{\mathcal{M}}_g$, curves on K3 surfaces and the slope conjecture*, J. Alg. Geom. **14** (2005).
- [15] F. Flamini, A. L. Knutsen, G. Pacienza, *On families of rational curves in the Hilbert square of a surface (with an appendix by Edoardo Sernesi)*, Michigan Math. J. **58** (2009), 639–682.
- [16] F. Flamini, A. L. Knutsen, G. Pacienza, E. Sernesi, *Nodal curves with general moduli on K3 surfaces*, Comm. Alg. **36** (2008), 3955–3971.
- [17] M. Halic, *Modular properties of nodal curves on K3 surfaces*, Math. Z. **270** (2012), 871–887; Erratum: **280** (2015), 1203–1211.
- [18] T. Keilen, *Irreducibility of equisingular families of curves*, Trans. Amer. math. Soc. **335** (2003), 3485–3512.
- [19] M. Kemeny, *The universal Severi variety of rational curves on K Surfaces*, Bull. London Math. Soc. **45** (2013), 159–174.
- [20] M. Kemeny, *The moduli of singular curves on K3 surfaces*, J. Math. Pures. Appl. **104** (2015), 882–920.
- [21] S. Mori, S. Mukai, *The uniruledness of the moduli space of curves of genus 11*, Algebraic Geometry, Proc. Tokyo/Kyoto, 334–353, Lecture Notes in Math. **1016**, Springer, Berlin, 1983.
- [22] S. Mukai, *Curves, K3 surfaces and Fano 3-folds of genus ≤ 10* . Algebraic geometry and commutative algebra, Vol. I, 357–377, Kinokuniya, Tokyo (1988).
- [23] S. Mukai, *Fano 3-folds*, in “Complex Projective Geometry”, LMS Lecture Notes Series, **179** (1992), 255–261.
- [24] S. Mukai, *Curves and K3 surfaces of genus eleven*, in “Moduli of vector bundles” (Sanda 1994–Kyoto 1994), Lecture Notes in Math. **179** (1996), 189–197.
- [25] C. Voisin, *Sur l'application de Wahl des courbes satisfaisant la condition de Brill-Noether-Petri*, Acta Math. **168** (1992), 249–272.
- [26] J. Wahl, *The Jacobian algebra of a graded Gorenstein singularity*, Duke Math. J. **55** (1987), 843–871.
- [27] J. Wahl, *On the cohomology of the square of an ideal sheaf*, J. Algebraic Geom. **6** (1997), 481–511.

The problem of the density of nodal curves in equigeneric families

THOMAS DEDIEU

(joint work with E. Sernesi)

Let S be a smooth projective surface, ξ a class in its Néron–Severi group, and g an integer. We consider V_g^ξ the locally closed subset of the Hilbert scheme of curves on S parametrizing reduced curves of genus g in the class ξ . The question is whether the subset of V_g^ξ corresponding to nodal curves (i.e. curves with only ordinary double points as possible singularities) is dense in V_g^ξ . The answer is known to be ‘yes’ when S is the projective plane [2, 1, 7], or a Del Pezzo or Hirzebruch surface [4], see also [3]. For $K3$ and abelian surfaces, only partial results are known [3, 5]; in particular, the general element of V_g^ξ is a curve with immersed singularities when $g > 0$ and $g > 2$ respectively.

Most of these results are obtained by a careful local study of the space $M_g^{\xi, \text{bir}}(S)$ of birational morphisms from smooth genus g curves to S with image in the class ξ , taking advantage of the fact that the deformation theory of such objects is well understood, and of a key observation [1] relating the geometry of $M_g^{\xi, \text{bir}}(S)$ to that of V_g^ξ (see [3, Lem. 2.5]). This method gives optimal results to prove that the general member of V_g^ξ is immersed, but is unfortunately artificially too expensive in terms of positivity of the canonical class K_S to answer the question in general.

One should therefore approach the question directly by studying deformations of integral curves $C \subset S$. Those preserving the geometric genus (resp. the topological type of the singularities) of C are governed by $N_{C/S} \otimes A$ (resp. $N_{C/S} \otimes I$), where A is the adjoint ideal of C and I is the equisingular ideal introduced by J. Wahl [6]. Following [3], we propose:

Question 25. *Use the fact that $I \not\subset A$ if and only if C has singularities worse than nodes to answer the question of the density of nodal curves.*

We also discuss implications of this question in the enumerative geometry of curves on surfaces.

REFERENCES

- [1] E. Arbarello, M. Cornalba, *Su una congettura di Petri*, Comment. Math. Helv. **56** (1981), 1–38.
- [2] E. Arbarello, M. Cornalba, *Su una proprietà notevole dei morfismi di una curva a moduli generali in uno spazio proiettivo*, Rend. Sem. Mat. Univ. Politec. Torino **38** (1980), 87–99 (1981).
- [3] T. Dedieu, E. Sernesi, *Equigeneric and equisingular families of curves on surfaces*, Publ. Mat. (to appear).
- [4] J. Harris, *On the Severi problem*, Invent. Math. **84** (1986), 445–461.
- [5] A. L. Knutsen, M. Lelli-Chiesa, G. Mongardi, *Severi varieties and Brill–Noether theory of curves on abelian surfaces*, arXiv:1503.04465.
- [6] J. Wahl, *Equisingular deformations of plane algebroid curves*, Trans. Amer. Math. Soc. **193** (1974), 143–170.
- [7] O. Zariski, *Dimension-theoretic characterization of maximal irreducible algebraic systems of plane nodal curves of a given order n and with a given number d of nodes*, Amer. J. Math. **104** (1982), 209–226.

Moduli spaces of hyperkähler manifolds and compactification problems. What do we know?

CONCETTINA GALATI

The following problem came up from discussions with C. Ciliberto, F. Flamini and A. L. Knutsen.

Besides its intrinsic interest, deformation theory of singular varieties has wide application in algebraic geometry. In particular, several recent papers about Severi varieties on $K3$ surfaces are based on degeneration arguments of curves and surfaces (cf., e.g., [1, 5, 7, 8, 9]). Semi-stable degenerations of $K3$ surfaces have been classified by Kulikov and Persson-Pinkham [6, 10] and are of three types. By [4, 3], we also know that there exists a *partial* compactification $\overline{\mathcal{K}}_p$ of the moduli space \mathcal{K}_p of polarized $K3$ surfaces of genus p , such that its boundary is a smooth divisor whose points correspond to "stable" Type II degenerations (see [3, (4.9) and (4.10)] and [2, Sect. 3] for details). The existence of this partially compactified moduli space $\overline{\mathcal{K}}_p$ is a key ingredient in [2] and is, more in general, useful for the study of the universal Severi varieties on $K3$ surfaces. Finally, [4, Thm. 5.10] provides a really interesting description of the versal deformation space of a degenerate $K3$ surface of type II and more generally of a "d-semi-stable" $K3$ surface.

Question 26. *What happens for hyperkähler manifolds of higher dimension? Is it possible to classify their degenerations? How many examples of degenerate hyperkähler manifolds are known and what do we know about their versal deformation space? Finally, in which cases is it possible to provide (possible partial) compactifications of moduli spaces of hyperkähler manifolds?*

REFERENCES

- [1] X. Chen, *Rational curves on K3 surfaces*, J. Alg. Geom. **8** (1999), 245–278.
- [2] C. Ciliberto, F. Flamini, C. Galati, A.L. Knutsen, *Moduli of nodal curves on K3 surfaces*, arXiv:1502.07378.
- [3] R. Friedman, *A new proof of the global Torelli theorem for K3 surfaces*, Annals of Math. **120** (1984), 237–269.
- [4] R. Friedman, *Global smoothings of varieties with normal crossings*, Annals of Math. **118** (1983), 75–114.
- [5] M. Kemeny, *The moduli of singular curves on K3 surfaces*, J. Math. Pures Appl. **104** (2015), 882–920.
- [6] V. Kulikov, *Degenerations of K3 surfaces and Enriques surfaces*, Math. USSR Izvestiya **11** (1977), 957–989.
- [7] C. Galati, *On the existence of curves with a triple point and nodes on K3-surfaces*, Rend. Lincei Mat. Appl. **23** (2012), 295–317.
- [8] C. Galati, A. L. Knutsen, *On the existence of curves with A_k -singularities on K3 surfaces*, Math. Res. Lett. **21** (2014), 1069–1109.
- [9] C. Galati, A. L. Knutsen, *Seshadri constants of K3 surfaces of degrees 6 and 8*, Int. Math. Res. Not. **2013** (2013), 4072–4084.
- [10] U. Persson, H. Pinkham, *Degenerations of surfaces with trivial canonical bundle*, Annals of Math. **113** (1981), 45–66.

Problems related to “fake” $K3$ surfaces

EDOARDO SERNESI

In [1] we consider a certain class of rational and ruled surfaces, sometimes called *fake $K3$ surfaces*. The relation between the smoothability of such surfaces to $K3$ surfaces and the existence of Petri general curves on them is investigated. The analysis in [1] suggests some interesting questions not yet investigated and related with moduli of curves, $K3$ surfaces and with classical problems on Del Pezzo surfaces of degree 1. Perhaps the most interesting one is:

Question 27. *Assume $g \geq 12$. Is the locus $(K3)_g \subset \mathcal{M}_g$ of curves lying on a $K3$ surface closed?*

For instance, there are specific curves lying on smoothable fake $K3$ s for which we could not decide whether they are Petri general or not. The study of such curves can be translated into questions concerning Severi varieties on Del Pezzo surfaces of degree 1. The all subject has connections with the study of moduli spaces of stable $K3$ surfaces. Another vague but suggestive problem is the following one:

Question 28. *Is it possible to generalize in some way Lazarsfeld’s proof [2] of Gieseker-Petri’s theorem to curves lying on a fake $K3$ surface?*

REFERENCES

- [1] E. Arbarello, A. Bruno, E. Sernesi, *On two conjectures by J. Wahl*, arXiv:1507.05002.
- [2] R. Lazarsfeld, *Brill-Noether-Petri without degenerations*, J. Diff. Geom. **23** (1986), 299–307.

Geometry of O’Grady’s 6 dimensional example

GIULIA SACCÀ

(joint work with G. Mongardi, A. Rapagnetta)

There are not many known examples of hyperkähler manifolds. Two series of examples appear in dimension $2n$, for every $n > 1$, and are related to the Hilbert scheme of points on a $K3$ or an abelian surface; and in dimension 6 and 10 there is one extra, or exceptional, deformation class, each of which was found by O’Grady. While considerable work has been devoted to studying hyperkähler manifolds belonging to the first two deformation classes, not much is known for the exceptional deformation classes. We present results from [2] regarding the geometry of O’Grady’s six dimensional example [3]. We use Lehn and Sorger’s local analytic description of O’Grady’s singularity [1], and results from [4] to realize these examples as “quotients” of another hyperkähler manifold by a birational involution. Since this hyperkähler manifold turns out to be deformation equivalent to the Hilbert scheme of 3 points on a $K3$ surface, we are able to compute all the Hodge numbers and study properties of their moduli spaces.

REFERENCES

- [1] M. Lehn, Ch. Sorger, *La singularité d'O'Grady*, J. Algebraic Geom. **15** (2006), 735–770.
- [2] G. Mongardi, A. Rapagnetta, G. Saccà, *The Hodge diamond of O'Grady's six dimensional example*, to appear, Fall 2015.
- [3] K. O'Grady, *A new six-dimensional irreducible symplectic variety* J. Algebraic Geom. **12** (2003), 435–505.
- [4] A. Rapagnetta, *Topological invariants of O'Grady's six dimensional irreducible symplectic variety*, Math. Z. **256** (2007), 1–34.

Syzygies of curves and K3 surfaces

MICHAEL KEMENY

(joint work with G. Farkas)

K3 surfaces have recently been used to make significant progress on several conjectures concerning syzygies of curves on a K3 surface. We discuss recent work in [5] and [6].

Let C be a curve and L an ample line bundle. Following Green, define the Koszul group $K_{i,j}(C, L)$ as the middle cohomology of

$$\bigwedge^{i+1} H^0(C, L) \otimes H^0(C, (j-1)L) \rightarrow \bigwedge^i H^0(C, L) \otimes H^0(C, jL) \rightarrow \bigwedge^{i-1} H^0(C, L) \otimes H^0(C, (j+1)L).$$

One says (C, L) satisfies *property* (N_p) if we have the vanishings

$$K_{i,j}(C, L) = 0 \text{ for } i \leq p, j \geq 2.$$

Then $\phi_L : C \hookrightarrow \mathbb{P}^r$ is projectively normal if and only if (C, L) satisfies (N_0) , whereas the ideal of C is generated by quadrics if, in addition, it satisfies (N_1) .

The line bundle L is called *p -very ample* if and only if for every effective divisor D of degree $p+1$ the evaluation map

$$ev : H^0(C, L) \rightarrow H^0(D, L|_D)$$

is surjective. Equivalently, L is *not p -very ample* if and only if $C \subseteq \mathbb{P}^r$ admits a $(p+1)$ -secant $(p-1)$ -plane.

The Secant Conjecture of Green–Lazarsfeld then states:

Conjecture 29 (Secant Conjecture). *Let L be a globally generated line bundle of degree d on a curve C of genus g such that*

$$d \geq 2g + p + 1 - 2h^1(C, L) - \text{Cliff}(C).$$

Then (C, L) fails property (N_p) if and only if L is not $p+1$ -very ample.

We discuss the recent proof of the Secant Conjecture for general curves. The proof uses moduli spaces of polarised K3 surfaces and relies on Voisin's proof of Green's conjecture for curves on suitably general K3 surfaces [3, 4]. It is rather straightforward to see that if L is not $p+1$ very ample, that is, L admits a $(p+2)$ -secant p -plane, then $K_{p,2}(C, L)$ is nonzero. The difficulty in establishing the above

conjecture is thus to go in the other direction, that is, to construct a secant plane out of a syzygy in $K_{p,2}(C, L)$

The second conjecture we consider is the *Prym–Green Conjecture*. Consider a smooth curve C of genus g , and fix a nontrivial torsion line bundle η of order l . A *paracanonical curve* is the embedded curve

$$\phi_{\omega_C \otimes \eta} : C \hookrightarrow \mathbb{P}^{g-2}.$$

For a general *canonical curve*

$$\phi_{\omega_C} : C \hookrightarrow \mathbb{P}^{g-1}$$

the famous work of Voisin [3, 4] suffices to describe the shape of the free resolution of the homogeneous coordinate ring of C . The Prym–Green conjecture likewise predicts a roughly similar shape for the homogeneous coordinate ring of a *general paracanonical curve*. We describe our proof of this when g is odd and l is either two (the classical case) or l large. The proof works by degenerating curves on special $K3$ surfaces and uses work of van Geemen–Sarti [2] resp. Barth–Verra [1] in the $l = 2$ resp. large l cases.

REFERENCES

- [1] W. Barth, A. Verra, *Torsions of $K3$ sections*, Problems in the theory of surfaces and their classification (Cortona, 1988), Symposia Math., Vol. XXXII, Academic Press 1991, 1–24.
- [2] B. van Geemen, A. Sarti, *Nikulin involutions on $K3$ surfaces*, Math. Z. **255** (2007), 731–753.
- [3] C. Voisin, *Green’s generic syzygy conjecture for curves of even genus lying on a $K3$ surface*, J. European Math. Soc. **4** (2002), 363–404.
- [4] C. Voisin, *Green’s canonical syzygy conjecture for generic curves of odd genus*, Compos. Math. **141** (2005), 1163–1190.
- [5] G. Farkas, M. Kemeny, *The generic Green–Lazarsfeld Secant conjecture*, Invent. Math. (to appear).
- [6] G. Farkas, M. Kemeny, *The Prym–Green conjecture for torsion bundles of high order*, arXiv:1509.07162.

Twisted cubics on a cubic fourfold and in involution on the associated 8-dimensional symplectic manifold

MANFRED LEHN

(joint work with Ch. Lehn, Ch. Sorger, D. van Straten; N. Addington; and I. Dolgachev)

A by now classical theorem of Beauville and Donagi states that the Fano variety of lines on a smooth cubic fourfold Y is an irreducible holomorphic symplectic manifold of dimension 4 that is deformation equivalent to the second Hilbert scheme of a $K3$ surface. With the intention to generalise this construction to curves of higher degree, we consider the moduli space of generalised twisted cubics on Y . If the fourfold Y does not contain a plane, this moduli space $M = M_3(Y)$ turns out to be smooth projective of dimension 10. The natural morphism $M \rightarrow G := \text{Grass}(3, \mathbb{P}^5)$ that sends a curve $C \subset Y \subset \mathbb{P}^5$ to the three-dimensional linear span $\langle C \rangle$, fibres as follows: $M \rightarrow Z' \rightarrow Z_{\text{Stein}} \rightarrow G$, where $Z_{\text{Stein}} \rightarrow G$ is the finite part of the Stein

factorisation, $Z' \rightarrow Z_{\text{Stein}}$ is a resolution of singularities, and, most importantly, $M \rightarrow Z'$ is a $\mathbb{P}^2 \rightarrow Z'$ -bundle.

According to a result of Piene and Schlessinger, twisted cubic curves in 3-space come in two flavours: the general curve is arithmetically Cohen-Macaulay and defined by three quadrics that are the minors of a (3×2) -matrix A_0 with linear entries. Points in a smooth divisor represent non-Cohen-Macaulay curves. Such curves are singular plane cubics with an embedded point at a singularity. Non-Cohen-Macaulay curves also form a smooth divisor in M , and this divisor is contracted to a smooth divisor $D \subset Z'$. Finally, there is a divisorial contraction $Z' \rightarrow Z = Z(Y)$ to an 8-dimensional holomorphic symplectic manifold Z . Under this map, the divisor D is mapped to a copy of Y in Z , and curve C with embedded point $p \in C$ is mapped to this point $p \in Y$.

Every twisted curve C of aCM-type determines an integral cubic surface $S = Y \cap \langle C \rangle$ and a linear determinantal representation $g = \det(A)$ of the equation g of S in the linear hull $\langle C \rangle$. A curve of nonCM-type corresponds to a skew-symmetric (3×3) -matrix A the entries of which define the position of the embedded point.

For surfaces S with at most ADE -singularities, families of twisted cubics are given by equivalence classes of roots in the E_6 -type root system of the lattice $K_{\tilde{S}}^{\perp} \subset H^2(\tilde{S}, \mathbb{Z})$ in the integral cohomology of the minimal resolution $\tilde{S} \rightarrow S$. Two roots are equivalent if they lie in the same orbit under the action of the Weyl group generated by effective roots, i.e. exceptional curves of the resolution $\tilde{S} \rightarrow S$.

The two stages of the contraction $M \rightarrow Z' \rightarrow Z$ can be interpreted in terms of left mutations from the derived category $\mathcal{D}(Y)$ onto Kuznetsov's subcategory $\mathcal{A} \subset \mathcal{D}(Y)$, and Z can be seen as a moduli space of objects in \mathcal{A} .

It is interesting to note that Z admits a non-symplectic involution i that can be described as follows: For any C the sections in $H^0(I_{C/S}(2))$ correspond to quadric Q that contain C . As the intersection $Q \cap S$ is curve of degree 6, it decomposes as $Q \cap S = C \cup C'$, and the involution takes (the family of) C to (the family of) C' . In terms of root systems, the involution is given by $\alpha \mapsto -\alpha$. Finally, in terms of matrices, it is $A \mapsto A^t$. From the last description it is clear that the fixed point locus has two components Y and Y' , the former corresponding to skew-symmetric matrices and hence nonCM-curves, the latter corresponding to symmetric matrices and thus symmetric linear determinantal representations. Such representations only exist for surfaces with four A_1 -singularities and for degenerations of such surfaces. So roughly, the second fixed point component is given by four-tuples of points on Y with the property that the 3-plane through the points is tangent to Y in each point. We hope that the quotient Y/i can be embedded into \mathbb{P}^{20} , by a map that restricts on the first fixed point component Y to the second Veronese map, and that this will allow to describe the quotient directly in terms of the projective geometry of Y leading to a construction of Z analogous to the double EPW-sextics of O'Grady.

REFERENCES

- [1] C. Lehn, M. Lehn, C. Sorger, D. van Straten, *Twisted cubics on cubic fourfolds*, *J. Reine Angew. Math.* (2015), DOI 10.1515/crelle-2014-0144.
- [2] N. Addington, M. Lehn, *On the symplectic 8-fold associated to a Pfaffian cubic fourfold*, *J. Reine Angew. Math.* (2015), DOI 10.1515/crelle-2014-0145.

On the problem of compactification of Lagrangian fibrations

DIMITRI MARKUSHEVICH

The following problem is studied: given an abelian fibration, when can one endow its total space with a symplectic structure in such a way that it becomes a Lagrangian fibration (LF), compactifiable to a possibly singular irreducible symplectic variety (ISV)?

Partial answers to this question were obtained for relative Jacobians of families of curves [5, 7] and relative Prymians [6, 1]. We give a survey of results of the thesis by J. Bouali [2] on the compactification of the Lagrangian fibrations in intermediate Jacobians associated to $K3$ -Fano flags [4]. Then an approach is presented toward a compactification of the Donagi–Markman LF [3] in intermediate Jacobians of cubic 3-folds that are hyperplane sections of a fixed cubic 4-fold, based upon the following almost-theorem:

Conjecture 30. *Let $X \subset \mathbb{P}^5$ be a generic cubic 4-fold, and let $\{V_t\}_{t \in \mathbb{P}^{5*}}$ denote the family of its hyperplane sections. Then for any $t_0 \in \mathbb{P}^{5*}$, the following properties are verified:*

- (1) $\text{Sing}(V_{t_0}) = \{p_1, \dots, p_k\}$ is finite, the singular points p_i are of types A_{μ_i} ($1 \leq \mu_i \leq 5$) or D_{μ_i} ($4 \leq \mu_i \leq 5$), and $\sum \mu_i \leq 5$.
- (2) The family $\mathcal{V} = \{V_t\}_{t \in \mathbb{P}^{5*}} \rightarrow \mathbb{P}^{5*}$ is locally analytically a multi-versal unfolding of $\text{Sing}(V_{t_0})$ near t_0 .

Thus, the degenerations of the family $\mathcal{V}/\mathbb{P}^{5*}$ are well-behaved and there is a hope to compactify the relative intermediate Jacobian $J = J(\mathcal{V}/\mathbb{P}^{5*})$ to an ISV. An evidence is given by the fact that the relative compactified Jacobian of a family of integral curves with plane singularities is non-singular, provided the family is multi-versal for each of its singular fibers. Remark that the monodromy of the versal deformation of an isolated curve singularity coincides with the monodromy of the versal deformation of the stably equivalent 3-dimensional singularity, which motivates the following question:

Question 31. *Is the relative compactified Jacobian of a family of curves over a polydisk in \mathbb{C}^k toroidal, provided that the family is a multi-versal unfolding of singularities of its central fiber, and that those singularities are only simple singularities of types A_{μ_i} ($1 \leq \mu_i \leq 5$) or D_{μ_i} ($4 \leq \mu_i \leq 5$), $\sum \mu_i \leq 5$?*

An alternative approach using a representation of the intermediate Jacobians as Prym varieties of double covers of curves is discussed.

REFERENCES

- [1] E. Arbarello, G. Saccà, A. Ferretti, *Relative Prym varieties associated to the double cover of an Enriques surface*, J. Diff. Geom. **100** (2015), 191–250.
- [2] J. Bouali, *Motifs des fibrés en quadriques et jacobiniennes intermédiaires relatives des paires K3-Fano*, Ph.D.-thesis (J. Nagel and D. Markushevich, Supervisors), Dijon, 2015.
- [3] R. Donagi, E. Markman, *Spectral covers, algebraically completely integrable, Hamiltonian systems, and moduli of bundles*, Lecture Notes in Math. **1620** (1996), 1–119.
- [4] D. Markushevich, *An integrable system of K3-Fano flags*, Math. Ann. **342** (2008), 145–156.
- [5] D. Markushevich, *Lagrangian families of Jacobians of genus 2 curves*, J. Math. Sci. **82** (1996), 3268–3284.
- [6] D. Markushevich, A.S. Tikhomirov, *New symplectic V-manifolds of dimension four via the relative compactified Prymian*, Internat. J. Math. **18** (2007), 1187–1224.
- [7] J. Sawon, *On Lagrangian fibrations by Jacobians I*, J. Reine Angew. Math. **701** (2015), 127–151.

Short report on discussion sessions

One group of participants, formed by Ciliberto, Dedieu, Galati, Kemeny, Knutsen, Saccà and Sernesi concentrated on some of the problems proposed by Ciliberto/Flamini and Sernesi. In particular, they figured out a way of possibly answering Question 13 affirmatively. Concerning Question 20, they observed that $V_{n,\delta}(X)$ is never rationally connected if $(X, L) \in \mathcal{K}_p$ is general.

Some time was also devoted to Question 28 towards extending Lazarsfeld’s vector bundle methods to fake K3 surfaces. A few days after the workshop, the problem was solved in [1], which in particular proves that certain hyperplane sections of fake K3 surfaces are indeed Brill-Noether-Petri general curves.

A second group of participants, formed by Boissière, Camere, Cattaneo, Flamini, Lehn, Markushevich, Ohashi and Sarti focused on finding geometric description of the involution on $S^{[2]}$ as in Question 1 proposed by Cattaneo. Various methods of attack were considered. The ideas developed were very useful and we believe that these will help in future research in finding the geometric description.

Some time was also devoted to discussing other problems, in various constellations of participants. Among those, considerable attention was devoted to degenerations of hyperkähler manifolds as in Question 26 proposed by Galati. Great progress in the study of deformations of degenerate hyperkähler manifolds was made in [2]. Nevertheless, up to now, very few examples of degenerate hyperkähler manifolds are known. Various attempts to find new examples were made.

REFERENCES

- [1] E. Arbarello, A. Bruno, G. Farkas, G. Saccà, *Explicit Brill-Noether-Petri general curves*, arXiv:1511.07321.
- [2] D. Hanke: *Logarithmically smooth deformations of strict normal crossing logarithmically symplectic varieties*, Ph.D.-thesis, Johannes Gutenberg-Universität Mainz, 2015.

Reporters: Concettina Galati, Andreas Leopold Knutsen, Alessandra Sarti

Participants

Prof. Dr. Samuel Boissière

Lab. de Mathématiques et Applications
Université de Poitiers
Blvd. Marie et Pierre Curie
86962 Chasseneuil Cedex
FRANCE

Prof. Dr. Chiara Camere

Dipartimento di Matematica
Università di Milano
Via C. Saldini, 50
20133 Milano
ITALY

Dr. Andrea Cattaneo

Dipartimento di Matematica
Università degli Studi di Parma
Parco Area delle Scienze, 53/A
43124 Parma
ITALY

Prof. Dr. Ciro Ciliberto

Dipartimento di Matematica
Università di Roma Tor Vergata
Via della Ricerca Scientifica, 1
00133 Roma
ITALY

Prof. Dr. Thomas Dedieu

Institut de Mathématiques de Toulouse
Université Paul Sabatier
118, route de Narbonne
31062 Toulouse Cedex 9
FRANCE

Prof. Dr. Flaminio Flamini

Dipartimento di Matematica
Università degli Studi di Roma II
Tor Vergata
Via della Ricerca Scientifica, 1
00133 Roma
ITALY

Prof. Dr. Concettina Galati

Dipartimento di Matematica
Università della Calabria
87036 Rende CS
ITALY

Dr. Michael Kemeny

Institut für Mathematik
Humboldt-Universität
Unter den Linden 6
10099 Berlin
GERMANY

Prof. Dr. Andreas L. Knutsen

Department of Mathematics
University of Bergen
Postboks 7800
5020 Bergen
NORWAY

Prof. Dr. Manfred Lehn

Institut für Mathematik
Johannes-Gutenberg Universität Mainz
55099 Mainz
GERMANY

Prof. Dr. Dimitri Markouchevitch

U. F. R. Mathématiques
Université de Lille 1
59655 Villeneuve d'Ascq Cedex
FRANCE

Prof. Dr. Hisanori Ohashi

Department of Mathematics
Faculty of Science and Technology
Tokyo University of Science
Yamazaki 2641
Chiba 278-8510
JAPAN

Dr. Giulia Saccà

Mathematics Department
James H. Simons Instructor
Stony Brook University
Math. Tower 3-116
Stony Brook NY 11794-3651
UNITED STATES

Prof. Dr. Edoardo Sernesi

Dipartimento di Matematica e Fisica
Università di Roma Tre
Largo San Leonardo Murialdo, 1
00146 Roma
ITALY

Prof. Dr. Alessandra Sarti

Département de Mathématiques
Université de Poitiers
Boite Postale 30179
86962 Futuroscope Chasseneuil Cedex
FRANCE

