

MATHEMATISCHES FORSCHUNGSINSTITUT OBERWOLFACH

Report No. 57/2018

DOI: 10.4171/OWR/2018/57

## Mini-Workshop: Mathematical and Numerical Analysis of Maxwell's Equations

Organised by  
Monique Dauge, Rennes  
Ulrich Langer, Linz  
Peter Monk, Newark  
Dirk Pauly, Essen

16 December – 22 December 2018

ABSTRACT. In this mini-workshop 17 leading mathematicians from Europe and United States met at the MFO to discuss and present new developments in the mathematical and numerical analysis of Maxwell's equations and related systems of partial differential equations. The report at hand offers the extended abstracts of their talks.

*Mathematics Subject Classification (2010):* MSC: 45xxx, 46xxx, 47xxx, 65xxx, 78xxx.

### Introduction by the Organisers

The mini-workshop *Mathematical and Numerical Analysis of Maxwell's Equations*, organised by Monique Dauge (Rennes), Ulrich Langer (Linz), Peter Monk (Newark), and Dirk Pauly (Essen) was well attended with 17 participants with broad geographic representation from Europe and United States. This workshop was a nice blend of researchers with various backgrounds from Maxwell's equations.

Maxwell's equations of electro-dynamics are of huge importance in mathematical physics, engineering, and especially in mathematics, leading since their discovery to interesting mathematical problems and even to new fields of mathematical research, particularly in the analysis and numerics of partial differential equations and applied functional analysis. The deep understanding of Maxwell's equations and the possibility of their numerical solution in complex geometries and different settings have led to very efficient and robust simulation methods in Computational Electromagnetics. Moreover, efficient simulation methods pave the way for optimizing electromagnetic devices and processes. Digital communication and

e-mobility are two fields where simulation and optimization techniques that are based on Maxwell's equations play a deciding role.

The workshop brought together different communities, namely people working in analysis of Maxwell's equations with those working in numerical analysis of Maxwell's equations and computational electromagnetics and acoustics.

*Acknowledgement:* The MFO and the workshop organizers would like to thank the National Science Foundation for supporting the participation of junior researchers in the workshop by the grant DMS-1641185, "US Junior Oberwolfach Fellows".

## Mini-Workshop: Mathematical and Numerical Analysis of Maxwell's Equations

### Table of Contents

Ana Alonso Rodríguez (joint with Jessika Camaño, Eduardo De Los Santos, Francesca Rapetti)	
<i>High order discrete potentials</i> .....	5
Anne-Sophie Bonnet-Ben Dhia (joint with Lucas Chesnel and Patrick Ciarlet)	
<i>Time-harmonic Maxwell equations with sign-changing coefficients</i> .....	6
Martin Costabel	
<i>On the regularity of electromagnetic fields on Lipschitz domains</i> .....	7
Monique Dauge (joint with Stéphane Balac and Zoïs Moitier)	
<i>Resonances of Optical Micro-Resonators</i> .....	8
Jay Gopalakrishnan (joint with Martin Neumüller, Panayot Vassilevski)	
<i>The HX Preconditioner in Higher Dimensions</i> .....	9
Lise-Marie Imbert-Gérard (joint with J.-F. Fritsch)	
<i>Generalized Plane Waves &amp; Maxwell's equations</i> .....	10
Peter Monk	
<i>Finite Element Methods for Maxwell's Equations</i> .....	11
Serge Nicaise (joint with Jérôme Tomezyk)	
<i>The time-harmonic Maxwell equations with impedance boundary conditions in domains with a polyhedral or an analytic boundary</i> .....	13
Frank Osterbrink (joint with Dirk Pauly)	
<i>Time-harmonic Scattering in exterior domains with mixed boundary conditions</i> .....	15
Dirk Pauly (joint with Sebastian Bauer, Michael Schomburg, and Walter Zulehner)	
<i>On Some Compact Embeddings in Various Hilbert Complexes</i> .....	17
Ilaria Perugia (joint with Lorenzo Mascotto, Alexander Pichler)	
<i>Nonconforming Trefftz virtual elements for the Helmholtz problem</i> .....	18
Rainer Picard	
<i>Memoryless Evolutionary Problems as Dynamic Abstract Friedrichs Systems</i> .....	19
Joachim Schöberl (joint with Astrid Pechstein, Michael Neunteufel, Philip Lederer)	
<i>Solving Elasticity in <math>H(\text{curl})</math></i> .....	21

Alberto Valli

*On the helicity of a bounded domain and the Biot–Savart operator* . . . . . 21

Marcus Waurick

*On Nonlocal  $H$ -convergence* . . . . . 23

Ragnar Winther (joint with Richard S. Falk)

*Geometric decompositions in finite element exterior calculus* . . . . . 24

## Abstracts

### High order discrete potentials

ANA ALONSO RODRÍGUEZ

(joint work with Jessika Camaño, Eduardo De Los Santos, Francesca Rapetti)

In applied computations the need often arises to define, for example, a discrete field with assigned curl or to represent a div-free field in a given discrete space. In the low degree case this need can be fulfilled by involving tree and co-tree techniques (see, e.g., [2]). Our aim is to extend this techniques to high order Whitney finite elements. The key point is to identify the degrees of freedom for which the matrices representing the divergence and the gradient operators are related with the incidence matrix of a connected oriented graph. This can be done using Bernstein polynomials in the definition of the classical moments (see, e.g. [4]) usually used as degrees of freedom. A convenient visualization of these graphs can be obtained using the so-called small simplices introduced in [5].

Once it have been proved that the matrix associated with the divergence operator is the incidence matrix of a connected oriented graph, it is possible to identify an invertible square submatrix of this incidence matrix by choosing a spanning tree of the graph. This allows to easily compute the moments of a field in the space of Raviart-Thomas finite elements with assigned divergence. This approach extends to finite elements of high degree the method introduced in [3] for finite elements of degree one and can be used to construct a basis of the space of divergence-free Raviart-Thomas finite elements. The numerical tests show that the performance of the algorithm does not depend on the topology of the domain or the polynomial degree.

### REFERENCES

- [1] A. Alonso Rodríguez, J. Camaño, E. De Los Santos, and F. Rapetti, *A graph approach for the construction of high order divergence-free Raviart-Thomas finite elements*, *Calcolo* **55**, (2018), 28 pp.
- [2] A. Alonso Rodríguez and A. Valli. *Finite element potentials* *Appl. Numer. Math.* **95** (2015), 2–14.
- [3] P. Alotto and I. Perugia. *Mixed finite element methods and tree-cotree implicit condensation*, *Calcolo* **36** (1999), 233–248.
- [4] D.N. Arnold, R.S. Falk, and R. Winther, *Finite element exterior calculus, homological techniques and applications*, *Acta Numerica* **15**, (2006), 1–155.
- [5] F. Rapetti and A. Bossavit, *Whitney forms of higher degree*, *SIAM J. Numer. Anal.* **47**, (2009), 2369–2386.

## Time-harmonic Maxwell equations with sign-changing coefficients

ANNE-SOPHIE BONNET-BEN DHIA

(joint work with Lucas Chesnel and Patrick Ciarlet)

In [1], we consider the time-harmonic Maxwell equations in a bounded domain  $\Omega \subset \mathbb{R}^3$ , when the dielectric permittivity  $\varepsilon$  and the magnetic permeability  $\mu$  are real functions whose sign can change ( $\varepsilon$ ,  $\mu$ ,  $1/\varepsilon$  and  $1/\mu$  are just supposed to be bounded functions). We assume for simplicity here that  $\mu = 1$  and that  $\Omega$  is topologically trivial, and we consider the variational problem:

$$(1) \quad \text{Find } E \in X \text{ such that } \forall E' \in X, \int_{\Omega} \text{curl } E \cdot \text{curl } E' - \omega^2 \varepsilon E \cdot E' = \int_{\Omega} J \cdot E'$$

where  $\omega \in \mathbb{R}$ ,  $J \in L^2(\Omega)^3$  is such that  $\text{div } J = 0$  in  $\Omega$ , and  $X$  can be one of these two functional spaces:

$$H_N = \{E \in L^2(\Omega)^3, \text{curl } E \in L^2(\Omega)^3 \text{ and } E \times n = 0 \text{ on } \partial\Omega\},$$

$$\text{and } X_N(\varepsilon) = \{E \in H_N, \text{div}(\varepsilon E) = 0 \text{ in } \Omega\}.$$

The usual approach (for positive coefficients) consists in proving the following two results:

- (1) Problem (1) with  $X = X_N(\varepsilon)$  is equivalent to Problem (1) with  $X = H_N$ .
- (2) The embedding of  $X_N(\varepsilon)$  in  $L^2(\Omega)^3$  is compact.

We prove that both assertions remain true for a sign-changing  $\varepsilon$  if

$$(2) \quad \forall f \in H^{-1}(\Omega), \exists! \varphi \in H_0^1(\Omega) \text{ such that } \forall \varphi' \in H_0^1(\Omega), \int_{\Omega} \varepsilon \nabla \varphi \cdot \nabla \varphi' = \int_{\Omega} f \varphi'$$

This scalar problem has been the subject of many studies in the case where the sign of  $\varepsilon$  changes across a Lipschitz boundary  $\Sigma$  (see for instance [2, 4, 3, 5]). Let us point out that this problem can be ill-posed, and even not Fredholm in some configurations (for instance, if a part of  $\Sigma$  is flat and if  $\varepsilon$  takes opposite values on each side).

To prove assertion 1, the idea is simply to notice that for every test field  $E' \in H_N$ , there exists  $\varphi \in H_0^1(\Omega)$  such that  $E' + \nabla \varphi \in X_N(\varepsilon)$ .

The proof of assertion 2 relies on a  $T$ -coercivity result. More precisely, we show that, if (2) is true, there exists an operator  $T \in \mathcal{L}(X_T)$  such that

$$\forall H, H' \in X_T, \int_{\Omega} \frac{1}{\varepsilon} \text{curl } H \cdot \text{curl}(TH') = \int_{\Omega} \text{curl } H \cdot \text{curl } H'$$

where  $X_T$  denotes the usual space for the magnetic fields:

$$X_T = \{H \in L^2(\Omega)^3, \text{curl } H \in L^2(\Omega)^3, \text{div } H = 0 \text{ in } \Omega \text{ and } H \cdot n = 0 \text{ on } \partial\Omega\}.$$

The compactness of the embedding is finally established by a proof *à la Weber* [6].

Finally, suppose that the scalar problem is Fredholm with a non trivial finite dimensional kernel, so that (2) is not satisfied. In that case, the first assertion is no longer true. Indeed, let  $\varphi$  belong to the kernel of the scalar problem. Then  $E = \nabla \varphi$  is a solution of (1) with  $J = 0$  for  $X = X_N(\varepsilon)$  and not for  $X = H_N$ . However the approach can be extended to this case by setting problem (1) in an

appropriate functional space  $X = \tilde{X}_N(\varepsilon)$  such that  $\tilde{X}_N(\varepsilon) = X_N(\varepsilon) \oplus Y_N(\varepsilon)$  where  $\dim(Y_N(\varepsilon)) < +\infty$ .

## REFERENCES

- [1] A.-S. Bonnet-Ben Dhia, L. Chesnel and P. Ciarlet Jr., *T-Coercivity for the Maxwell Problem with Sign-Changing Coefficients*, Communications in Partial Differential Equations, **39** (2014), 1007-1031.
- [2] M. Costabel and E. Stephan, *A direct boundary integral equation method for transmission problems*, J. Math. Anal. Appl. **106** (1985), 367-413.
- [3] A.-S. Bonnet-Bendhia, M. Dauge and K. Ramdani, *Analyse spectrale et singularités d'un problème de transmission non coercif*, Comptes Rendus de l'Académie des Sciences - Series I - Mathematics, **328** (1999), 717-720.
- [4] A.-S. Bonnet-Ben Dhia, L. Chesnel and P. Ciarlet Jr., *T-coercivity for scalar interface problems between dielectrics and metamaterials*, ESAIM: M2AN **46** (2012), 1363-1387.
- [5] S. Nicaise and J. Venel, *A posteriori error estimates for a finite element approximation of transmission problems with sign changing coefficients*, J. Comput. Appl. Math. **235** (2011) 4272-4282.
- [6] C. Weber, *A local compactness theorem for Maxwell's equations*, Math. Meth. Appl. Sci., **2** (1980), 12-25.

## On the regularity of electromagnetic fields on Lipschitz domains

MARTIN COSTABEL

The usual variational spaces  $X_N$  and  $X_T$  for time-harmonic electric or magnetic fields with PEC boundary conditions on a bounded Lipschitz domain  $\Omega \subset \mathbb{R}^3$  are defined as

$$X_N = H(\operatorname{div}, \Omega) \cap H_0(\operatorname{curl}, \Omega) \quad \text{and} \quad X_T = H_0(\operatorname{div}, \Omega) \cap H(\operatorname{curl}, \Omega).$$

These spaces have long been known to be contained in the Sobolev space  $H^{1/2}(\Omega)$  [3, 6]. If the domain is, in addition, piecewise smooth, then there exists always an  $\varepsilon > 0$  such that these spaces are contained in  $H^{1/2+\varepsilon}(\Omega)$ , and this additional smoothness is known to be useful in the analysis of some numerical algorithms [2, 1].

At a conference in 2017, Alberto Valli brought up the question whether such an additional regularity is also present for every Lipschitz domain. The answer turns out to be negative.

Using ideas from Nikolai Filonov's construction of a  $C^{3/2}$  domain for which the usual Birman-Solomyak decomposition of  $X_T$  is not possible [5], a bounded domain  $\Omega$  can be constructed that is of class  $C^1$  and for which such an  $\varepsilon > 0$  does not exist. This domain has the property that for functions in  $W^{1+\frac{1}{p}+\varepsilon, p}(\Omega)$  with any  $\varepsilon > 0$  and  $p \geq 1$ , the vanishing of the trace on  $\partial\Omega$  implies the vanishing of the normal derivative. As a consequence, there exist smooth right hand sides for which the solutions of the Poisson equation with homogeneous Dirichlet or Neumann conditions (known to belong to  $W^{1+\frac{1}{p}, p}(\Omega)$  for  $p > 1$ ) do not belong to  $W^{1+\frac{1}{p}+\varepsilon, p}(\Omega)$  with any  $\varepsilon > 0$  and  $p \geq 1$ . Taking gradients of these solutions and

$p = 2$  gives the non-regularity result for  $X_N$  and  $X_T$ . This construction is the subject of the article [4].

#### REFERENCES

- [1] M. AINSWORTH, J. GUZMÁN, AND F.-J. SAYAS, *Discrete extension operators for mixed finite element spaces on locally refined meshes*, Math. Comp., 85 (2016), pp. 2639–2650.
- [2] A. ALONSO AND A. VALLI, *An optimal domain decomposition preconditioner for low-frequency time-harmonic Maxwell equations*, Math. Comp., 68 (1999), pp. 607–631.
- [3] M. COSTABEL, *A remark on the regularity of solutions of Maxwell’s equations on Lipschitz domains*, Math. Methods Appl. Sci., 12 (4) (1990), 365–368.
- [4] M. COSTABEL, *On the limit Sobolev regularity for Dirichlet and Neumann problems on Lipschitz domains*, Math. Nachr. (2019), accepted.
- [5] N. FILONOV, *Principal singularities of the magnetic field component in resonators with a boundary of a given class of smoothness*, Algebra i Analiz, 9 (1997), pp. 241–255.
- [6] D. MITREA AND M. MITREA, *Finite energy solutions of Maxwell’s equations and constructive Hodge decompositions on nonsmooth Riemannian manifolds*, J. Funct. Anal., 190 (2002), pp. 339–417.

## Resonances of Optical Micro-Resonators

MONIQUE DAUGE

(joint work with Stéphane Balac and Zoïa Moitier)

Optical micro-resonators are dielectric cavities of small size (at  $\mu\text{m}$  scale) coupled to waveguides or fibers for light input and output [5]. The frequencies of interaction have to be close to resonance frequencies of the cavity alone (i.e. without neighboring fibers). Of particular interest are the whispering gallery modes: These special resonant modes concentrate along the boundary of the cavity and are mainly generated by curvature effects at high frequency.

As a simplified model for the three-dimensional Maxwell system (effective index method [11]), we consider two-dimensional [2D] Helmholtz equations governing transverse electric [TE] or magnetic [TM] modes. Even in this 2D framework, very few results provide asymptotic expansion of WGM at high frequency, see [6] for disk micro-resonators using expansions of Bessel functions [8].

In [3] we present a unified procedure to construct WGM in 2D cavities  $\Omega$  with smooth boundaries and varying optical index  $n$ . Note that  $n$  is defined as 1 outside the cavity (in  $\mathbb{R}^2 \setminus \overline{\Omega}$ ) and is assumed to be smooth and away from 1 in  $\overline{\Omega}$ . The problem under consideration is: Find non-zero  $u$  and complex number  $k$  such that—here the integer  $p \in \{+1, -1\}$  distinguishes TM modes ( $p = 1$ ) and TE modes ( $p = -1$ ):

$$(1) \quad -\operatorname{div}(n^{p-1} \nabla u) = k^2 n^{p+1} u \quad \text{in } \mathbb{R}^2$$

complemented by suitable radiation condition (expansion at infinity in terms of Hankel functions of the first kind). Equation (1) includes the jump conditions  $[u] = 0$  and  $[n^{p-1} \partial_\nu u] = 0$  on  $\partial\Omega$ . Couples  $(k, u)$  are the resonant modes. In contrast with impenetrable obstacles [10], transparent obstacles or dielectric cavities may have non-real resonances super-algebraically close to the real axis [9, 7, 4].

We have found that the variations of  $n$  give rise to a new effective curvature index  $\check{\kappa} = \kappa + n'/n$  (where  $\kappa$  is the usual curvature of  $\partial\Omega$ ) whose sign governs the asymptotic type:  $\check{\kappa} > 0$  causes modes to concentrate along the boundary with Airy-type profiles,  $\check{\kappa} = 0$  still causes concentration along the boundary but with Harmonic Oscillator profiles, whereas  $\check{\kappa} < 0$  may produce internal WGM types. By multiscale expansions based on a WKB approximation, we construct asymptotic quasi-resonances that have the WGM structure and satisfy suitable estimates. Our formulas are reminiscent of others obtained for eigenmodes with Dirichlet boundary conditions [2, 1]. Relying on general theory [12] we deduce that our quasi-resonances are asymptotically close to true resonances.

#### REFERENCES

- [1] S. Ancey, A. Folacci, and P. Gabrielli, *Whispering-gallery modes and resonances of an elliptic cavity*, Journal of Physics. A. Mathematical and General, 34 (2001), pp. 1341–1359.
- [2] V. M. Babič and V. S. Buldyrev, *Short-wavelength diffraction theory*, vol. 4 of Springer Series on Wave Phenomena, Springer-Verlag, Berlin, 1991.
- [3] S. Balac, M. Dauge and Z. Moitier, *Asymptotic expansions of whispering gallery modes in optical micro-cavities*, In preparation (2019).
- [4] J. Galkowski, *The quantum Sabine law for resonances in transmission problems*, Pure and Applied Analysis, 1 (2019), pp. 27–100.
- [5] J. Heebner, R. Grover, and T. Ibrahim, *Optical Microresonators: Theory, Fabrication, and Applications*, Optical Sciences, 138, Springer, 2008.
- [6] C. C. Lam, P. T. Leung, and K. Young, *Explicit asymptotic formulas for the positions, widths, and strengths of resonances in Mie scattering*, J. Opt. Soc. Am. B, 9 (1992), pp. 1585–1592.
- [7] A. Moiola and E. A. Spence, *Acoustic transmission problems: wavenumber-explicit bounds and resonance-free regions*, Math. Models Methods Appl. Sci., (2019) To appear.
- [8] F. Olver, D. Lozier, R. Boisvert, and C. Clark, *NIST Handbook of Mathematical Functions*, Cambridge University Press, 2010.
- [9] G. Popov and G. Vodev, *Resonances near the real axis for transparent obstacles*, Communications in Mathematical Physics, 207 (1999), pp. 411–438.
- [10] J. Sjöstrand and M. Zworski, *Asymptotic distribution of resonances for convex obstacles*, Acta Mathematica, 183 (1999), pp. 191–253.
- [11] E. I. Smotrova, A. I. Nosich, T. M. Benson, and P. Sewell, *Cold-cavity thresholds of microdisks with uniform and nonuniform gain: quasi-3-D modeling with accurate 2-D analysis*, IEEE Journal of Selected Topics in Quantum Electronics, 11 (2005), pp. 1135–1142.
- [12] S.-H. Tang and M. Zworski, *From quasimodes to resonances*, Mathematical Research Letters, 5 (1998), pp. 261–272.

### The HX Preconditioner in Higher Dimensions

JAY GOPALAKRISHNAN

(joint work with Martin Neumüller, Panayot Vassilevski)

In calculations using the  $n$ -dimensional finite element subcomplex of the de Rham complex, we often need a general purpose preconditioner for the stiffness matrix of the inner products. We report on the results obtained in [1] that generalize the construction and analysis of auxiliary space preconditioners, or HX preconditioners.

The first part of the talk focuses on developing simple computational proxies for  $k$ -forms and exterior derivatives in four dimensions (4D). For example, motivated by the fact the curl operator in two and three dimensions takes the form  $\text{curl } w = \varepsilon^{ij} \partial_i w_j$  and  $[\text{curl } w]_i = \varepsilon^{ijk} \partial_j w_k$ , respectively, where we have used the Levi-Civita tensor  $\varepsilon$  and the summation convention, we define curl in 4D by  $[\text{Curl } w]_{ij} = \varepsilon^{ijkl} \partial_k w_l$ . This produces a  $4 \times 4$  skew symmetric matrix, given a 4-vector  $w$ . Similar definitions yield simple 4D proxies, implementable using scalar, vector, and matrix algebraic structures that are usually already present in current software.

The second half of the talk focuses on the generalized HX preconditioners. The construction of these preconditioners is motivated using a generalization of a decomposition of Sobolev space functions into a regular component and a potential. A discrete version of such a decomposition can now be quickly established using the modern tools of finite element exterior calculus. Extensive numerical experiments in 4D illustrate the performance of the preconditioners, practical scalability, and parameter robustness, all in accordance with the theory.

#### REFERENCES

- [1] J. Gopalakrishnan, M. Neumüller, and P. S. Vassilevski, *The auxiliary space preconditioner for the de Rham complex*, SIAM J Numer Anal, 56 (2018), pp. 3196–3218.

### Generalized Plane Waves & Maxwell's equations

LISE-MARIE IMBERT-GÉRARD

(joint work with J.-F. Fritsch)

Modeling for wave propagation in magnetically confined plasma motivates the development of numerical methods for smooth variable coefficient time-harmonic Maxwell's equations. The simplest of these models, the cold plasma model, reads

$$\text{curl curl } \mathbf{E} - \left(\frac{\omega}{c}\right)^2 \epsilon \mathbf{E} = 0$$

where the tensor  $\epsilon$  is both homogeneous and anisotropic. Generalized Plane Waves (GPWs) are then introduced in the 2D variable refractive index Helmholtz framework [1, 2]. These functions are constructed to satisfy approximately the PDE, and a set of linearly independent can easily be constructed for discretization purposes. They are designed as exponential of polynomials, using Taylor expansions. The first extension of the GPW construction to the 3D vector-valued Maxwell's equation is introduced, specifying a particular ansatz for the amplitude and phase functions, and emphasizing the challenges related to the construction algorithm.

#### REFERENCES

- [1] L.-M. Imbert-Gérard, B. Després, *A generalized plane wave numerical method for smooth non constant coefficients*, IMA Journal of Numerical Analysis **34** (2014).  
 [2] L.-M. Imbert-Gérard, *Interpolation properties of generalized plane waves*, Numerische Mathematik **131**(4), (2015), pp.683–711.

## Finite Element Methods for Maxwell's Equations

PETER MONK

In the last few decades, conforming finite element methods for approximating the time harmonic Maxwell system governing electromagnetic wave propagation have undergone profound changes. Whereas in the 1980s there was confusion about how to choose the sesquilinear form and the appropriate finite elements to obtain a convergent solver, it is now clear not only how to discretize the Maxwell system using edge elements, but also how to analyze the resulting method [11, 13]. Important spin-offs from this analysis include the Finite Element Exterior Calculus [2] and the realization that the discrete de Rham diagram is a useful tool to guarantee conservation of charge, as well as for understanding the error analysis.

Of course edge elements are not the only possibility. But alternative approaches using continuous elements require special care to control the divergence of the solution and modification at discontinuities in material properties [6]. Problems facing the use of continuous elements were first recognized by Costabel and Dauge [8] who also proposed a fix for the problem (see also [6]). Recently a different approach has been proposed by Bonito and Guermond [5].

The simplest model problem for Maxwell's equations that captures several of the difficulties associated with their solution is as follows. Suppose  $\Omega \subset \mathbb{R}^3$  is a bounded polyhedral domain such that its boundary has two connected components  $\Gamma$  and  $\Sigma$  where  $\Sigma$  is the boundary of the unbounded connected complement of  $\Omega$ . Let  $\boldsymbol{\nu}$  denote the unit outward normal to  $\Omega$  and assume  $\Omega$  is simply connected. Then, given a suitable tangential vector field  $\mathbf{g} \in L^2(\Sigma)$  we seek  $\mathbf{E}$  such that

$$\begin{aligned} \nabla \times (\nabla \times \mathbf{E}) - \kappa^2 \varepsilon \mathbf{E} &= 0 \text{ in } \Omega \\ (\nabla \times \mathbf{E}) \times \boldsymbol{\nu} - i\kappa \mathbf{E}_T &= \mathbf{g} \text{ on } \Sigma, \\ \mathbf{E} \times \boldsymbol{\nu} &= 0 \text{ on } \Gamma. \end{aligned}$$

Here  $\mathbf{E}_T = (\boldsymbol{\nu} \times \mathbf{E}) \times \boldsymbol{\nu}$  is the tangential trace of  $\mathbf{E}$  and  $\kappa > 0$  is the constant wave-number of the radiation. The function  $\varepsilon$  denotes the electric permittivity of the material inside  $\Omega$  and is assumed here to be a real valued piecewise smooth coefficient which is uniformly positive and bounded.

The appropriate function space for the solution is built from the standard space

$$H(\text{curl}; \Omega) = \{\mathbf{u} \in (L^2(\Omega))^3 \mid \nabla \times \mathbf{u} \in (L^2(\Omega))^3\}$$

as follows

$$X = \{\mathbf{u} \in H(\text{curl}; \Omega) \mid \boldsymbol{\nu} \times \mathbf{u}|_{\Sigma} \in (L^2(\Sigma))^3, \boldsymbol{\nu} \times \mathbf{u} = 0 \text{ on } \Gamma\}$$

with norms

$$\begin{aligned} \|\mathbf{u}\|_{H(\text{curl}; \Omega)} &= \sqrt{\|\mathbf{u}\|_{(L^2(\Omega))^3}^2 + \|\nabla \times \mathbf{u}\|_{(L^2(\Omega))^3}^2} \\ \|\mathbf{u}\|_X &= \sqrt{\|\mathbf{u}\|_{H(\text{curl}; \Omega)}^2 + \|\boldsymbol{\nu} \times \mathbf{u}\|_{(L^2(\Sigma))^3}^2}. \end{aligned}$$

The associated inner products are

$$(\mathbf{u}, \mathbf{v}) = \int_{\Omega} \mathbf{u} \cdot \bar{\nabla} dV, \quad \langle \mathbf{u}, \mathbf{v} \rangle_{\Sigma} = \int_{\Sigma} \mathbf{u} \cdot \bar{\nabla} dA.$$

Following the usual Galerkin philosophy we obtain the variational problem of finding  $\mathbf{u} \in X$  such that

$$(\nabla \times \mathbf{E}, \nabla \times \phi) - \kappa^2(\varepsilon \mathbf{E}, \phi) - i\kappa \langle \mathbf{E}_T, \phi_T \rangle_{\Sigma} = \langle \mathbf{g}, \phi_T \rangle_{\Sigma}$$

for all  $\phi \in X$ . Using unique continuation it is possible to prove uniqueness of any solution to this problem. Then using the Helmholtz decomposition, the Webber compactness result, and the Fredholm Alternative existence of a solution can be obtained [3, 13]. The key here is that it is essential to use a Helmholtz decomposition in the analysis. This suggests that a successful finite element space should also possess a suitable discrete Helmholtz decomposition.

Let  $\mathcal{T}_h$  denote a conforming and regular tetrahedral grid covering  $\Omega$ . A suitable finite element discretization is provided by the Nédélec elements [14]. In particular the lowest order space is

$$X_h = \{ \mathbf{u}_h \in H(\text{curl}; \Omega) \mid \mathbf{u}_h|_K = \mathbf{a}_K + \mathbf{b}_K \times \mathbf{x} \\ \mathbf{a}_K, \mathbf{b}_K \in \mathbb{C}^3, \quad \forall K \in \mathcal{T}_h \}.$$

The degrees of freedom (unknowns) for this element are  $\int_e \mathbf{u}_h \cdot \boldsymbol{\tau}_h ds$  for each edge  $e$  of each tetrahedron where  $\boldsymbol{\tau}_e$  is an appropriately oriented tangent vector.

An important property of Nédélec's elements is that they contain many gradients. In the lowest order case, if

$$S_h = \{ p_h \in S \mid p_h|_K \in P_1, \quad \forall K \in \mathcal{T}_h, p = 0 \text{ on } \Gamma, p \text{ is constant on } \Sigma \},$$

then  $\nabla S_h \subset X_h$ . Furthermore the curl satisfies a Friedrich's type inequality on the orthogonal complement of  $\nabla S_h$  in  $X_h$  weighted by  $\epsilon$  [13].

Using the properties of edge finite element spaces: in particular a discrete analogue of compactness [12] and an extension theorem, Gatica and Meddahi [10, 13] prove that if  $h$  is small enough then there exists a unique finite element solution  $\mathbf{E}_h \in X_h$  and

$$\|\mathbf{E} - \mathbf{E}_h\|_{H(\text{curl}; \Omega)} \rightarrow 0 \text{ as } h \rightarrow 0.$$

For sufficiently smooth solutions, lowest order Nédélec elements give  $O(h)$  convergence. In fact it is very beneficial to use higher order elements when  $\kappa$  is large [9].

For other approaches to error estimates using mixed method or the regular decomposition see [4, 11].

More recently several open source implementations of common finite element families for the Maxwell system have appeared making edge elements much more accessible. An example is the NGSolve package [15] that contains edge elements of arbitrary order, and a sophisticated interface based on Python that allows the user easy access to the underlying components of the software. This allows the easy use of edge elements in applications, for example to photonic crystals [1], in which  $\epsilon$  varies markedly from place to place. This in turn raises interesting questions of approximation theory and the regularity of the solutions [7, 6].

## REFERENCES

- [1] Ammari, H. and Bao, G., *Maxwell's equations in periodic chiral structures*, Math. Nachr. **251** (2003) 3-18.
- [2] Arnold, D., Falk, R. and Winther, R. *Finite element exterior calculus: from Hodge theory to numerical stability*, Math. Comp. **47** (2010), 281-354.
- [3] Assous, F., Ciarlet, P., and Labrunie, S., *Mathematical foundations of computational electromagnetism*, Springer International Publishing (2018).
- [4] D. Boffi. *Finite element approximation of eigenvalue problems*. Acta Numerica, **19** (2010) 1-120.
- [5] Bonito, A., Guermond, J.-L., *Approximation of the eigenvalue problem for the time harmonic Maxwell system by continuous Lagrange finite elements*, Math. Comp. **80** (2011), 1887-1910.
- [6] Bonito, A., Guermond, J.-L., and Luddens, F., *An Interior Penalty Method with  $C^0$  Finite Elements for the Approximation of the Maxwell Equations in Heterogeneous Media: Convergence Analysis with Minimal Regularity*, Math. Modelling and Numer. Anal. (M2AN), **50** (2016) 1457-1489.
- [7] Ciarlet, P., *On the approximation of electromagnetic fields by edge finite elements. Part 1: Sharp interpolation results for low-regularity fields*, Computers & Mathematics with Applications, **71** (2016) 85-104.
- [8] Costabel, M., and Dauge, M., *Weighted regularization of Maxwell equations in polyhedral domains*, Numer. Math. **93** (2002) 239-277.
- [9] Demkowicz, L., *Computing with hp-Adaptive Finite Elements*, vol 1, CRC Press, 2006
- [10] Gatica, G.N. and Meddahi, S., *Finite element analysis of a time harmonic Maxwell problem with an impedance boundary condition*. IMA J. Numer. Anal., **32**, (2012) 534–52.
- [11] Hiptmair, R., *Finite elements in computational electromagnetism*, Cambridge University Press (2002) 237-339.
- [12] Kikuchi, F. *On a discrete compactness property for the Nédélec finite elements*. J. Fac. Sci. Univ. Tokyo, Sect. 1A Math., **36** (1989), 479–90.
- [13] Monk, P., *Finite Element Methods for Maxwell's Equation*, Oxford University Press (2003).
- [14] Nédélec, J.C., *Mixed finite elements in  $\mathbb{R}^3$* . Numer. Math., **35**, (1980) 315–41.
- [15] Schöberl, J., *NGSolve* available online at <https://ngsolve.org> (2019).

## The time-harmonic Maxwell equations with impedance boundary conditions in domains with a polyhedral or an analytic boundary

SERGE NICAISE

(joint work with Jérôme Tomezyk)

We are interested in properties of solutions of the Maxwell system with impedance boundary condition

$$(1) \quad \begin{cases} \operatorname{curl} \mathbf{E} - i\omega \mathbf{H} = \mathbf{0} & \text{and} & \operatorname{curl} \mathbf{H} + i\omega \mathbf{E} = \mathbf{J} & \text{in } \Omega, \\ \mathbf{H} \times \mathbf{n} - \lambda \mathbf{E}_t = \mathbf{0} & & & \text{on } \Gamma := \partial\Omega, \end{cases}$$

where  $\mathbf{E}$  is the electric part and  $\mathbf{H}$  is the magnetic part of the electromagnetic field,  $\omega > 0$  corresponds to the wave number or frequency. The right hand side  $\mathbf{J}$  is the current density which – in the absence of free electric charges – is divergence free, namely

$$\operatorname{div} \mathbf{J} = \mathbf{0} \quad \text{in } \Omega.$$

The impedance  $\lambda$  is supposed to be a smooth function satisfying

$$(2) \quad \lambda : \Gamma \rightarrow \mathbb{R}, \quad \text{such that} \quad \lambda(x) \neq 0, \forall x \in \partial\Omega,$$

see for instance [4, 3]. The case  $\lambda_{\text{imp}} \equiv 1$  is also called the Silver-Müller boundary condition [1].

Note that the boundary condition in (1) is an absorbing boundary condition that is used to reduce the full space into a bounded domain.

We present two variational formulations of this problem [2, §4.4.d] and analyse more carefully the second one that consists in keeping the full electromagnetic field  $(\mathbf{E}, \mathbf{H})$  as unknown, using the variational space

$$(3) \quad \mathbf{V} = \{(\mathbf{E}, \mathbf{H}) \in (\mathbf{H}(\text{curl}, \Omega) \cap \mathbf{H}(\text{div}, \Omega))^2 : \mathbf{H} \times \mathbf{n} = \lambda \mathbf{E}_{\mathbf{t}} \text{ on } \partial\Omega\},$$

considering the impedance condition as an essential boundary condition.

Hence the proposed variational formulation is: Find  $(\mathbf{E}, \mathbf{H}) \in \mathbf{V}$  such that

$$(4) \quad a(\mathbf{E}, \mathbf{H}; \mathbf{E}', \mathbf{H}') = \int_{\Omega} \left( i\omega \mathbf{J} \cdot \bar{\mathbf{E}}' + \mathbf{J} \cdot \text{curl} \bar{\mathbf{H}}' \right) dx, \forall (\mathbf{E}', \mathbf{H}') \in \mathbf{V},$$

with the choice

$$a(\mathbf{E}, \mathbf{H}; \mathbf{E}', \mathbf{H}') = a_{\omega, s}(\mathbf{E}, \mathbf{E}') + a_{\omega, s}(\mathbf{H}, \mathbf{H}') - i\omega \int_{\partial\Omega} (\lambda \mathbf{E}_{\mathbf{t}} \cdot \bar{\mathbf{E}}'_{\mathbf{t}} + \frac{1}{\lambda} \mathbf{H}_{\mathbf{t}} \cdot \bar{\mathbf{H}}'_{\mathbf{t}}) d\sigma,$$

with a positive real parameter  $s \in [1, 2]$  appropriately fixed and

$$a_{\omega, s}(\mathbf{u}, \mathbf{v}) = \int_{\Omega} (\text{curl} \mathbf{u} \cdot \text{curl} \bar{\mathbf{v}} + s \text{div} \mathbf{u} \text{div} \bar{\mathbf{v}} - \omega^2 \mathbf{u} \cdot \bar{\mathbf{v}}) dx.$$

The interest of this formulation stays on the fact that if  $\Omega$  has a  $C^2$  boundary or if it is a convex polyhedron, then  $\mathbf{V}$  is embedded in  $\mathbf{H}^1(\Omega) \times \mathbf{H}^1(\Omega)$  and  $a$  is (weakly) coercive on  $\mathbf{V}$ , see [2, §4.4.d] for the smooth case and [5] for the polyhedral case. Consequently problem (4) induces a Fredholm operator of index zero from  $\mathbf{V}$  into its dual and uniqueness implies existence and uniqueness.

The coerciveness of  $a$  implying that the corresponding system is an elliptic system in the Agmon-Douglis-Nirenberg sense, standard shift regularity results hold. Furthermore if the right-hand side  $\mathbf{J}$  belongs to  $\mathbf{H}(\text{div} = 0; \Omega)$  and  $-\omega^2/s$  is not an eigenvalue of the Laplace operator  $\Delta_{\text{Dir}}$ , then any solution  $(\mathbf{E}, \mathbf{H}) \in \mathbf{V}$  of the variational problem (4) satisfies (1).

Another advantage of this formulation concerns its numerical approximation. Indeed as  $\mathbf{V}$  is a subspace of  $\mathbf{H}^1(\Omega)^2$ , we can consider an  $hp$   $C^0$  finite element approximation, see [5] for polyhedral domains. But if the boundary is analytic, we cannot impose the impedance boundary condition in the finite element space. Hence we need to adopt a nonconforming approximation [6].

## REFERENCES

- [1] H. Barucq and B. Hanouzet. Etude asymptotique du système de Maxwell avec la condition aux limites absorbante de Silver-Müller II. *C. R. Acad. Sci. Paris Sér. I*, 316:1019–1024, 1993.
- [2] M. Costabel, M. Dauge, and S. Nicaise. *Corner Singularities and Analytic Regularity for Linear Elliptic Systems. Part I: Smooth domains*. <http://hal.archives-ouvertes.fr/hal-00453934/en/> Online version of Chapters 1 to 5. (2010).
- [3] P. Monk. *Finite element methods for Maxwell's equations*. Numerical Mathematics and Scientific Computation. Oxford University Press, 2003.
- [4] J.-C. Nédélec. *Acoustic and electromagnetic equations*, volume 144 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 2001.
- [5] S. Nicaise and J. Tomezyk. The time-harmonic Maxwell equations with impedance boundary conditions in polyhedral domains. *Maxwell's Equations: Analysis and Numerics*, U. Langer, D. Pauly and S. Repin eds, De Gruyter, 2019.
- [6] S. Nicaise and J. Tomezyk. Convergence analysis of a  $hp$ -finite element approximation of the time-harmonic Maxwell equations with impedance boundary conditions in domains with an analytic boundary. 2019, in preparation.

### Time-harmonic Scattering in exterior domains with mixed boundary conditions

FRANK OSTERBRINK

(joint work with Dirk Pauly)

Let  $\Omega \subset \mathbb{R}^3$  be an exterior weak Lipschitz domain with weak Lipschitz interface  $\partial\Omega = \bar{\Gamma}_1 \cup \bar{\Gamma}_2$  in the sense of [1, Definition 2.3, Definition 2.5]. We study the boundary value problem

$$(1) \quad \begin{aligned} -\operatorname{rot} H + i\omega\mu E &= -F & \text{in } \Omega, & & E \times \nu &= 0 & \text{on } \Gamma_1, \\ \operatorname{rot} E + i\omega\mu H &= G & \text{in } \Omega, & & H \times \nu &= 0 & \text{on } \Gamma_2, \end{aligned}$$

where  $\omega \in \mathbb{C} \setminus \{0\}$  and  $\varepsilon, \mu$  are symmetric, uniformly positive definite  $L^\infty$ -matrix fields, which are asymptotically a multiple of the identity, i.e.,  $\varepsilon = \varepsilon_0 \cdot \mathbb{1} + \hat{\varepsilon}$ ,  $\mu = \mu_0 \cdot \mathbb{1} + \hat{\mu}$  with  $\varepsilon_0, \mu_0 \in \mathbb{R}^+$  and

$$\hat{\varepsilon}, \hat{\mu} = \mathcal{O}(r^{-\kappa}) \quad \text{for } r \longrightarrow \infty, \quad \kappa > 1.$$

As shown in [5], working in the framework of polynomially weighted Sobolev spaces, the same methods as in [2] (see also [3]) are sufficient for solving system (1). In fact, the linear operator

$$\begin{aligned} \mathcal{M} &: \mathbf{R}_{\Gamma_1}(\Omega) \times \mathbf{R}_{\Gamma_2}(\Omega) \subset \mathbf{L}_\varepsilon^2(\Omega) \times \mathbf{L}_\mu^2(\Omega) &\longrightarrow & \mathbf{L}_\varepsilon^2(\Omega) \times \mathbf{L}_\mu^2(\Omega) \\ & & (E, H) & \longmapsto & (-i\varepsilon^{-1} \operatorname{rot} H, i\mu^{-1} \operatorname{rot} E) \end{aligned}$$

is selfadjoint, hence  $(\mathcal{M} - \omega)^{-1}$  is continuous for  $\omega \in \mathbb{C} \setminus \mathbb{R}$ . For  $\omega \in \mathbb{R} \setminus \{0\}$ , we are solving in the continuous spectrum of  $\mathcal{M}$  and therefore  $(\mathcal{M} - \omega)^{-1}$  exists only on a dense subset of  $\mathbf{L}_\varepsilon^2(\Omega) \times \mathbf{L}_\mu^2(\Omega)$ . However, using Eidus' limiting absorption

principle [4] we are still able to obtain weaker so called *radiating solutions*  $E, H$  by restricting ourselves to data  $F, G$  in

$$\mathbf{L}_s^2(\Omega) := \{ u \in \mathbf{L}_{\text{loc}}^2(\Omega) \mid (1 + r^2)^{s/2} u \in \mathbf{L}^2(\Omega) \}$$

for some  $s > 1/2$ . These solutions are then elements of

$$\mathbf{R}_t(\Omega) := \{ u \in \mathbf{L}_t^2(\Omega) \mid \text{rot } u \in \mathbf{L}_t^2(\Omega) \} \quad \forall t < -1/2$$

and satisfy the radiation condition

$$(\varepsilon_0 E - \sqrt{\varepsilon_0 \mu_0} \xi \times H, \mu_0 H + \sqrt{\varepsilon_0 \mu_0} \xi \times E) \in \mathbf{L}_s^2(\Omega) \times \mathbf{L}_s^2(\Omega), \quad \xi := x/r,$$

for some  $s > -1/2$ . In other words the resolvent  $(\mathcal{M} - \omega)^{-1}$  of  $\mathcal{M}$  may indeed be extended continuously to the real axis. The essential ingredients needed for the underlying limit process are the polynomial decay of eigensolutions, an a-priori estimate for solutions corresponding to non-real frequencies, a Helmholtz-type decomposition and *Weck's local selection theorem*, i.e.,

$$\mathbf{R}_{\Gamma_1}(\Omega) \cap \mathbf{D}_{\Gamma_2}(\Omega) \hookrightarrow \mathbf{L}_{\text{loc}}^2(\overline{\Omega}) \text{ is compact.}$$

While the first two are obtained by transferring well known results for the scalar Helmholtz equation to the time-harmonic Maxwell equations using a suitable decomposition of the fields  $E$  and  $H$  the last one is an assumption on the quality of the boundary, which holds true in weak Lipschitz domains.

## REFERENCES

- [1] S. Bauer, D. Pauly, and M. Schomburg, *The Maxwell Compactness Property in Bounded Weak Lipschitz Domains with Mixed Boundary Conditions*, SIAM J. Math. Anal. **48**(4), (2016), 2912 – 2943.
- [2] D. Pauly, *Low frequency asymptotics for time-harmonic generalized Maxwell equations in nonsmooth exterior domains*, Adv. Math. Sci. Appl. **16**(2), (2006), 591 – 622.
- [3] R. Picard, N. Weck, and K. J. Witsch, *Time-Harmonic Maxwell Equations in the Exterior of Perfectly Conducting, Irregular Obstacles*, Analysis (Munich) **21**(3), (2001), 231 – 264.
- [4] D. M. Eidus, *The principle of limiting absorption*, Amer. Math. Soc. Transl. Ser. 2 **47**, (1965), 157 – 191.
- [5] F. Osterbrink and D. Pauly, *Time-Harmonic Electro-Magnetic Scattering in Exterior Weak Lipschitz Domains with Mixed Boundary Conditions*, Maxwells Equations: Analysis and Numerics (Radon Series on Computational and Applied Mathematics), De Gruyter (2019), accepted.

## On Some Compact Embeddings in Various Hilbert Complexes

DIRK PAULY

(joint work with Sebastian Bauer, Michael Schomburg, and Walter Zulehner)

We study Hilbert complexes

$$(1) \quad \begin{aligned} \dots &\xrightarrow{\quad} D(A_0) \xrightarrow{A_0} D(A_1) \xrightarrow{A_1} H_2 \xrightarrow{\quad} \dots, \\ \dots &\xleftarrow{\quad} H_0 \xleftarrow{A_0^*} D(A_0^*) \xleftarrow{A_1^*} D(A_1^*) \xleftarrow{\quad} \dots \end{aligned}$$

for two densely defined and closed linear operators

$$A_0 : D(A_0) \subset H_0 \longrightarrow H_1, \quad A_1 : D(A_1) \subset H_1 \longrightarrow H_2$$

with Hilbert space adjoints  $A_0^*$  and  $A_1^*$ . The corresponding mathematical analysis, such as closed ranges, bounded inverses, Friedrichs/Poincaré type estimates, Helmholtz type decompositions, solution theories, finite cohomology groups, benefits strongly from a certain compact embedding, i.e.,

$$(2) \quad D(A_1) \cap D(A_0^*) \hookrightarrow H_1.$$

In the Hilbert complexes arising in mathematical physics a proper Sobolev setting with respective boundary conditions on bounded weak or strong Lipschitz domains is most important. In these cases a compact embedding (2) can be shown with the very elegant technique of regular potentials.

Classical examples are the de Rham complex

$$(3) \quad \begin{aligned} \{0\} &\xrightarrow{\iota_{\{0\}}} \mathring{H}^1 \xrightarrow{\mathring{\nabla}} \mathring{R} \xrightarrow{\mathring{\text{rot}}} \mathring{D} \xrightarrow{\mathring{\text{div}}} L^2 \xrightarrow{\pi_{\mathbb{R}}} \mathbb{R}, \\ \{0\} &\xleftarrow{\pi_{\{0\}}} L^2 \xleftarrow{-\mathring{\text{div}}} \mathring{D} \xleftarrow{\mathring{\text{rot}}} \mathring{R} \xleftarrow{-\mathring{\nabla}} \mathring{H}^1 \xleftarrow{\iota_{\mathbb{R}}} \mathbb{R}, \end{aligned}$$

which may also be generalised using alternating differential forms (even with mixed boundary conditions), the elasticity complex

$$(4) \quad \begin{aligned} \{0\} &\xrightarrow{\iota_{\{0\}}} \mathring{H}^1 \xrightarrow{\mathring{\text{sym}}\mathring{\nabla}} \mathring{R}\mathring{R}_{\mathbb{S}}^{\top} \xrightarrow{\mathring{\text{Rot}}\mathring{\text{Rot}}_{\mathbb{S}}^{\top}} \mathring{D}_{\mathbb{S}} \xrightarrow{\mathring{\text{Div}}_{\mathbb{S}}} L^2 \xrightarrow{\pi_{\text{RM}}} \text{RM}, \\ \{0\} &\xleftarrow{\pi_{\{0\}}} L^2 \xleftarrow{-\mathring{\text{Div}}_{\mathbb{S}}} \mathring{D}_{\mathbb{S}} \xleftarrow{\mathring{\text{Rot}}\mathring{\text{Rot}}_{\mathbb{S}}^{\top}} \mathring{R}\mathring{R}_{\mathbb{S}}^{\top} \xleftarrow{-\mathring{\text{sym}}\mathring{\nabla}} \mathring{H}^1 \xleftarrow{\iota_{\text{RM}}} \text{RM}, \end{aligned}$$

and the biharmonic (or  $\text{divDiv}$ ) complex

$$(5) \quad \begin{aligned} \{0\} &\xrightarrow{\iota_{\{0\}}} \mathring{H}^2 \xrightarrow{\mathring{\nabla}\mathring{\nabla}} \mathring{R}_{\mathbb{S}} \xrightarrow{\mathring{\text{Rot}}_{\mathbb{S}}} \mathring{D}_{\mathbb{T}} \xrightarrow{\mathring{\text{Div}}_{\mathbb{T}}} L^2 \xrightarrow{\pi_{\text{RT}}} \text{RT}, \\ \{0\} &\xleftarrow{\pi_{\{0\}}} L^2 \xleftarrow{\mathring{\text{div}}\mathring{\text{Div}}_{\mathbb{S}}} \mathring{D}\mathring{D}_{\mathbb{S}} \xleftarrow{\mathring{\text{sym}}\mathring{\text{Rot}}_{\mathbb{T}}} \mathring{R}_{\mathbb{T},\text{sym}} \xleftarrow{-\mathring{\text{dev}}\mathring{\nabla}} \mathring{H}^1 \xleftarrow{\iota_{\text{RT}}} \text{RT}. \end{aligned}$$

Typical compact embeddings (2) read as follows:

$$\begin{aligned} \text{in (3) : } & \mathring{H}^1 \hookrightarrow L^2, & \mathring{R} \cap \mathring{D} \hookrightarrow L^2, & \mathring{D} \cap \mathring{R} \hookrightarrow L^2, & \mathring{H}^1 \hookrightarrow L^2 \\ \text{in (4) : } & \mathring{H}^1 \hookrightarrow L^2, & \mathring{R}\mathring{R}_{\mathbb{S}}^{\top} \cap \mathring{D}_{\mathbb{S}} \hookrightarrow L_{\mathbb{S}}^2, & \mathring{D}_{\mathbb{S}} \cap \mathring{R}\mathring{R}_{\mathbb{S}}^{\top} \hookrightarrow L_{\mathbb{S}}^2, & \mathring{H}^1 \hookrightarrow L^2 \\ \text{in (5) : } & \mathring{H}^2 \hookrightarrow L^2, & \mathring{R}_{\mathbb{S}} \cap \mathring{D}\mathring{D}_{\mathbb{S}} \hookrightarrow L_{\mathbb{S}}^2, & \mathring{D}_{\mathbb{T}} \cap \mathring{R}_{\mathbb{T},\text{sym}} \hookrightarrow L_{\mathbb{T}}^2, & \mathring{H}^1 \hookrightarrow L^2 \end{aligned}$$

## REFERENCES

- [1] S. Bauer, D. Pauly, and M. Schomburg, *The Maxwell Compactness Property in Bounded Weak Lipschitz Domains with Mixed Boundary Conditions*, SIAM Journal on Mathematical Analysis **48**(4), (2016), 2912-2943.
- [2] D. Pauly, *Solution Theory, Variational Formulations, and Functional a Posteriori Error Estimates for General First Order Systems with Applications to Electro-Magneto-Statics and More*, Numerical Functional Analysis and Optimization (2019), accepted.
- [3] D. Pauly, W. Zulehner, *The divDiv-Complex and Applications to Biharmonic Equations*, Applicable Analysis (2019), accepted.
- [4] D. Pauly, *On the Maxwell and Friedrichs/Poincare Constants in ND*, Mathematische Zeitschrift (2019), accepted.
- [5] S. Bauer, D. Pauly, and M. Schomburg, *Weck's Selection Theorem: The Maxwell Compactness Property for Bounded Weak Lipschitz Domains with Mixed Boundary Conditions in Arbitrary Dimensions*, Maxwells Equations: Analysis and Numerics (Radon Series on Computational and Applied Mathematics), De Gruyter (2019), accepted.
- [6] D. Pauly, *A Global div-curl-Lemma for Mixed Boundary Conditions in Weak Lipschitz Domains and a Corresponding Generalized  $A_0^*$ - $A_1$ -Lemma in Hilbert Spaces*, Analysis (Munich) (2019), accepted.

**Nonconforming Trefftz virtual elements for the Helmholtz problem**

ILARIA PERUGIA

(joint work with Lorenzo Mascotto, Alexander Pichler)

The virtual element method (VEM) is a recent generalization of the finite element method to polytopal grids [1, 2]. The main feature of VEM is that test and trial spaces consist of functions that are not known in closed form, but that are solutions to local differential problems mimicking the target one. Despite this fact, the method is made fully computable by defining two tools, namely suitable mappings from local approximation spaces into spaces of known functions (typically polynomials), and suitable bilinear/sesquilinear stabilization forms.

I have presented a novel VEM for the two dimensional Helmholtz problem endowed with impedance boundary conditions, which was introduced in [3]. The local approximation spaces consist of Trefftz functions, namely, functions belonging to the kernel of the Helmholtz operator. The global trial and test spaces are not fully discontinuous, as in most Trefftz methods [4], but rather interelement continuity is imposed in a nonconforming fashion (à la Crouzeix-Raviart). Although their functions are only implicitly defined, as typical of the VEM framework, they contain discontinuous subspaces made of functions known in closed form and with good approximation properties (plane waves, in the considered case).

An essential ingredient in the implementation of the method is an edgewise orthogonalization-and-filtering process described in [5]. This process allows to dramatically reduce the number of basis functions without deteriorating the accuracy. It also have a positive effect on the conditioning of the overall method.

After carrying out an  $h$ -version error analysis of the method, I have presented numerical tests reported in [5], which *i*) demonstrate the  $h$ -version theoretical convergence rates; *ii*) show that, in case of smooth solutions, the  $p$ -version achieves

exponential convergence; *iii*) show that, in case of singular solutions, the *hp*-version on graded meshes achieves exponential convergence; *iv*) show that the new method is competitive with other plane wave-based methods in terms of number of degrees of freedom needed in order to achieve a given accuracy, and is able to reach a higher accuracy, before onset of instabilities.

## REFERENCES

- [1] L. Beirão da Veiga, F. Brezzi, A. Cangiani, G. Manzini, L.D. Marini, and A. Russo, *Basic principles of virtual element methods*, Math. Models Methods Appl. Sci., **23** (2013), 199-214.
- [2] L. Beirão da Veiga, F. Brezzi, L.D. Marini, and A. Russo, *The hitchhiker's guide to the virtual element method*, Math. Models Methods Appl. Sci., **24** (2014), 1541-1573.
- [3] L. Mascotto, I. Perugia, and A. Pichler, *A nonconforming Trefftz virtual element method for the Helmholtz problem*, arXiv:1805.05634 [math.NA].
- [4] R. Hiptmair, A. Moiola, and I. Perugia, A survey of Trefftz methods for the Helmholtz equation, in *Building bridges: connections and challenges in modern approaches to numerical partial differential equations*, pages 237-279, Springer, 2016.
- [5] L. Mascotto, I. Perugia, and A. Pichler, *A nonconforming Trefftz virtual element method for the Helmholtz problem: numerical aspects*, Comp. Meth. Appl. Mech. Engrg., **347** (2019), 445-476.

## Memoryless Evolutionary Problems as Dynamic Abstract Friedrichs Systems

RAINER PICARD

Following the setting presented in [2] many models of mathematical physics share a common form

$$\partial_t V + AU = F,$$

where  $\partial_t$  denotes time-differentiation and  $A$  is a maximal accretive linear operator. Indeed, in standard cases  $A$  is simply skew-selfadjoint in an underlying real Hilbert space  $H$ . The unknowns  $U, V$  are here linked by a so-called material law

$$V = \mathcal{M}U.$$

In a number of studies, see e.g. [1, 4, 5], it has been illustrated, that this simple framework is indeed suitable for a large number of complex applications including even time-delay problems and fractional time derivatives. A typical and simple case, which we shall focus on here is

$$(1) \quad \mathcal{M} = M_0 + \partial_t^{-1} M_1,$$

where  $M_0, M_1$  are continuous linear operators in  $H$ , in particular,  $M_0$  is selfadjoint. The operator  $\partial_t^{-1}$  appearing here is forward causal time-integration, which can be properly realized in a weighted Hilbert space

$$H_{\rho,0}(\mathbb{R}, H) := \left\{ f \in L^{2,\text{loc}}(\mathbb{R}, X) \mid \int_{\mathbb{R}} \langle f(t) | f(t) \rangle_H \exp(-2\rho t) dt < \infty \right\}$$

with the natural weighted inner product

$$(f, g) \mapsto \langle f | g \rangle_{\rho,0,0} := \int_{\mathbb{R}} \langle f(t) | g(t) \rangle_H \exp(-2\rho t) dt.$$

In this setting the time-derivative  $\partial_0$  is a normal, strictly positive operator, [2], [3, Chapter 6], indeed we have

$$(2) \quad \partial_t \geq \rho$$

in the sense that we have for the numerical range of  $\partial_t$  that

$$\langle U | \partial_t U \rangle_{\rho,0,0} \geq \rho \langle U | U \rangle_{\rho,0,0}$$

for all  $U$  in the domain  $\text{dom}(\partial_t)$  of  $\partial_t$ . This strict positivity carries over to a basic well-posedness constraint for material laws of the form (1) such that

$$(3) \quad \rho M_0 + M_1 \geq c_0 > 0$$

holds for some real number  $c_0$  and all sufficiently large positive  $\rho \in \mathbb{R}$ . Since in this case  $\partial_t M(\partial_t) = \partial_t M_0 + M_1$  is indeed a local operator in time, we speak of a memoryless material law.

For  $A$  skew-selfadjoint these systems can be transformed into operator equations of the form

$$1 + \mathcal{A},$$

where now  $\mathcal{A}$  is skew-selfadjoint. Since the classical Friedrichs systems can also be brought into this formal form, we speak of abstract Friedrichs systems. With the skew-selfadjointness of  $A$  playing a central role, we present a number of tools, useful for modeling concrete problems, to construct such operators. These tools as well as the utility of the general setting are illustrated with various applications from mathematical physics.

## REFERENCES

- [1] A. J. Mulholland, R. Picard, S. Trostorff, and M. Waurick. On well-posedness for some thermo-piezoelectric coupling models. *Mathematical Methods in the Applied Sciences*, 39(15):4375–4384, 2016. mma.3866.
- [2] R. Picard. A structural observation for linear material laws in classical mathematical physics. *Mathematical Methods in the Applied Sciences*, 32(14):1768–1803, 2009.
- [3] R. Picard and D. McGhee. *Partial differential equations. A unified Hilbert space approach.*, volume 55 of *De Gruyter Expositions in Mathematics*. de Gruyter Expositions in Mathematics 55. Berlin: de Gruyter. xviii, 469 p., 2011.
- [4] R. Picard, S. Trostorff, and M. Waurick. On evolutionary equations with material laws containing fractional integrals. *Mathematical Methods in the Applied Sciences*, 38(15):3141–3154, 2015.
- [5] R. Picard, S. Trostorff, and M. Waurick. On some models for elastic solids with microstructure. *ZAMM - Journal of Applied Mathematics and Mechanics / Zeitschrift für Angewandte Mathematik und Mechanik*, 95(7):664–689, 2015.

## Solving Elasticity in $\mathbf{H}(\text{curl})$

JOACHIM SCHÖBERL

(joint work with Astrid Pechstein, Michael Neunteufel, Philip Lederer)

Vector valued finite elements with tangential or normal continuity are well established in the electromagnetics community. We show how to solve the elasticity equation with tangentially continuous  $\mathbf{H}(\text{curl})$  finite elements, and why this make sense. The TDNNS mixed method proposed by A. Pechstein and J. Schöberl uses the  $\mathbf{H}(\text{curl})$  function space for the displacement, and  $\mathbf{H}(\text{div div})$  for the stresses. The canonical finite elements for the stress space are normal-normal continuous symmetric matrix valued elements. It is shown that these elements are robust for anisotropic meshes, and hand over to plate and shell models. A recent proposal by Gopalakrishnan, Lederer and Schöberl switches the continuity: The vector variable  $\mathbf{u}$  is normal-continuous, while the matrix variable  $\boldsymbol{\sigma}$  is now normal-tangential continuous, and non-symmetric. This method provides an exactly divergence vector field  $\mathbf{u}$ , and is thus useful for incompressible flows. We show how all these spaces fit into a complex similar to the de Rham complex.

### REFERENCES

- [1] J. Gopalakrishnan, P. L. Lederer, and J. Schöberl, *A mass conserving mixed stress formulation for the Stokes equations*, arXiv:1806.07173, (2018).
- [2] A. S. Pechstein and J. Schöberl, *An analysis of the TDNNS method using natural norms*, Numer. Math. **139**(1), (2018), 93-120.
- [3] A. S. Pechstein and J. Schöberl, *Tangential-Displacement and Normal-Normal-Stress Continuous Mixed Finite Elements for Elasticity*, Mathematical Models and Methods in Applied Sciences, **21**(8), (2011), 1761-1782.

## On the helicity of a bounded domain and the Biot–Savart operator

ALBERTO VALLI

Let  $\Omega \subset \mathbb{R}^3$  be a bounded domain, with Lipschitz boundary  $\partial\Omega$  and unit outward normal vector  $\mathbf{n}$  on  $\partial\Omega$ . Consider the Hilbert space

$$\mathcal{V} = \{\mathbf{v} \in (L^2(\Omega))^3 \mid \text{div } \mathbf{v} = 0 \text{ in } \Omega, \mathbf{v} \cdot \mathbf{n} = 0 \text{ on } \partial\Omega\}.$$

The helicity of a vector field  $\mathbf{v} \in \mathcal{V}$ , a concept introduced by Woltjer (1958) and named by Moffatt (1969), is given by

$$(1) \quad H(\mathbf{v}) = \frac{1}{4\pi} \int_{\Omega} \int_{\Omega} \mathbf{v}(\mathbf{x}) \times \mathbf{v}(\mathbf{y}) \cdot \frac{\mathbf{x} - \mathbf{y}}{|\mathbf{x} - \mathbf{y}|^3} d\mathbf{x} d\mathbf{y}.$$

It is a “measure of the extent to which the field lines wrap and coil around one another” (Cantarella et al. (2001)).

The Biot–Savart operator  $BS$  is defined in  $\mathcal{V}$  as

$$(2) \quad BS(\mathbf{v})(\mathbf{x}) = \frac{1}{4\pi} \int_{\Omega} \mathbf{v}(\mathbf{y}) \times \frac{\mathbf{x} - \mathbf{y}}{|\mathbf{x} - \mathbf{y}|^3} d\mathbf{y}.$$

Hence we can rewrite  $H(\mathbf{v}) = \int_{\Omega} \mathbf{v} \cdot BS(\mathbf{v})$ .

It is well-known that  $BS(\mathbf{v})$  satisfies in  $\Omega$  the relations  $\operatorname{curl} BS(\mathbf{v}) = \mathbf{v}$  and  $\operatorname{div} BS(\mathbf{v}) = 0$ . Let us define  $\widehat{BS}(\mathbf{v})$  the  $(L^2(\Omega))^3$ -orthogonal projection of  $BS(\mathbf{v})$  over  $\mathcal{V}$ ; since it differs from  $BS(\mathbf{v})$  by a gradient of a scalar function, it satisfies

$$(3) \quad \begin{cases} \operatorname{curl} \widehat{BS}(\mathbf{v}) = \mathbf{v} & \text{in } \Omega \\ \operatorname{div} \widehat{BS}(\mathbf{v}) = 0 & \text{in } \Omega \\ \widehat{BS}(\mathbf{v}) \cdot \mathbf{n} = 0 & \text{on } \partial\Omega, \end{cases}$$

and one also easily sees that  $H(\mathbf{v}) = \int_{\Omega} \mathbf{v} \cdot \widehat{BS}(\mathbf{v})$ .

In Valli (2019) it is shown that the couple  $(\widehat{BS}(\mathbf{v}), \mathbf{0})$  is the solution  $(\mathbf{u}, \mathbf{q}) \in \mathcal{Z} \times \mathcal{H}$  of the following constrained least-square problem:

$$(4) \quad \begin{aligned} \int_{\Omega} \operatorname{curl} \mathbf{u} \cdot \operatorname{curl} \mathbf{w} + \int_{\Omega} \mathbf{q} \cdot \mathbf{w} &= \int_{\Omega} \mathbf{v} \cdot \operatorname{curl} \mathbf{w} & \forall \mathbf{w} \in \mathcal{Z} \\ \int_{\Omega} \mathbf{u} \cdot \mathbf{p} &= 0 & \forall \mathbf{p} \in \mathcal{H}, \end{aligned}$$

where

$$\begin{aligned} \mathcal{X} &= \{ \mathbf{w} \in H(\operatorname{curl}; \Omega) \mid \operatorname{curl} \mathbf{w} \cdot \mathbf{n} = 0 \text{ on } \partial\Omega \}, \\ \mathcal{Z} &= \{ \mathbf{w} \in \mathcal{X} \mid \oint_{\gamma_j} \mathbf{w} \cdot \mathbf{t}_j = 0 \text{ for } j = 1, \dots, g \}, \\ \mathcal{H} &= \operatorname{grad} H^1(\Omega), \end{aligned}$$

and the closed curves  $\gamma_j \subset \partial\Omega$  are a basis of the first homology group of  $\overline{\Omega}$ .

It is known that  $\widehat{BS}$  is a self-adjoint and compact operator in  $\mathcal{V}$  (see, e.g., Cantarella et al. (2001)), thus its spectrum is discrete. Defining the helicity of a domain  $\Omega$  by

$$(5) \quad H_{\Omega} = \sup_{\mathbf{v} \in \mathcal{V} \setminus \mathbf{0}} \frac{|H(\mathbf{v})|}{\|\mathbf{v}\|_{L^2(\Omega)}^2},$$

we easily obtain the spectral representation

$$H_{\Omega} = |\lambda_{\max}^{\Omega}|,$$

where  $\lambda_{\max}^{\Omega}$  is the eigenvalue of  $\widehat{BS}$  in  $\Omega$  of maximum absolute value.

A finite element scheme for calculating the eigenvalues of  $\widehat{BS}$  has been proposed in Alonso Rodríguez et al. (2018), and permits to compute the helicity of  $\Omega$ .

## REFERENCES

- [1] A. Alonso Rodríguez, J. Camaño, R. Rodríguez, A. Valli and P. Venegas, *Finite element approximation of the spectrum of the curl operator in a multiply-connected domain*, *Found. Comput. Math.*, **18** (2018), 1493–1533.
- [2] J. Cantarella, D. DeTurck and H. Gluck, *The Biot–Savart operator for application to knot theory, fluid dynamics, and plasma physics*, *J. Math. Phys.*, **42** (2001), 876–905.
- [3] H.K. Moffatt, *The degree of knottedness of tangled vortex lines*, *J. Fluid Mech.*, **35** (1969), 117–129.
- [4] A. Valli, *A variational interpretation of the Biot–Savart operator and the helicity of a bounded domain*, *J. Math. Phys.*, **60** (2019), to appear.
- [5] L. Woltjer, *A theorem on force-free magnetic fields*, *Proc. Nat. Acad. Sci. U.S.A.*, **44** (1958), 489–491.

## On Nonlocal $H$ -convergence

MARCUS WAURICK

In classical homogenisation problems for elliptic-type equations, one studies coefficients

$$M(\alpha, \beta, \Omega) := \{a \in L^\infty(\Omega; \mathbb{R}^{d \times d}); \Re a(x) \geq \alpha, \Re a(x)^{-1} \geq 1/\beta \text{ (a.e. } x \in \Omega)\}$$

for some fixed  $0 < \alpha < \beta$  and  $\Omega \subseteq \mathbb{R}^d$  open and bounded. The inequalities are understood in the sense of positive definiteness. Then a sequence  $(a_n)_n$  in  $M(\alpha, \beta, \Omega)$  is said to *locally  $H$ -converge* to  $a \in M(\alpha, \beta, \Omega)$ , if for all  $f \in H^{-1}(\Omega)$  and corresponding solutions  $(u_n)_n$  in  $H_0^1(\Omega)$  of the Dirichlet problem

$$\langle a_n \text{grad } u_n, \text{grad } \phi \rangle = f(\phi) \quad (\phi \in H_0^1(\Omega))$$

we have that  $u_n \rightharpoonup u \in H_0^1(\Omega)$  and  $a_n \text{grad } u_n \rightharpoonup a \text{grad } u \in L^2(\Omega)^d$ , where  $u$  is the unique solution of

$$\langle a \text{grad } u, \text{grad } \phi \rangle = f(\phi) \quad (\phi \in H_0^1(\Omega)).$$

In this talk, we address the situation, where the condition on  $(a_n)_n$  to be  $L^\infty$ -matrices is relaxed to the extent that  $(a_n)_n$  is now only assumed to live in

$$\mathcal{M}(\alpha, \beta, \Omega) := \{a \in \mathcal{B}(L^2(\Omega)^d); \Re a \geq \alpha, \Re a^{-1} \geq 1/\beta\}.$$

It is relatively easy to see that the above definition has to be amended to be fitted also for this potentially nonlocal situation. In fact, for the definition of nonlocal  $H$ -convergence, one needs to introduce a second variational problem with the curl-operator (without boundary condition) replacing the above grad.

Having introduced the new notion of nonlocal  $H$ -convergence, we show that this convergence induces a compact, metrisable Hausdorff topology on (subsets of)  $\mathcal{M}(\alpha, \beta, \Omega)$ . Moreover, we show that nonlocal  $H$ -convergence coincides with local  $H$ -convergence if restricted to  $M(\alpha, \beta, \Omega)$ . It turns out that different choices for the boundary conditions for grad and curl lead to different topologies. Thus, nonlocal  $H$ -convergence ‘sees’ the boundary, a property not shared by local  $H$ -convergence. Furthermore, we provide a div-curl-type characterisation for nonlocal  $H$ -convergence, which facilitates the computation of nonlocal  $H$ -limits in practice. The results have applications to Maxwell's equations with nonlocal, that is, convolution-type material law relevant in nonlocal response theory for electromagnetics; see also [1, Chapter 10] and for the application part see [2] and the survey paper [4].

The talk is based on results in [3].

### REFERENCES

- [1] O. Keller. *Quantum Theory of Near-Field Electrodynamics*. Springer, Berlin Heidelberg, 2011.
- [2] M. Waurick. *On the homogenization of partial integro-differential-algebraic equations*. Operators and Matrices, 10(2):247–283, 2016.
- [3] M. Waurick. *Nonlocal  $H$ -convergence*. Calculus of Variations and Partial Differential Equations, 57(6):46, 2018.
- [4] M. Waurick *Homogenisation and the Weak Operator Topology*, arXiv:1901.06670, 2019.

## Geometric decompositions in finite element exterior calculus

RAGNAR WINTHER

(joint work with Richard S. Falk)

The theory for approximation of Hilbert complexes, presented in [2], explains the importance of bounded cochain projections for the construction of stable numerical methods for the Hodge Laplace problems. The purpose of this talk is to give a review of various constructions of such operators, and to discuss their properties.

The classical cochain projections are the canonical projections constructed from the degrees of freedom. However, these operators are not bounded in the desired Sobolev norms. Motivated by this observation we reviewed the construction of the so-called "smoothed projections," as it was presented in [1]. This gives  $L^2$  bounded projections, but in contrast to the canonical projections, they are nonlocal. We then followed up by presented the construction of local cochain projections given in [3]. In the final part of the talk we briefly reviewed some results for the bubble transform, see [4], which potentially can lead to cochain projections with bounds independent of the polynomial degree.

### REFERENCES

- [1] D.N. Arnold, R.S. Falk, and R. Winther, *Finite element exterior calculus, homological techniques and applications*, Acta Numerica **15**, (2006), 1 - 155.
- [2] D.N. Arnold, R.S. Falk, and R. Winther, *Finite element exterior calculus: from Hodge theory to numerical stability*, Bull. Amer. Math. Soc. **47**(2), (2010), 281 - 354.
- [3] R.S. Falk, and R. Winther, *Local bounded cochain projections*, Math. Comp. **83**(290), (2014), 2631 - 2656.
- [4] R.S. Falk, and R. Winther, *The bubble transform: A new tool for analysis of finite element methods*, Found. of Comp. Math. **16**(1), (2016), 1 - 32.

Reporter: Dirk Pauly