



Mathematisches
Forschungsinstitut
Oberwolfach

Member of the



Oberwolfach Preprints

OWP 2021 - 08

TIM DE LAAT AND SAFOURA ZADEH

Weak*-Continuity of Invariant Means on
Spaces of Matrix Coefficients

Mathematisches Forschungsinstitut Oberwolfach gGmbH
Oberwolfach Preprints (OWP) ISSN 1864-7596

Oberwolfach Preprints (OWP)

The MFO publishes a preprint series **Oberwolfach Preprints (OWP)**, ISSN 1864-7596, which mainly contains research results related to a longer stay in Oberwolfach, as a documentation of the research work done at the MFO. In particular, this concerns the Research in Pairs-Programme (RiP) and the Oberwolfach-Leibniz-Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

A preprint can have a size from 1 - 200 pages, and the MFO will publish it on its website as well as by hard copy. Every RiP group or Oberwolfach-Leibniz-Fellow may receive on request 20 free hard copies (DIN A4, black and white copy) by surface mail.

The full copyright is left to the authors. With the submission of a manuscript, the authors warrant that they are the creators of the work, including all graphics. The authors grant the MFO a perpetual, irrevocable, non-exclusive right to publish it on the MFO's institutional repository.

In case of interest, please send a **pdf file** of your preprint by email to rip@mfo.de or owlf@mfo.de, respectively. The file should be sent to the MFO within 12 months after your stay as RiP or OWLF at the MFO.

There are no requirements for the format of the preprint, except that the introduction should contain a short appreciation and that the paper size (respectively format) should be DIN A4, "letter" or "article".

On the front page of the hard copies, which contains the logo of the MFO, title and authors, we shall add a running number (20XX - XX). Additionally, each preprint will get a Digital Object Identifier (DOI).

We cordially invite the researchers within the RiP or OWLF programme to make use of this offer and would like to thank you in advance for your cooperation.

Imprint:

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany

Tel +49 7834 979 50
Fax +49 7834 979 55
Email admin@mfo.de
URL www.mfo.de

The Oberwolfach Preprints (OWP, ISSN 1864-7596) are published by the MFO.
Copyright of the content is held by the authors.

DOI 10.14760/OWP-2021-08

WEAK*-CONTINUITY OF INVARIANT MEANS ON SPACES OF MATRIX COEFFICIENTS

TIM DE LAAT AND SAFOURA ZADEH

ABSTRACT. With every locally compact group G , one can associate several interesting bi-invariant subspaces $X(G)$ of the weakly almost periodic functions $\text{WAP}(G)$ on G , each of which captures parts of the representation theory of G . Under certain natural assumptions, such a space $X(G)$ carries a unique invariant mean and has a natural predual, and we view the weak*-continuity of this mean as a rigidity property of G . Important examples of such spaces $X(G)$, which we study explicitly, are the algebra $M_{\text{cb}}A_p(G)$ of p -completely bounded multipliers of the Figà-Talamanca–Herz algebra $A_p(G)$ and the p -Fourier-Stieltjes algebra $B_p(G)$. In the setting of connected Lie groups G , we relate the weak*-continuity of the mean on these spaces to structural properties of G . Our results generalise results of Bekka, Kaniuth, Lau and Schlichting.

1. INTRODUCTION

With every locally compact group G , which in this article we assume to be second countable and Hausdorff, one can associate several interesting function spaces consisting of matrix coefficients of different classes of representations. In the setting of unitary representations, notable examples, which were introduced by Eymard [14], are the Fourier-Stieltjes algebra $B(G)$, consisting of all matrix coefficients of unitary representations of G , and the Fourier algebra $A(G)$, consisting of all matrix coefficients of the left-regular representation of G .

A much more general class of representations is the class of uniformly bounded representations of G on reflexive Banach spaces. Albeit being a very large class, these representations still have nice analytic properties. For instance, if $\pi: G \rightarrow \mathcal{B}(E)$ is such a representation, then E decomposes as the direct sum of the closed subspace of $\pi(G)$ -invariant vectors and a canonical invariant complement [2] (see also [5], [29]). The algebra of matrix coefficients of such representations coincides with the algebra $\text{WAP}(G)$ of weakly almost periodic functions on G [24] (see also [22], [4]).

As a well-known consequence of the Ryll-Nardzewski fixed point theorem, the space $\text{WAP}(G)$ carries a unique (two-sided) invariant mean (see e.g. [16,

TdL is supported by the Deutsche Forschungsgemeinschaft – Project-ID 427320536 – SFB 1442, as well as under Germany’s Excellence Strategy – EXC 2044 – 390685587, Mathematics Münster: Dynamics – Geometry – Structure.

SZ is supported through the “Oberwolfach Leibniz Fellows” program – Project-ID 2105q, by Mathematisches Forschungsinstitut Oberwolfach in 2021.

§3.1]). In [17, Theorem 3.6], Haagerup, Knudby and the first-named author showed that under mild conditions, a bi-invariant linear subspace $X(G)$ of $\text{WAP}(G)$ has a unique invariant mean, arising as the restriction of the mean on $\text{WAP}(G)$. A particular case of this result, namely the case $B(G)$, was already known from [15]. We recall some background on weakly almost periodic functions and invariant means in Section 2.

The Fourier-Stieltjes algebra $B(G)$ can be identified with the dual space of the universal group C^* -algebra $C^*(G)$ of G . It is known that the mean on $B(G)$ is weak*-continuous if and only if G has Kazhdan's property (T) [1], [30] (see also [17]), which is a well-known rigidity property for groups with many applications (see [6]).

This characterisation of property (T) raises the question under which conditions a space $X(G)$ as above has a predual. We establish sufficient conditions for this in Theorem 3.1 and describe an explicit predual $Y(G)$ for these spaces. Under the conditions that a (sufficiently large) function space $X(G)$ has a (unique) invariant mean m_X and a natural predual $Y(G)$, we can study the analogue of property (T) corresponding to the weak*-continuity of m_X . More precisely, we say that G has property (T_X^*) if the invariant mean m_X on $X(G)$ is weak*-continuous with respect to the weak*-topology coming from the predual $Y(G)$ (see also Definition 4.1). This idea generalises both property (T), corresponding to $X(G) = B(G)$, and property (T^*) from [17], corresponding to $X(G)$ being the space $M_{\text{cb}}A(G)$ of completely bounded Fourier multipliers.

We study property (T_X^*) explicitly in the concrete cases of $X(G)$ being the algebra $M_{\text{cb}}A_p(G)$ of p -completely bounded multipliers of the Figà-Talamanca–Herz algebra $A_p(G)$, and $X(G)$ being the p -Fourier-Stieltjes algebra $B_p(G)$, with $1 < p < \infty$. We recall these algebras in Section 2. For these $X(G)$ and connected Lie groups G , we relate property (T_X^*) to the structure of G ; see Theorem 7.3 and Theorem 8.2. In these theorems, the space $X(G) \cap C_0(G)$ is of special interest. Indeed, as can be expected, under certain natural assumptions on G , property (T_X^*) is equivalent to the fact that $X(G) \cap C_0(G)$ is weak*-closed in $X(G)$. This follows directly from the weak*-continuity of the mean in combination with a very general Howe-Moore type theorem due to Veech [32]. Note that the space $X(G) \cap C_0(G)$ can be viewed as a generalisation of the Rajchman algebra $B(G) \cap C_0(G)$.

Theorem 8.2 generalises a result of Bekka, Kaniuth, Lau and Schlichting [7, Theorem 2.7], where for connected Lie groups G , the weak*-closedness of the space $B(G) \cap C_0(G)$ in $B(G)$ was related to the structure of G . Let us point out that the characterisation of property (T) in terms of the weak*-continuity of the mean on $B(G)$ was not used there.

Our initial aim was to prove results along the lines of Theorem 7.3 and Theorem 8.2 for more abstract classes of function spaces/algebras (of matrix coefficients) that carry an invariant mean and have a natural predual. However, in order to relate the weak*-continuity of the mean to the structure of

groups, we would have needed to put too many additional assumptions on $X(G)$, which would not have justified the level of abstraction.

ACKNOWLEDGEMENTS

The work leading to this paper started whilst the second-named author was visiting the Westfälische Wilhelms-Universität Münster. She would like to gratefully thank the WWU Münster for the hospitality and excellent working conditions during the visit.

2. PRELIMINARIES

2.1. Invariant means. Let G be a locally compact group and $X(G)$ a linear subspace of $L^\infty(G)$ that is closed under complex conjugation and that contains all constant functions. A positive linear functional $m: X(G) \rightarrow \mathbb{C}$ satisfying $\|m\| = m(1) = 1$ is said to be a mean on $X(G)$.

We assume that $X(G)$ is invariant under left and right translations, i.e. if $\varphi \in X(G)$, then for all $g \in G$, the functions $L_g\varphi: h \mapsto \varphi(g^{-1}h)$ and $R_g\varphi: h \mapsto \varphi(hg)$ lie in $X(G)$. A mean m is called left invariant (resp. right invariant) if $m(L_g\varphi) = m(\varphi)$ (resp. $m(R_g\varphi) = m(\varphi)$) for all $\varphi \in X(G)$ and $g \in G$. A mean that is both left and right invariant is called two-sided invariant. In this article, invariant means are always assumed to be two-sided invariant, unless explicitly mentioned otherwise.

2.2. Weakly almost periodic functions. For a locally compact group G , a function $\varphi \in C_b(G)$ is called weakly almost periodic if the left orbit $O(\varphi, L) = \{L_g\varphi \mid g \in G\}$ (or equivalently, the right orbit $O(\varphi, R) = \{R_g\varphi \mid g \in G\}$) is relatively weakly compact, i.e. its closure is compact in the weak topology on $C_b(G)$. We denote the space of weakly almost periodic functions by $\text{WAP}(G)$. This space is a closed bi-invariant and inversion invariant subalgebra of $C_b(G)$ that contains the constants.

It is well known (see e.g. [16, §3.1]) that for every locally compact group G , there exists a unique left invariant mean m on $\text{WAP}(G)$. Given an element $\varphi \in \text{WAP}(G)$, its mean $m(\varphi)$ is explicitly given by the unique constant in the weakly closed convex hull $C(\varphi, L)$ of $O(\varphi, L)$ (which equals the unique constant in the analogously defined set $C(\varphi, R)$). Moreover, the mean m is right invariant and inversion invariant. For details on weakly almost periodic functions, see [16], [9].

2.3. An invariant mean on subspaces of $\text{WAP}(G)$. It is known that certain classes of subspaces of $\text{WAP}(G)$ also carry a unique invariant mean. The following result is [17, Theorem 3.6].

Theorem 2.1. Let G be a locally compact group and $X(G)$ a linear subspace of $\text{WAP}(G)$ that is closed under left translations and conjugation and that contains the constants. Then $X(G)$ carries a unique left invariant mean m_X , which is in fact the restriction of the mean on $\text{WAP}(G)$. If additionally,

$X(G)$ is closed under right translations, then m_X is right invariant as well. Moreover, if $X(G)$ is closed under inversion, then m_X is also invariant under inversion.

2.4. The space $M_{\text{cb}}A_p(G)$. For $1 < p < \infty$, we denote by $M_{\text{cb}}A_p(G)$ the space of p -completely bounded multipliers of the Figà-Talamanca–Herz algebra $A_p(G)$ of G . We mainly work with the following characterisation of p -completely bounded multipliers from [13]. Let QSL^p denote the class of quotients of closed subspaces (or, equivalently, closed subspaces of quotients) of L^p -spaces.

Proposition 2.2 ([13, Theorem 8.3]). Let G be a locally compact group and $1 < p < \infty$. A function $\varphi: G \rightarrow \mathbb{C}$ is a p -completely bounded multiplier of $A_p(G)$ if there exists a space $E \in QSL^p$ and bounded, continuous maps $\alpha: G \rightarrow E$ and $\beta: G \rightarrow E^*$ such that for all $g, h \in G$,

$$(1) \quad \varphi(hg^{-1}) = \langle \beta(h), \alpha(g) \rangle.$$

Given $\varphi \in M_{\text{cb}}A_p(G)$, its norm $\|\varphi\|_{M_{\text{cb}}A_p(G)}$ is given by the infimum of the numbers $\|\alpha\|_\infty \|\beta\|_\infty$ over all choices of E , α and β for which (1) holds. The space $M_{\text{cb}}A_2(G)$ corresponds with the usual completely bounded multipliers of the Fourier algebra $A(G)$.

It is known that for $1 < p < \infty$, the space $M_{\text{cb}}A_p(G)$ is a subalgebra of $\text{WAP}(G)$ [34] (see also [17, Proposition 3.3] for an explicit proof in the case $p = 2$). Also, for $1 < p \leq q \leq 2$ or $2 \leq q \leq p < \infty$, the algebra $M_{\text{cb}}A_q(G)$ embeds contractively into $M_{\text{cb}}A_p(G)$ [3, Proposition 6.1].

In general, a Fourier multiplier that is not completely bounded is not necessarily a weakly almost periodic function (see [8]), which is why we cannot consider the space $MA_p(G)$ of such multipliers in the setting of this article.

2.5. The space $B_p(G)$. For a locally compact group G and $1 < p < \infty$, let $\text{Rep}_p(G)$ denote the collection of all (isometric equivalence classes of) isometric representations of G on a QSL^p -space. Examples of elements in $\text{Rep}_p(G)$ are the trivial representation (on every QSL^p -space) and the left-regular representation $\lambda_p: G \rightarrow \mathcal{B}(L^p(G))$.

Let $1 < p < \infty$. The p -Fourier-Stieltjes algebra $B_p(G)$ is defined as the space of matrix coefficients of (isometric equivalence classes of) isometric representations on a QSL^p -space, i.e. functions of the form

$$(2) \quad g \mapsto \langle \eta, \pi(g)\xi \rangle,$$

where $\pi: G \rightarrow \mathcal{B}(E)$ is a representation in $\text{Rep}_p(G)$ and $\xi \in E$, $\eta \in E^*$. The space $B_p(G)$ carries a natural norm given by the infimum of the numbers $\|\xi\| \|\eta\|$ over all representations π in $\text{Rep}_p(G)$ and $\xi \in E$, $\eta \in E^*$ such that (2) holds.

This definition of $B_p(G)$ is due to Runde [28], who showed that $B_p(G)$ is a Banach algebra. However, the conventions used above are slightly different, in order to be better suitable to the purposes of this article.

It is known that for $1 < p \leq q \leq 2$ or $2 \leq q \leq p < \infty$, the algebra $B_q(G)$ embeds contractively into $B_p(G)$ [28].

For fixed p , we have the following norm-decreasing inclusions among the aforementioned spaces:

$$A_p(G) \subset B_p(G) \subset M_{\text{cb}}A_p(G) \subset \text{WAP}(G) \subset L^\infty(G).$$

3. A PREDUAL OF $X(G)$

Let G be a locally compact group, and let $\text{WAP}(G)$ be the algebra of weakly almost periodic functions on G . In what follows, let $X(G)$ be a bi-invariant linear subspace of $\text{WAP}(G)$ that is closed under complex conjugation and that contains the constants. Suppose that $X(G)$ has a natural norm $\|\cdot\|_X$, that the Fourier-Stieltjes algebra $B(G)$ embeds contractively into $X(G)$ and that $\|\cdot\|_\infty \leq \|\cdot\|_X \leq \|\cdot\|_B$. The assumption that $B(G)$ is contained in $X(G)$ guarantees that $X(G)$ “contains the unitary representation theory of G ” and that it is sufficiently large for our purposes. In particular, $X(G)$ is $\sigma(L^\infty(G), L^1(G))$ -dense in $L^\infty(G)$ (i.e. weak*-dense with respect to the weak*-topology on $L^\infty(G)$).

For $f \in L^1(G)$, set

$$\|f\|_Y := \sup \left\{ \left| \int_G f(g)\varphi(g)dg \right| \mid \varphi \in X(G), \|\varphi\|_X \leq 1 \right\}.$$

Then $\|\cdot\|_Y$ defines a norm on $L^1(G)$, and we denote by $Y(G)$ the completion of $L^1(G)$ with respect to $\|\cdot\|_Y$. The following result provides a criterion for $X(G)$ to have a predual.

Theorem 3.1. Let $X(G)$ be as above. Then the following are equivalent:

- (i) The spaces $X(G)$ and $Y(G)^*$ are isometrically isomorphic.
- (ii) The unit ball $(X(G))_1$ of $X(G)$ is $\sigma(L^\infty(G), L^1(G))$ -closed.

Moreover, in this case, every bounded linear functional $\alpha: Y(G) \rightarrow \mathbb{C}$ is of the form

$$(3) \quad \alpha(f) = \int_G f(g)\varphi(g)dg, \quad f \in L^1(G),$$

for some $\varphi \in X(G)$, and $\|\alpha\| = \|\varphi\|_X$.

Proof. (i) \implies (ii): Suppose that $Y(G)^*$ and $X(G)$ are isometrically isomorphic. There is a contractive map $\iota: L^1(G) \rightarrow Y(G)$. Its adjoint $\iota^*: X(G) \rightarrow L^\infty(G)$ is the inclusion map, which is weak*-weak*-continuous. Hence, it maps the unit ball of $X(G)$ to a weak*-compact subset of $L^\infty(G)$.

(ii) \implies (i): First note that for every $\varphi \in X(G)$, the linear functional

$$\alpha_\varphi: L^1(G) \rightarrow \mathbb{C}, \quad f \mapsto \int_G f(g)\varphi(g)dg$$

uniquely extends to a bounded linear functional on $Y(G)$, and we have that $\|\alpha_\varphi\| \leq \|\varphi\|_X$. Let $\Psi: X(G) \rightarrow Y(G)^*$ denote the contractive linear map given by $\Psi(\varphi) = \alpha_\varphi$. We show that Ψ is, in fact, a surjective isometry.

Let $\alpha \in Y(G)^*$ with $\|\alpha\| = 1$. Since $X(G)$ embeds contractively into $L^\infty(G)$, we observe that for every $f \in L^1(G)$, we have $\|f\|_Y \leq \|f\|_1$, and therefore, $\alpha' := \alpha|_{L^1(G)}$ is a bounded linear functional on $L^1(G)$. Hence, there exists a $\varphi \in L^\infty(G)$ such that

$$\alpha'(f) = \int_G f(g)\varphi(g)dg, \quad f \in L^1(G).$$

We now show that $\varphi \in (X(G))_1$, by means of a version of the bipolar theorem. We consider $(X(G))_1$ as a subset of $L^\infty(G)$ equipped with the $\sigma(L^\infty(G), L^1(G))$ -topology. By definition of $\|\cdot\|_Y$, we observe that the prepolar ${}^\circ(X(G))_1$ of $(X(G))_1$ is given by

$${}^\circ(X(G))_1 = \{f \in L^1(G) \mid \|f\|_Y \leq 1\}.$$

Since $\|\alpha\| = 1$, we have that for every $f \in {}^\circ(X(G))_1$,

$$\left| \int f(g)\varphi(g)dg \right| = |\alpha'(f)| = |\alpha(f)| \leq 1.$$

Therefore, φ belongs to the polar of ${}^\circ(X(G))_1$. Since $(X(G))_1$ is a convex balanced subset of $L^\infty(G)$ that is $\sigma(L^\infty(G), L^1(G))$ -closed by assumption, the polar of ${}^\circ(X(G))_1$ coincides with $(X(G))_1$ by the bipolar theorem (see [11, Corollary V.1.9]), so φ belongs to $(X(G))_1$. Moreover, $\Psi(\varphi) = \alpha$ and $\|\Psi(\varphi)\| \leq \|\varphi\|_X \leq 1 = \|\alpha\|$. This implies that Ψ is a surjective isometry, as desired. \square

Remark 3.2.

- (i) Note that in Theorem 3.1, the existence of a predual does not come for free. For example, $WAP(G)$ itself does not have a predual in general. Indeed, the algebra $WAP(G)$ is a C^* -algebra. If it would have a predual, then by Sakai's theorem, it would be a von Neumann algebra, which is in general not the case.
- (ii) A subspace $X(G)$ as in Theorem 3.1 may have several preduals that are not isometrically isomorphic. An easy example of this is the Fourier-Stieltjes algebra $B(\mathbb{T})$ of the circle group \mathbb{T} , which in fact coincides with the completely bounded Fourier multipliers $M_{cb}A(\mathbb{T})$. It is known that

$$B(\mathbb{T}) \cong C^*(\mathbb{T})^* \cong c_0(\mathbb{Z})^* \cong \ell^1(\mathbb{Z}).$$

The space $\ell^1(\mathbb{Z})$ is known to have many preduals different from $c_0(\mathbb{Z})$.

- (iii) In the case that $X(G)$ is $M_{cb}A(G)$, the predual from Theorem 3.1 coincides with the predual described for this space in [10, Proposition 1.10]. Miao generalised this to the p -completely bounded multipliers $M_{cb}A_p(G)$ [25], [26], [27]. Indeed, he proved that the space $Q_{p,cb}(G)$ defined as the completion of $L^1(G)$ with respect to the norm

$$\|f\|_{Q_{p,cb}} = \sup \left\{ \left| \int_G f(g)\varphi(g)dg \right| \mid \varphi \in (M_{cb}A_p(G))_1 \right\}$$

is a predual of $M_{cb}A_p(G)$.

As a matter of fact, the predual of $M_{\text{cb}}A_p(G)$ already occurred earlier in the literature. To see this, note that in [13, Theorem 8.6], Daws establishes an isometric isomorphism

$$(4) \quad HS_p(G) \cong M_{\text{cb}}A_p(G),$$

where $HS_p(G)$ is the Banach space of “ p -Herz-Schur multipliers” as defined by Herz in [21]. (In the notation of [21], however, the space $HS_p(G)$ is denoted by $B_p(G)$.) For the space $HS_p(G)$, Herz had already constructed a predual. Indeed, in [21], Herz introduces a Banach space $QF_p(G)$ and a contractive map $Q: L^1(G) \rightarrow QF_p(G)$ with dense range [21, Proposition 1]. Then, in [21, Proposition 2], he shows that $HS_p(G)$ coincides with the range of the adjoint map Q^* , and that Q^* induces an isometric isomorphism

$$HS_p(G) \cong QF_p(G)^*.$$

Combined with (4), this yields the desired predual.

- (iv) For $B_p(G)$, the predual of Theorem 3.1 coincides with the predual of this space described in [28, Theorem 6.6].

4. PROPERTY (T_X^*)

In this section, we study rigidity properties formulated in terms of the weak*-continuity of invariant means on appropriate function spaces of G .

Let G be a locally compact group, and let $X(G)$ be as in Theorem 3.1. In particular, $X(G)$ has a unique invariant mean m_X and the space $Y(G)$ as defined in Theorem 3.1 is a predual of $X(G)$.

Definition 4.1. A locally compact group G has property (T_X^*) if m_X is $\sigma(X(G), Y(G))$ -continuous.

When it is clear which space $X(G)$ we consider, we often just use the terminology “weak*-topology” instead of $\sigma(X(G), Y(G))$ -topology. Note that this topology depends on the chosen predual $Y(G)$. Unless explicitly stated otherwise, we always consider the natural predual from Theorem 3.1.

As mentioned in Section 1, for $X(G) = B(G)$, this property corresponds to Kazhdan’s property (T) (see [1], [30]), and for $X(G) = M_{\text{cb}}A(G)$, this property corresponds to property (T^*) of Haagerup, Knudby and the first-named author (see [17]).

The following proposition shows that if $X(G)$ is as in Theorem 3.1 (in particular we assume that $X(G)$ “includes the unitary representation theory” of G , in the sense that $B(G)$ embeds contractively into $X(G)$), then property (T_X^*) is a strengthening of property (T).

Proposition 4.2. Let $X(G)$ be a subspace of $WAP(G)$ satisfying the conditions of Theorem 3.1. If G has property (T_X^*) , then G has property (T).

Proof. The identity map $\text{id}: L^1(G) \rightarrow L^1(G)$ extends to a linear contraction $Y(G) \rightarrow C^*(G)$. Its adjoint is the inclusion map $\iota: B(G) \rightarrow X(G)$, and the map ι is weak*-weak*-continuous.

Suppose that the mean m_X on $X(G)$ is $\sigma(X(G), Y(G))$ -continuous. Since the mean m_B on $B(G)$ coincides with the composition $m_X \circ \iota$, the mean m_B is $\sigma(B(G), C^*(G))$ -continuous. \square

Remark 4.3. By the results recalled in Section 2, we know that for every $1 < p < \infty$, the space $B(G)$ embeds contractively into $B_p(G)$, and $B_p(G)$ embeds contractively into $M_{\text{cb}}A_p(G)$. Hence, the assumptions of Proposition 4.2 hold for the spaces $X(G) = B_p(G)$ and $X(G) = M_{\text{cb}}A_p(G)$.

It is clear that if $X(G)$ is as in Theorem 3.1 and G is a compact group, then G has property (T_X^*) . Indeed, the map $\varphi \mapsto \langle \varphi, 1 \rangle$ defines the weak*-continuous unique invariant mean on $X(G)$.

In case G is a non-compact group, it is in general difficult to show that the invariant mean on $X(G)$ is weak*-continuous. Indeed, establishing (a strengthening) of property (T) for a group G is usually hard. We study specific cases of $X(G)$ in the setting of Lie groups G in the next sections.

5. THE MEAN ON $M_{\text{cb}}A_p(G)$ AND PROPERTY $(T_{M_{\text{cb}}A_p}^*)$

In this section, we consider the case of $X(G)$ being the space $M_{\text{cb}}A_p(G)$ of p -completely bounded multipliers of $A_p(G)$ (with $1 < p < \infty$), and we study property $(T_{M_{\text{cb}}A_p}^*)$ and its permanence properties. We write $Q_{p,\text{cb}}(G)$ for the predual of $M_{\text{cb}}A_p(G)$ as described in Theorem 3.1. Some of the arguments in this section are modifications of results from [17].

The following generalises [17, Lemma 5.8]. The proof follows mutatis mutandis from the proof given there.

Lemma 5.1. Let G and H be locally compact groups and $\rho: H \rightarrow G$ a continuous group homomorphism, and let $1 < p < \infty$. If $u \in C_c(G)$ is a non-negative function with $\|u\|_1 = 1$ and $u^* = u$ (where $u^*(g) = \overline{u(g^{-1})}\Delta(g^{-1})$), then the linear map $T: M_{\text{cb}}A_p(G) \rightarrow M_{\text{cb}}A_p(H)$ defined by $\varphi \mapsto (u * \varphi) \circ \rho$ is weak*-weak*-continuous.

Proposition 5.2. Let $\rho: H \rightarrow G$ be a continuous group homomorphism with dense image, and let $1 < p < \infty$. If H has property $(T_{M_{\text{cb}}A_p}^*)$, then so has G .

Proof. Let $u \in C_c(G)$ be a non-negative function with $\|u\|_1 = 1$ and $u^* = u$, and let $T: M_{\text{cb}}A_p(G) \rightarrow M_{\text{cb}}A_p(H)$ be as in Lemma 5.1. In particular, the map T is weak*-weak*-continuous. Let m denote the (unique) invariant mean on $M_{\text{cb}}A_p(H)$. Then $m'(\varphi) = m(T\varphi)$ defines a linear functional on $M_{\text{cb}}A_p(G)$. In the same way as in the proof of [17, Proposition 5.9], we can show that m' is the (unique) invariant mean on $M_{\text{cb}}A_p(G)$. If m is weak*-continuous, then m' is weak*-continuous as well by the weak*-weak*-continuity of T . \square

Corollary 5.3. If G is a locally compact group with property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$ and N is a closed normal subgroup of G , then G/N has property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$.

Lemma 5.4. Let G_1 and G_2 be locally compact groups, let $1 < p < \infty$, and let $f_1 \in Q_{p,\text{cb}}(G_1)$ and $f_2 \in Q_{p,\text{cb}}(G_2)$. The function $f_1 \times f_2: G_1 \times G_2 \rightarrow \mathbb{C}$ given by

$$f_1 \times f_2(g_1, g_2) = f_1(g_1)f_2(g_2)$$

is an element of $Q_{p,\text{cb}}(G_1 \times G_2)$ and

$$(5) \quad \|f_1 \times f_2\|_{Q_{p,\text{cb}}(G_1 \times G_2)} \leq \|f_1\|_{Q_{p,\text{cb}}(G_1)} \|f_2\|_{Q_{p,\text{cb}}(G_2)}.$$

Proof. The proof is a simple modification of the proof of [12, Lemma 1.4] up to the following changes. Let $p' = \frac{p}{p-1}$. The tensor product space $T(G)$ in the proof of [12, Lemma 1.4] is replaced by

$$T_p(G) = \left(L^p(G) \widehat{\otimes} L^{p'}(G) \right) \widehat{\otimes} \left(L^{p'} \check{\otimes} L^p(G) \right)$$

(as is the case in Herz [21]). Use the fact that when $L: L^p(G_1) \rightarrow L^p(G_1)$ and $M: L^p(G_2) \rightarrow L^p(G_2)$ are bounded operators,

$$L \otimes M: L^p(G_1 \times G_2) \rightarrow L^p(G_1 \times G_2)$$

is also a bounded operator with $\|L \otimes M\| = \|L\| \|M\|$. \square

The following lemma generalises [17, Lemma 5.11] with the same proof.

Lemma 5.5. Let G be a locally compact group with property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$, and let $m \in Q_{p,\text{cb}}(G)$ denote the (unique) invariant mean on $M_{\text{cb}}A_p(G)$. Then there exists a sequence $f_n \in L^1(G)$ of non-negative functions with $\|f_n\|_1 = 1$ for all $n \in \mathbb{N}$ such that $\|f_n - m\|_{Q_{p,\text{cb}}(G)} \rightarrow 0$.

Proposition 5.6. Let G_1 and G_2 be two locally compact groups. The direct product $G = G_1 \times G_2$ has property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$ if and only if G_1 and G_2 have property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$.

Proof. Suppose that G_1 and G_2 have property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$. For $i = 1, 2$, let m_i be the invariant mean on $M_{\text{cb}}A_p(G_i)$. For $i = 1, 2$, by Lemma 5.5, there are sequences $(f_n^{(i)})$ in $L^1(G_i)_{\geq 0}$ with $\|f_n^{(i)}\|_1 = 1$ for all n such that $\|f_n^{(i)} - m_i\|_{Q_{p,\text{cb}}(G_i)} \rightarrow 0$ as $n \rightarrow \infty$. For all $g_i \in G_i$, we have

$$\|L_{g_i} f_n^{(i)} - f_n^{(i)}\|_{Q_{p,\text{cb}}(G_i)} \rightarrow 0,$$

because m_i is left invariant. By Lemma 5.4, the sequence $f_n^{(1)} \times f_n^{(2)}$ is a Cauchy sequence in $Q_{p,\text{cb}}(G)$, and its limit M is a weak*-continuous mean on $M_{\text{cb}}A_p(G)$. It follows that

$$\|L_g(f_n^{(1)} \times f_n^{(2)}) - f_n^{(1)} \times f_n^{(2)}\|_{Q_{p,\text{cb}}(G)} \rightarrow 0 \quad \text{for all } g \in G.$$

Therefore, the mean M is left invariant, so G has property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$.

The other direction follows directly from Corollary 5.3. \square

The proof of the following result is a modification of [17, Proposition 5.13].

Proposition 5.7. Let G be a locally compact group, and let K be a compact closed normal subgroup of G . Then G has property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$ if and only if G/K has property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$.

To conclude this section, we establish a relation between property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$ and property $(\mathbb{T}_{M_{\text{cb}}A_q}^*)$ for different p and q .

Proposition 5.8. Let G be a locally compact group and $1 < p \leq q \leq 2$ or $2 \leq q \leq p < \infty$. If G has property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$, then it has property $(\mathbb{T}_{M_{\text{cb}}A_q}^*)$.

Proof. Let $1 < p \leq q \leq 2$ or $2 \leq q \leq p < \infty$. Suppose that G has property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$. By [3, Proposition 6.1], the inclusion map

$$\iota: M_{\text{cb}}A_q(G) \rightarrow M_{\text{cb}}A_p(G)$$

is a contraction. Its adjoint $\iota^*: M_{\text{cb}}A_p(G)^* \rightarrow M_{\text{cb}}A_q(G)^*$, also a contraction, maps $Q_{p,\text{cb}}(G)$ to $Q_{q,\text{cb}}(G)$ (see [33, Proposition 2.1]). Therefore, $\iota: M_{\text{cb}}A_q(G) \rightarrow M_{\text{cb}}A_p(G)$ is weak*-weak*-continuous. Let m denote the (unique) invariant mean on $M_{\text{cb}}A_p(G)$. Since $m|_{M_{\text{cb}}A_q(G)} = m \circ \iota$ is the invariant mean on $M_{\text{cb}}A_q(G)$ and m and ι are weak*-continuous, we obtain that G has property $(\mathbb{T}_{M_{\text{cb}}A_q}^*)$. \square

6. SIMPLE LIE GROUPS WITH PROPERTY $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$

In this section, we determine exactly which connected simple Lie groups with finite center have property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$ for $1 < p < \infty$.

Recall that a (connected) Lie group is called simple if its Lie algebra is simple and that it is called semisimple if its Lie algebra is a direct sum of simple Lie algebras. Let G be a connected semisimple Lie group with finite center, and let \mathfrak{g} denote its Lie algebra. Then \mathfrak{g} has a Cartan decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$, where \mathfrak{k} is the Lie algebra of a maximal compact subgroup K of G . Furthermore, the Lie group G has a decomposition $G = KAK$, where A is an abelian Lie group, whose Lie algebra \mathfrak{a} is a maximal abelian subspace of \mathfrak{p} . The real rank of G is defined as the dimension of \mathfrak{a} . For details, see e.g. [31]. The real rank is an important invariant for our purposes.

Let us first recall two examples of simple Lie groups and show that they satisfy property $(\mathbb{T}_{M_{\text{cb}}A_p}^*)$.

First, let $\text{SL}(3, \mathbb{R})$ denote the special linear group, i.e. the group of 3×3 -matrices with real entries and determinant 1. The special orthogonal group $\text{SO}(3)$ is the natural maximal compact subgroup of $\text{SL}(3, \mathbb{R})$.

For the second example, let J be the matrix defined by

$$J = \begin{pmatrix} 0 & I_2 \\ -I_2 & 0 \end{pmatrix},$$

where I_2 is the identity 2×2 -matrix. The symplectic group $\text{Sp}(2, \mathbb{R})$ is defined as

$$\text{Sp}(2, \mathbb{R}) = \{g \in \text{GL}(4, \mathbb{R}) \mid g^t J g = J\},$$

where g^t denotes the transpose of g . The group

$$K = \left\{ \begin{pmatrix} A & -B \\ B & A \end{pmatrix} \in \text{Mat}_4(\mathbb{R}) \mid A + iB \in \text{U}(2) \right\} \cong \text{U}(2)$$

is a maximal compact subgroup of $\text{Sp}(2, \mathbb{R})$.

Proposition 6.1. For $1 < p < \infty$, the groups $\text{SL}(3, \mathbb{R})$ and $\text{Sp}(2, \mathbb{R})$ satisfy property $(\text{T}_{M_{\text{cb}}A_p}^*)$.

The proposition essentially follows from [33, Theorem 1.3 and Theorem 1.4]. Indeed, in these theorems, Vergara proves that the groups $\text{SL}(3, \mathbb{R})$ and $\text{Sp}(2, \mathbb{R})$ do not satisfy the p -AP for $1 < p < \infty$. The p -AP was introduced in [3] as a p -analogue of the Approximation Property of Haagerup and Kraus (AP), which goes back to [18]. It was proved by Lafforgue and de la Salle that $\text{SL}(3, \mathbb{R})$ does not have the AP [23]. This was generalised to all connected simple Lie groups with real rank at least 2, including the group $\text{Sp}(2, \mathbb{R})$, by Haagerup and the first-named author in [19], [20].

We now present a brief proof sketch of Proposition 6.1, relying on Vergara's work.

Proof sketch of Proposition 6.1. In fact, what is proved in [33, Theorem 1.3 and Theorem 1.4] is stronger than the fact that the groups $\text{SL}(3, \mathbb{R})$ and $\text{Sp}(2, \mathbb{R})$ do not have the p -AP for $1 < p < \infty$. We explain this for the group $\text{SL}(3, \mathbb{R})$. The case $\text{Sp}(2, \mathbb{R})$ follows analogously.

Let $G = \text{SL}(3, \mathbb{R})$, and let $K = \text{SO}(3)$ (its maximal compact subgroup). For $g \in G$, let m_g be the measure defined by

$$\int_G f dm_g = \int_K \int_K f(k_1 g k_2) dk_1 dk_2$$

Vergara shows that (m_g) is a Cauchy net in $M_{\text{cb}}A_p(G)^*$, which converges to a mean m on $M_{\text{cb}}A_p(G)$. Additionally, he proves that m is actually an element of $Q_{p, \text{cb}}(G)$, i.e. the mean m is weak*-continuous. What remains to be shown is the fact that m is actually an invariant mean, which is straightforward. \square

Remark 6.2. Note that the fact that $\text{SL}(3, \mathbb{R})$ and $\text{Sp}(2, \mathbb{R})$ have property (T^*) (which is exactly property $(\text{T}_{M_{\text{cb}}A_p}^*)$ for $p = 2$) was proved in [17]. Alternative to the proof sketch of Proposition 6.1 given above, one can modify the proofs from [17] to the p -setting. This would require some straightforward technical modifications.

Before we can characterise the connected simple Lie groups with property $(\text{T}_{M_{\text{cb}}A_p}^*)$, we recall the following powerful result of Veech [32, Theorem 1.4], which can be seen as a strong version of the Howe-Moore property. First, we recall some additional structure theory for Lie groups. Recall that a Lie group G is semisimple if its Lie algebra \mathfrak{g} decomposes as a direct sum $\mathfrak{g} = \mathfrak{s}_1 \oplus \dots \oplus \mathfrak{s}_n$, where the algebras \mathfrak{s}_i are simple Lie algebras. We say that a connected semisimple Lie group does not have compact simple factors if

for all $i = 1, \dots, n$, the analytic subgroup S_i of G corresponding to the Lie algebra \mathfrak{s}_i is not compact.

Theorem 6.3 (Veech). Let G be a connected semisimple Lie group with finite center and without compact simple factors. Then

$$\text{WAP}(G) = C_0(G) \oplus \mathbb{C}1$$

and for every $\varphi \in \text{WAP}(G)$, we have

$$m(\varphi) = \lim_{g \rightarrow \infty} \varphi(g),$$

where m is the unique invariant mean on $\text{WAP}(G)$.

We are now ready to prove the main result of this section.

Theorem 6.4. Let G be a connected simple Lie group with finite center, and let $1 < p < \infty$. Then G has property $(T_{M_{cb}A_p}^*)$ if and only if the real rank of G is 0 or at least 2.

Proof. Let G be a connected simple Lie group with finite center. Then G has real rank 0 if and only if it is compact, in which case it has property $(T_{M_{cb}A_p}^*)$ (see Section 4). If G has real rank 1, then it is well known that G is weakly amenable [12], and hence it has the AP of Haagerup and Kraus, so it cannot have property (T^*) , since by [17, Proposition 5.5] a locally compact group having the AP and property (T^*) has to be compact. By Proposition 5.8, the group G cannot have property $(T_{M_{cb}A_p}^*)$ for any $p \in (1, \infty)$.

Hence, it only remains to consider the case of real rank at least 2, which follows in the same way as [17, Theorem D]. Let us give a brief argument. Let G be a connected simple Lie group with finite center and real rank at least 2. It is well known that G contains a closed subgroup H that is locally isomorphic to $\text{SL}(3, \mathbb{R})$ or $\text{Sp}(2, \mathbb{R})$ (i.e. the Lie algebra of H is isomorphic to the Lie algebra of $\text{SL}(3, \mathbb{R})$ or to the Lie algebra of $\text{Sp}(2, \mathbb{R})$). Let m_H denote the invariant mean on $\text{WAP}(H)$, the restriction of which to $M_{cb}A_p(H)$ is weak*-continuous by Proposition 6.1.

We define a map from $\text{WAP}(G)$ to $\text{WAP}(H)$ in the following way, similar to Lemma 5.1. Let $u \in C_c(G)$ be a non-negative function such that $\|u\|_1 = 1$ and $u^* = u$. Define the map $T: \text{WAP}(G) \rightarrow \text{WAP}(H)$ by $T\varphi = (u * \varphi)|_H$. Since $T(C_0(G)) \subset C_0(H)$, we have $(m_H \circ T)|_{C_0(G)} \equiv 0$. Using Theorem 6.3, it follows that the mean m_G on $\text{WAP}(G)$ is given by $m_G = m_H \circ T$. By Lemma 5.1, we see that the restriction of the map T to $M_{cb}A_p(G)$ is weak*-weak*-continuous, and the result follows. \square

7. PROPERTY $(T_{M_{cb}A_p}^*)$ FOR CONNECTED LIE GROUPS

In this section, we study property $(T_{M_{cb}A_p}^*)$ for connected Lie groups. Let us first recall some structure theory for connected Lie groups, additional to the structure theory for semisimple Lie groups which we recalled before.

Let G be a connected Lie group with Lie algebra \mathfrak{g} . If \mathfrak{r} is the solvable radical (i.e. the largest solvable ideal) of \mathfrak{g} , then the Lie algebra \mathfrak{g} can be

written as $\mathfrak{g} = \mathfrak{r} \ltimes \mathfrak{s}$, where \mathfrak{s} is a semisimple Lie subalgebra. This decomposition is called the Levi decomposition. Furthermore, as was recalled at the beginning of Section 6, we can decompose \mathfrak{s} as $\mathfrak{s} = \mathfrak{s}_1 \oplus \cdots \oplus \mathfrak{s}_n$, where the \mathfrak{s}_i 's, with $i = 1, \dots, n$, are simple Lie algebras. Let R , S and S_i , with $i = 1, \dots, n$, be the analytic subgroups of G corresponding to the Lie algebras \mathfrak{r} , \mathfrak{s} and \mathfrak{s}_i , respectively. Thus, R is a solvable closed normal subgroup of G , and S is a (maximal) semisimple analytic subgroup of G . However, S is not necessarily a closed subgroup, and in general, we have $G = RS$, but $R \cap S$ can consist of more than one element. However, for simply connected G , the subgroup S is, in fact, closed, and the Levi decomposition becomes a semidirect product $G = R \ltimes S$. For details, see [31, Section 3.18].

Before we get to the main theorem, we prove some auxiliary results.

Lemma 7.1. Let G be a locally compact group with a compact normal subgroup K . Then $M_{\text{cb}A_p}(G) \cap C_0(G)$ is weak*-closed in $M_{\text{cb}A_p}(G)$ if and only if $M_{\text{cb}A_p}(G/K) \cap C_0(G/K)$ is weak*-closed in $M_{\text{cb}A_p}(G/K)$.

Proof. Suppose that $M_{\text{cb}A_p}(G) \cap C_0(G)$ is weak*-closed in $M_{\text{cb}A_p}(G)$. Let (φ_i) be a net in $M_{\text{cb}A_p}(G/K) \cap C_0(G/K)$ that converges to $\varphi \in M_{\text{cb}A_p}(G/K)$ in the weak*-topology on $M_{\text{cb}A_p}(G/K)$. Let $q: G \rightarrow G/K$ be the canonical quotient map. From the proof of [33, Proposition 5.1], it follows that the operator

$$T: M_{\text{cb}A_p}(G/K) \rightarrow M_{\text{cb}A_p}(G), \quad \psi \mapsto \psi \circ q$$

is weak*-weak*-continuous. Therefore, $(T(\varphi_i))$ is a net in $M_{\text{cb}A_p}(G) \cap C_0(G)$ that converges in the weak*-topology to $T(\varphi) \in M_{\text{cb}A_p}(G)$. Since $M_{\text{cb}A_p}(G) \cap C_0(G)$ is weak*-closed in $M_{\text{cb}A_p}(G)$, we have $T(\varphi) \in C_0(G)$. Hence, φ belongs to $C_0(G/K)$.

For the converse, let (φ_i) be a net in $M_{\text{cb}A_p}(G) \cap C_0(G)$ that converges to φ in $M_{\text{cb}A_p}(G)$ in the weak*-topology of $M_{\text{cb}A_p}(G)$. Again, it follows from the proof of [33, Proposition 5.1] that the operator

$$\tilde{T}: M_{\text{cb}A_p}(G) \rightarrow M_{\text{cb}A_p}(G/K),$$

defined by $\tilde{T}(\psi)(gK) = \int_K \psi(gk)dk$ for $\psi \in M_{\text{cb}A_p}(G)$ and $g \in G$, is weak*-weak*-continuous. Hence, the net $(\tilde{T}(\varphi_i))$ in $M_{\text{cb}A_p}(G/K) \cap C_0(G/K)$ converges in the weak*-topology to $\tilde{T}(\varphi)$ in $M_{\text{cb}A_p}(G/K)$. By assumption, $\tilde{T}(\varphi)$ belongs to $C_0(G/K)$. Hence, $\varphi \in C_0(G)$. \square

The following is a generalisation of [7, Lemma 2.5].

Lemma 7.2. Let G be a connected Lie group with connected closed normal subgroup N . Suppose that $M_{\text{cb}A_p}(G) \cap C_0(G)$ is weak*-closed in $M_{\text{cb}A_p}(G)$. Then the center $Z(N)$ of N is compact and $N/Z(N)$ is semisimple.

Proof. Since the identity map $L^1(G) \rightarrow L^1(G)$ extends to a contraction $Q_{p,\text{cb}}(G) \rightarrow C^*(G)$, its Banach adjoint is a weak*-weak*-continuous contraction $B(G) \rightarrow M_{\text{cb}A_p}(G)$ that is easily verified to be the inclusion map.

Hence, $B(G) \cap C_0(G)$ is weak*-closed in $B(G)$. The claim now directly follows from [7, Lemma 2.5]. \square

We can now state and prove the main theorem. For the purpose of this article, a connected (real) Lie group G is said to be reductive if the solvable radical R of G is abelian. (The definition of reductive Lie group is subject to conventions, which we will not further discuss here.) The proof of this theorem is inspired by [7, Section 2].

Theorem 7.3. Let G be a connected Lie group, and let $1 < p < \infty$. Suppose that the semisimple part S of the Levi decomposition of G has finite center. Then the following are equivalent:

- (i) The group G is a reductive Lie group with compact center satisfying property $(T_{M_{cb}A_p}^*)$.
- (ii) The space $M_{cb}A_p(G) \cap C_0(G)$ is closed in $M_{cb}A_p(G)$ in the $\sigma(M_{cb}A_p(G), Q_{p,cb}(G))$ -topology.
- (iii) The group G is a reductive Lie group with compact centre, in which every simple factor has real rank 0 or at least 2.

Proof of Theorem 7.3. (i) \implies (ii): Suppose that G is a reductive Lie group with compact center satisfying property $(T_{M_{cb}A_p}^*)$. Using that the center $Z(G)$ is a compact normal subgroup of G , it is enough to show that $M_{cb}A_p(G/Z(G)) \cap C_0(G/Z(G))$ is weak*-closed in $M_{cb}A_p(G/Z(G))$ by Lemma 7.1. To this end, let (φ_i) be a net in $M_{cb}A_p(G/Z(G)) \cap C_0(G/Z(G))$ that converges to $\varphi \in M_{cb}A_p(G/Z(G))$ in the weak*-topology. By assumption, the group $G/Z(G)$ is semisimple. From Theorem 6.3, it follows that

$$m(\varphi) = \lim_{g \rightarrow \infty} \varphi(g) \quad \text{and} \quad m(\varphi_i) = \lim_{g \rightarrow \infty} \varphi_i(g) \quad \text{for all } i,$$

where m denotes the unique invariant mean on $M_{cb}A_p(G/Z(G))$. Since φ_i belongs to $C_0(G/Z(G))$, we have $m(\varphi_i) = 0$ for all i . Since property $(T_{M_{cb}A_p}^*)$ passes to quotients by Proposition 5.7, the quotient group $G/Z(G)$ also has property $(T_{M_{cb}A_p}^*)$, and hence, we also have $m(\varphi) = \lim_i m(\varphi_i) = 0$, so φ belongs to $C_0(G/Z(G))$.

(ii) \implies (i): Suppose that $M_{cb}A_p(G) \cap C_0(G)$ is weak*-closed in $M_{cb}A_p(G)$. By Lemma 7.2, we know that $Z(G)$ is compact (abelian) and $G/Z(G)$ is semisimple. Hence, G is a reductive Lie group with compact center, and by Lemma 7.1, the space $M_{cb}A_p(G/Z(G)) \cap C_0(G/Z(G))$ is weak*-closed in $M_{cb}A_p(G/Z(G))$.

By the fact that $G/Z(G)$ is semisimple, the kernel of the unique invariant mean on $M_{cb}A_p(G/Z(G))$ coincides with $M_{cb}A_p(G/Z(G)) \cap C_0(G/Z(G))$ by Theorem 6.3. Since this kernel is weak*-closed, the mean on $M_{cb}A_p(G/Z(G))$ is weak*-continuous, so the group $G/Z(G)$ has property $(T_{M_{cb}A_p}^*)$. Hence, by Proposition 5.7, the group G has property $(T_{M_{cb}A_p}^*)$ as well.

(i) \iff (iii): Let $\tilde{G} = \mathbb{R}^n \times \tilde{S}_1 \times \dots \times \tilde{S}_n$ be the universal covering group of G , where $\tilde{S}_1, \dots, \tilde{S}_n$ are simply connected simple Lie groups. There exists

a discrete subgroup Γ of the center $Z(\tilde{G}) = \mathbb{R}^n \times Z(\tilde{S}_1) \times \dots \times Z(\tilde{S}_n)$ of \tilde{G} such that $G = \tilde{G}/\Gamma$. Hence, we have

$$G/Z(G) = \tilde{G}/Z(\tilde{G}) = \left(\tilde{S}_1/Z(\tilde{S}_1) \right) \times \dots \times \left(\tilde{S}_n/Z(\tilde{S}_n) \right).$$

Since $Z(G)$ is compact, the group G has property $(T_{M_{cb}A_p}^*)$ if and only if $G/Z(G)$ has property $(T_{M_{cb}A_p}^*)$, by Proposition 5.7. It follows from Proposition 5.6 that $G/Z(G)$ has property $(T_{M_{cb}A_p}^*)$ if and only if each $\tilde{S}_i/Z(\tilde{S}_i)$ has property $(T_{M_{cb}A_p}^*)$. From Theorem 6.4, we know that $\tilde{S}_i/Z(\tilde{S}_i)$ has property $(T_{M_{cb}A_p}^*)$ if and only if $\tilde{S}_i/Z(\tilde{S}_i)$, which is a simple Lie group with trivial center, has real rank 0 or real rank at least 2. \square

Remark 7.4. It is an open question whether there exists a group with property (T^*) , but without property $(T_{M_{cb}A_p}^*)$ for some $p \neq 2$.

8. PROPERTY $(T_{B_p}^*)$ FOR CONNECTED LIE GROUPS

In this section, we study property $(T_{B_p}^*)$ for connected Lie groups, and we prove Theorem 8.2, which is analogous to (but less explicit than) Theorem 7.3. Indeed, we prove the equivalence of the analogues of the first two equivalent assertions of Theorem 7.3.

First, we establish the following lemma.

Lemma 8.1. Let G be a locally compact group with a compact normal subgroup K . Then $B_p(G) \cap C_0(G)$ is weak*-closed in $B_p(G)$ if and only if $B_p(G/K) \cap C_0(G/K)$ is weak*-closed in $B_p(G/K)$.

Proof. By the same arguments as given in the proof of [33, Proposition 5.1], the operator $T: B_p(G/K) \rightarrow B_p(G)$, $\psi \mapsto \psi \circ q$ and the operator $\tilde{T}: B_p(G) \rightarrow B_p(G/K)$ defined by $\tilde{T}(\psi)(gK) = \int_K \psi(gk)dk$ are weak*-weak*-continuous. \square

We have now established the right structural properties for $B_p(G)$ and permanence properties for property $(T_{B_p}^*)$. The proof of the following theorem is similar to the corresponding parts of the proof of Theorem 7.3. We only explain the relevant points of the proof.

Theorem 8.2. Let G be a connected Lie group, and let $1 < p < \infty$. Suppose that the semisimple part S of the Levi decomposition of G has finite center. Then the following are equivalent:

- (i) The group G is a reductive Lie group with compact center satisfying property $(T_{B_p}^*)$.
- (ii) The space $B_p(G) \cap C_0(G)$ is closed in $B_p(G)$ in the $\sigma(B_p(G), B_p(G)_*)$ -topology.

The proof of this theorem follows mutatis mutandis from the proof of Theorem 7.3.

Remark 8.3. Theorem 8.2 generalises [7, Theorem 2.7]. An important difference with the situation of property (T), as covered in [7], however, is that it is known exactly which simple Lie groups have property (T). Indeed, let G be a connected simple Lie group. Then G has property (T) if and only if G has real rank 0, real rank at least 2 or if G is locally isomorphic to $\mathrm{Sp}(n, 1)$ (with $n \geq 2$) or to $F_{4(-20)}$ (in the latter two cases, G has real rank 1).

It is clear that if G has real rank 0 or at least 2, then G has property $(\mathrm{T}_{B_p}^*)$, which follows from the fact that in these cases, G has property $(\mathrm{T}_{M_{\mathrm{cb}A_p}}^*)$ (see Theorem 6.4), because property $(\mathrm{T}_{M_{\mathrm{cb}A_p}}^*)$ is stronger than property $(\mathrm{T}_{B_p}^*)$.

However, if G is locally isomorphic to $\mathrm{Sp}(n, 1)$ (with $n \geq 2$) or to $F_{4(-20)}$, we do not know what happens, although we expect that these groups have property $(\mathrm{T}_{B_p}^*)$ for $1 < p < \infty$. In the remaining cases of simple Lie groups with real rank 1, i.e. groups locally isomorphic to $\mathrm{SO}_0(n, 1)$ or $\mathrm{SU}(n, 1)$ (with $n \geq 2$), we know that these groups have the Haagerup property, in which case they cannot have property $(\mathrm{T}_{B_p}^*)$ for $1 < p < \infty$.

REFERENCES

- [1] C.A. Akemann and M.E. Walter, *Unbounded negative definite functions*, Canadian J. Math. **33** (1981), 862–871.
- [2] L. Alaoglu and G. Birkhoff, *General ergodic theorems*, Ann. of Math. (2) **41** (1940), 293–309.
- [3] G. An, J.-J. Lee and Z.-J. Ruan, *On p -approximation properties for p -operator spaces*, J. Funct. Anal. **259** (2010), 933–974.
- [4] U. Bader and T. Gelander, *Equicontinuous actions of semisimple groups*, Groups Geom. Dyn. **11** (2017), 1003–1039.
- [5] U. Bader, C. Rosenthal and R. Sauer, *On the cohomology of weakly almost periodic group representations*, J. Topol. Anal. **6** (2014), 153–165.
- [6] B. Bekka, P. de la Harpe and A. Valette, *Kazhdan’s property (T)*, Cambridge University Press, Cambridge, 2008.
- [7] B. Bekka, E. Kaniuth, A.T. Lau and G. Schlichting, *Weak*-closedness of subspaces of Fourier-Stieltjes algebras and weak*-continuity of the restriction map*, Trans. Amer. Math. Soc. **350** (1998), 2277–2296.
- [8] M. Bożejko, *Positive and negative definite kernels on discrete groups*, Lectures at Heidelberg University, 1987.
- [9] R.B. Burckel, *Weakly almost periodic functions on semigroups*, Gordon and Breach, New York, 1970.
- [10] J. De Cannière and U. Haagerup, *Multipliers of the Fourier algebras of some simple Lie groups and their discrete subgroups*, Amer. J. Math. **107** (1985), 455–500.
- [11] J.B. Conway, *A course in functional analysis*, Springer-Verlag, New York, 1990.
- [12] M. Cowling and U. Haagerup, *Completely bounded multipliers of the Fourier algebra of a simple Lie group of real rank one*, Invent. Math. **96** (1989), 507–549.
- [13] M. Daws, *p -operator spaces and Figà-Talamanca–Herz algebras*, J. Operator Theory **63** (2010), 47–83.
- [14] P. Eymard, *L’algèbre de Fourier d’un groupe localement compact*, Bull. Soc. Math. France **92** (1964), 181–236.
- [15] R. Godement, *Les fonctions de type positif et la théorie des groupes*, Trans. Amer. Math. Soc. **63** (1948), 1–84.

- [16] F. Greenleaf, *Invariant means on topological groups and their applications*, Van Nostrand Reinhold, New York, 1969.
- [17] U. Haagerup, S. Knudby and T. de Laat, *A complete characterization of connected Lie groups with the Approximation Property*, Ann. Sci. Éc. Norm. Supér. (4) **49** (2016), 927–946.
- [18] U. Haagerup and J. Kraus, *Approximation properties for group C^* -algebras and group von Neumann algebras* Trans. Amer. Math. Soc. **344** (1994), 667–699.
- [19] U. Haagerup and T. de Laat, *Simple Lie groups without the Approximation Property*, Duke Math. J. **162** (2013), 925–964.
- [20] U. Haagerup and T. de Laat, *Simple Lie groups without the Approximation Property II*, Trans. Amer. Math. Soc. **368** (2016), 3777–3809.
- [21] C. Herz, *Une généralisation de la notion de transformée de Fourier-Stieltjes*, Ann. Inst. Fourier (Grenoble) **24** (1974), 145–157.
- [22] S. Kaijser, *On Banach modules. I.*, Math. Proc. Cambridge Philos. Soc. **90** (1981), 423–444.
- [23] V. Lafforgue and M. de la Salle, *Noncommutative L^p -spaces without the completely bounded approximation property*, Duke Math. J. **160** (2011), 71–116.
- [24] M. Megrelishvili, *Fragmentability and representations of flows*, Proceedings of the 17th Summer Conference on Topology and its Applications, Topology Proc. **27** (2003), 497–544.
- [25] T. Miao, *Predual of the multiplier algebra of $A_p(G)$ and amenability*, Canad. J. Math. **56** (2004), 344–355.
- [26] T. Miao, *Approximation properties and approximate identity of $A_p(G)$* , Trans. Amer. Math. Soc. **361** (2009) 1581–1595.
- [27] T. Miao, Private communication.
- [28] V. Runde, *Representations of locally compact groups on QSL_p -spaces and a p -analog of the Fourier-Stieltjes algebra*, Pacific J. Math. **221** (2005), 379–397.
- [29] T. Shulman, *On subspaces of invariant vectors*, Studia Math. **236** (2017), 1–11.
- [30] A. Valette, *Minimal projections, integrable representations and property (T)*, Arch. Math. (Basel) **43** (1984), 397–406.
- [31] V.S. Varadarajan, *Lie groups, Lie algebras, and their representations*, Springer-Verlag, New York, 1984.
- [32] W.A. Veech, *Weakly almost periodic functions on semisimple Lie groups*, Monatsh. Math. **88** (1979), 55–68.
- [33] I. Vergara, *The p -approximation property for simple Lie groups with finite center*, J. Funct. Anal. **273** (2017), 3463–3503.
- [34] G. Xu, *Herz-Schur multipliers and weakly almost periodic functions on locally compact groups*, Trans. Amer. Math. Soc. **349** (1997), 2525–2536.

TIM DE LAAT,
WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, MATHEMATISCHES INSTITUT
EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY
Email address: tim.delaat@uni-muenster.de

SAFOURA ZADEH,
MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111, BONN, GERMANY
Email address: jsafoora@gmail.com