



Mathematisches
Forschungsinstitut
Oberwolfach



Oberwolfach Preprints

Number 2024-10

ANNA ABBATIELLO
DANICA BASARIĆ
NILASIS CHAUDHURI
EDUARD FEIREISL

Local Existence and Conditional Regularity
for the Navier-Stokes-Fourier System Driven
by Inhomogeneous Boundary Conditions

September 2024

Oberwolfach Preprints (OWP)

The MFO publishes the Oberwolfach Preprints (OWP) as a series which mainly contains research results related to a longer stay in Oberwolfach, as a documentation of the research work done at the MFO. In particular, this concerns the Oberwolfach Research Fellows program (and the former Research in Pairs program) and the Oberwolfach Leibniz Fellows (OWLF), but this can also include an Oberwolfach Lecture, for example.

All information about the publication process can be found at <https://www.mfo.de/scientific-programme/publications/owp>

All published Oberwolfach Preprints can be found at <https://publications.mfo.de>

ISSN 1864-7596

License Information

This document may be downloaded, read, stored and printed for your own use within the limits of § 53 UrhG but it may not be distributed via the internet or passed on to external parties.

The series Oberwolfach Preprints is published by the MFO. Copyright of the content is held by the authors.

Mathematisches Forschungsinstitut Oberwolfach gGmbH (MFO)
Schwarzwaldstrasse 9-11
77709 Oberwolfach-Walke
Germany
<https://www.mfo.de>

DOI 10.14760/OWP-2024-10



Local existence and conditional regularity for the Navier–Stokes–Fourier system driven by inhomogeneous boundary conditions

Anna Abbatiello* Danica Basarić † Nilasis Chaudhuri ‡ Eduard Feireisl §

University of Campania “L. Vanvitelli”, Department of Mathematics and Physics
Viale A. Lincoln 5, 81100 Caserta, Italy
anna.abbatiello@unicampania.it

Politecnico di Milano, Department of Mathematics
Via E. Bonardi 9, 20133 Milano, Italy
danica.basaric@polimi.it

University of Warsaw, Faculty of Mathematics, Informatics and Mechanics
Stefana Banacha 2, 02-097 Warsaw, Poland
nchaudhuri@mimuw.edu.pl

Institute of Mathematics, Czech Academy of Sciences
Žitná 25, 115 67 Praha 1, Czech Republic
feireisl@math.cas.cz

Abstract

We consider the Navier–Stokes–Fourier system with general inhomogeneous Dirichlet–Neumann boundary conditions. We propose a new approach to the local well-posedness problem based on conditional regularity estimates. By conditional regularity we mean that any strong solution belonging to a suitable class remains regular as long as its amplitude remains bounded. The result holds for general Dirichlet–Neumann boundary conditions provided the material derivative of the velocity field vanishes on the boundary of the physical domain. As a corollary of this result we obtain:

- Blow up criteria for strong solutions
- Local existence of strong solutions in the optimal $L^p - L^q$ framework
- Alternative proof of the existing results on local well posedness

*The work of A.A. is performed under the auspices of the Italian National Group of the Mathematical Physics (GNFM) of INdAM.

†The work of D.B. was supported by the PRIN project 2022 “Partial differential equations and related geometric-functional inequalities”, financially supported by the EU, in the framework of the “Next Generation EU initiative”. The Department of Mathematics of Politecnico di Milano is supported by MUR “Excellence Department 2023-2027”.

‡The work of N.C. was supported by the “Excellence Initiative Research University (IDUB)” program at the University of Warsaw.

§The work of E.F. was partially supported by the Czech Sciences Foundation (GAČR), Grant Agreement 24–11034S. The Institute of Mathematics of the Academy of Sciences of the Czech Republic is supported by RVO:67985840. E.F. is member of the Nečas Center for Mathematical Modelling.

2020 Mathematics Subject Classification: (primary)35Q35, 35Q30; (secondary) 76N10, 76N06 .

Keywords: Navier–Stokes–Fourier system, strong solution, Dirichlet problem, local existence, conditional regularity, blow–up criterion

Contents

1	Introduction	2
2	Problem formulation and main results	3
2.1	Navier–Stokes–Fourier system	3
2.2	Strong solutions in the $L^p - L^q$ framework	4
3	New formulation and minimum principle	6
3.1	Minimum principle for the density	6
3.2	Minimum principle for the temperature	7
4	Classes of strong solutions	7
4.1	Hilbert space – energy framework	8
4.2	Mixed setting	9
4.3	$L^p - L^q$ setting	11
5	Conditional regularity estimates	12
5.1	Conditional regularity in the mixed and Hilbertian setting	12
5.1.1	Energy estimates	13
5.1.2	Estimates on the material derivative	13
5.2	Conditional regularity estimates in the $L^p - L^q$ setting	18
6	Local existence of strong solutions	22
6.1	Local existence $L^p - L^q$ setting	22
6.1.1	Data approximation	22
6.1.2	Control functionals and hitting times	22
6.2	Local existence in the mixed setting	25
7	Conditional regularity in the $L^p - L^q$ framework	26
7.1	Estimates for more regular data	26
7.2	Initial smoothing	27

1 Introduction

We propose a new approach to problems of well posedness of systems of equations arising in fluid dynamics based on *a priori* bounds resulting from conditional regularity estimates. The method consists in several steps:

1. First, a local in time existence of smooth solutions is established in a class of very regular functions in the spirit of Nash’s original paper [22]. In fact, we use a more recent “energy framework” proposed by Matsumura and Nishida [20], later developed by Valli [26], [27], Valli and Zajackowski [28], or Kagei and Kawashima [17]. The energy method is based on a simple Hilbertian framework based on suitable scales of Sobolev spaces, where the desired level of regularity is achieved by successive

differentiation of the field equations in the time variable. As a result, high regularity of the data is necessary accompanied by several levels of compatibility conditions to be satisfied.

2. Probably optimal results can be achieved in the more recent framework of $L^p - L^q$ regularity spaces used in the seminal paper of Solonnikov [24], and later revisited and developed by Danchin [8], [9], and Kotschote [18]. The $L^p - L^q$ class seems optimal both with respect to the required regularity of the data and the number of compatibility conditions.
3. We propose a new approach to the local existence in the $L^p - L^q$ setting based on suitable *a priori* bounds in terms of the data and a suitable *control* functional evaluated in terms of suitable norms of the local solution. The essential ingredient of the method is weak continuity (compactness) of the control functional with respect to the norm of the function space in which the desired solution lives. A typical example of such a functional is the amplitude – L^∞ -norm of the solution – provided the underlying solution space is compactly embedded in the space of continuous function. With this type of estimates at hand, the local solution can be obtained in the desired class on a time interval determined by “hitting” time of the control functional.
4. Besides the new local existence results, our approach yield a blow-up criterion in terms of the amplitude of the local solution in the spirit of the celebrated Nash’s conjecture, cf. Nash [21], and [15].

The paper is organized as follows. In Section 2, we introduce the Navier–Stokes–Fourier system along with the relevant initial/boundary conditions and state our main results. Various concepts of strong solution are introduced in Section 4. Section 5 presents preliminary conditional regularity estimates. In Section 6 we establish the local existence result in the $L^p - L^q$ framework. The proof of conditional regularity and blow-up criterion are completed in Section 7.

2 Problem formulation and main results

We introduce the initial/boundary value problem for the Navier–Stokes–Fourier system and state our main results.

2.1 Navier–Stokes–Fourier system

The time evolution of the mass density $\varrho = \varrho(t, x)$, the (absolute) temperature $\vartheta = \vartheta(t, x)$, and the velocity $\mathbf{u} = \mathbf{u}(t, x)$ of a viscous, compressible, and heat conducting fluid is governed by the *Navier–Stokes–Fourier* (NSF) *system* of equations:

$$\partial_t \varrho + \operatorname{div}_x(\varrho \mathbf{u}) = 0, \tag{2.1}$$

$$\partial_t(\varrho \mathbf{u}) + \operatorname{div}_x(\varrho \mathbf{u} \otimes \mathbf{u}) + \nabla_x p(\varrho, \vartheta) = \operatorname{div}_x \mathbb{S}(\mathbb{D}_x \mathbf{u}) + \varrho \nabla_x G, \tag{2.2}$$

$$\partial_t(\varrho e(\varrho, \vartheta)) + \operatorname{div}_x(\varrho e(\varrho, \vartheta) \mathbf{u}) + \operatorname{div}_x \mathbf{q}(\nabla_x \vartheta) = \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \mathbb{D}_x \mathbf{u} - p(\varrho, \vartheta) \operatorname{div}_x \mathbf{u} \tag{2.3}$$

satisfied for $(t, x) \in (0, T) \times \Omega$, $\Omega \subset \mathbb{R}^3$, where $\mathbb{D}_x \mathbf{u}$ denotes the symmetric velocity gradient:

$$\mathbb{D}_x \mathbf{u} \equiv \frac{1}{2} (\nabla_x \mathbf{u} + \nabla_x^t \mathbf{u}).$$

The pressure p and the internal energy e are related to the density and the temperature through the conventional *Boyle–Mariotte* equation of state (EOS):

$$p = p(\varrho, \vartheta) = \varrho\vartheta, \quad e = e(\vartheta) = c_v\vartheta, \quad c_v > 0. \quad (2.4)$$

The viscous stress \mathbb{S} is given by *Newton’s rheological law*, while the heat flux \mathbf{q} obeys *Fourier’s law*:

$$\mathbb{S} = \mathbb{S}(\mathbb{D}_x \mathbf{u}) = \mu \left(\nabla_x \mathbf{u} + \nabla_x^t \mathbf{u} - \frac{2}{3} \operatorname{div}_x \mathbf{u} \mathbb{I} \right) + \lambda \operatorname{div}_x \mathbf{u}, \quad (2.5)$$

$$\mathbf{q} = \mathbf{q}(\nabla_x \vartheta) = -\kappa \nabla_x \vartheta, \quad (2.6)$$

with constant transport coefficients $\mu > 0$, $\lambda \geq 0$, $\kappa > 0$.

The original state of the system is determined by the initial conditions

$$\varrho(0, \cdot) = \varrho_0, \quad \vartheta(0, \cdot) = \vartheta_0, \quad \mathbf{u}(0, \cdot) = \mathbf{u}_0. \quad (2.7)$$

In addition, we suppose the fluid occupies a bounded sufficiently smooth domain $\Omega \subset \mathbb{R}^3$ and impose the Dirichlet boundary conditions for the velocity

$$\mathbf{u}|_{\partial\Omega} = \mathbf{u}_B, \quad \mathbf{u}_B \cdot \mathbf{n} = 0 \quad (2.8)$$

where \mathbf{n} denotes the outer normal vector to $\partial\Omega$. Finally, we assume the boundary $\partial\Omega$ can be decomposed as

$$\partial\Omega = \Gamma_D \cup \Gamma_N, \quad \Gamma_D \cap \Gamma_N = \emptyset, \quad \Gamma_D, \Gamma_N \text{ compact}, \quad (2.9)$$

where

$$\vartheta|_{\Gamma_D} = \vartheta_B, \quad \nabla_x \vartheta \cdot \mathbf{n}|_{\Gamma_N} = q_B. \quad (2.10)$$

The boundary data $\mathbf{u}_B = \mathbf{u}_B(x)$, $\vartheta_B = \vartheta_B(x)$, and $q_B = q_B(x)$ are independent of the time. The hypothesis \mathbf{u}_B is tangential to the boundary plays a crucial role in the analysis and could possibly be relaxed by imposing a more restrictive regularity criterion.

2.2 Strong solutions in the $L^p - L^q$ framework

Following the seminal paper by Solonnikov [24], we look for solutions in the $L^p - L^q$ class. Specifically,

$$\varrho \in C([0, T]; W^{1,q}(\Omega)), \quad \partial_t \varrho \in L^p(0, T; L^q(\Omega)), \quad (2.11)$$

$$\vartheta \in L^p(0, T; W^{2,q}(\Omega)), \quad \partial_t \vartheta \in L^p(0, T; L^q(\Omega)), \quad (2.12)$$

$$\mathbf{u} \in L^p(0, T; W^{2,q}(\Omega; \mathbb{R}^3)), \quad \partial_t \mathbf{u} \in L^p(0, T; L^q(\Omega; \mathbb{R}^3)). \quad (2.13)$$

Accordingly,

$$\vartheta \in C([0, T]; B_{q,p}^{2(1-\frac{1}{p})}(\Omega)), \quad \mathbf{u} \in C([0, T]; B_{q,p}^{2(1-\frac{1}{p})}(\Omega; \mathbb{R}^3)),$$

where the Besov space $B_{q,p}^{2(1-\frac{1}{p})}(\Omega)$ can be identified with the real interpolation space

$$B_{q,p}^{2(1-\frac{1}{p})}(\Omega) = [L^q(\Omega); W^{2,q}(\Omega)]_{1-\frac{1}{p}, p},$$

see Amann [2, Section 5], [3, Chapter III, Theorem 4.10.2].

Our main result reads as follows.

Theorem 2.1 (NSF system - local solutions, blow up criterion). *Let $\Omega \subset \mathbb{R}^3$ be a bounded domain, with the boundary of class C^4 admitting the decomposition (2.9). Let the exponents p, q ,*

$$3 < q < \infty, \quad \frac{2q}{2q-3} < p < \infty, \quad (2.14)$$

be given.

Let $G = G(x)$, $G \in W^{1,q}(\Omega)$ be a given potential. Let the initial data $(\varrho_0, \vartheta_0, \mathbf{u}_0)$ and the boundary data $(\vartheta_B, \mathbf{u}_B, q_B)$ belong to the class

$$\varrho_0 \in W^{1,q}(\Omega), \quad \inf_{\Omega} \varrho_0 \geq \underline{\varrho} > 0, \quad (2.15)$$

$$\vartheta_0 \in B_{q,p}^{2(1-\frac{1}{p})}(\Omega), \quad \inf_{\Omega} \vartheta_0 \geq \underline{\vartheta} > 0, \quad (2.16)$$

$$\mathbf{u}_0 \in B_{q,p}^{2(1-\frac{1}{p})}(\Omega; \mathbb{R}^3), \quad (2.17)$$

$$\vartheta_B \in W^{2-\frac{1}{q},q}(\Gamma_D), \quad \inf_{\Gamma_D} \vartheta_B \geq \underline{\vartheta} > 0, \quad (2.18)$$

$$\mathbf{u}_B \in W^{2-\frac{1}{q},q}(\partial\Omega; \mathbb{R}^3), \quad \mathbf{u}_B \cdot \mathbf{n} = 0, \quad (2.19)$$

$$q_B \in W^{1-\frac{1}{q},q}(\Gamma_N), \quad q_B \geq 0. \quad (2.20)$$

In addition let the compatibility conditions

$$\mathbf{u}_0|_{\partial\Omega} = \mathbf{u}_B, \quad (2.21)$$

$$\vartheta_0|_{\Gamma_D} = \vartheta_B, \quad \nabla_x \vartheta_0 \cdot \mathbf{n}|_{\Gamma_N} = q_B \quad (2.22)$$

hold. Finally, suppose either $\Gamma_D \neq \emptyset$ or $q_B = 0$.

Then the following holds.

- **Local existence.** *There exists $T > 0$ such that the NSF system (2.1)–(2.6), with the initial conditions (2.7), and the boundary conditions (2.8), (2.10) admits a unique solution $(\varrho, \vartheta, \mathbf{u})$ in the class (2.11)–(2.13). In addition,*

$$\min_{[0,T] \times \bar{\Omega}} \varrho > 0, \quad \min_{[0,T] \times \bar{\Omega}} \vartheta > 0. \quad (2.23)$$

- **Blow-up criterion.** *Suppose $G \in W^{2,q}(\Omega)$. Let*

$$T_{\max} = \sup \left\{ T > 0 \mid \text{the solution } (\varrho, \vartheta, \mathbf{u}) \text{ exists in } [0, T] \right\}.$$

Then either $T_{\max} = \infty$ or

$$\limsup_{t \rightarrow (T_{\max})^-} \|(\varrho, \vartheta, \mathbf{u})(t, \cdot)\|_{C(\bar{\Omega}; \mathbb{R}^5)} = \infty. \quad (2.24)$$

Remark 2.2. The required regularity of $\partial\Omega$ is enforced by the method of construction and is not optimal. The result should hold for Ω of class C^2 or even $C^{1,1}$.

Remark 2.3. Hypothesis (2.14) allows to control the quadratic term $\mathbb{S}(\mathbb{D}_x \mathbf{u}) : \mathbb{D}_x \mathbf{u}$ in terms of the available parabolic estimates, cf. the proof of Proposition 5.4. It is also necessary for the (compact) embedding

$$B_{q,p}^{2(1-\frac{1}{p})}(\Omega) \hookrightarrow \hookrightarrow C(\bar{\Omega}),$$

cf. Amann [4]. In particular, T_{\max} is independent of p, q as long as (2.14) holds.

Remark 2.4. While the positivity of the flux q_B is used in the proof of minimum principle, see Section 3 below, the alternative $\Gamma_D \neq \emptyset$ or $q_B = 0$ is possibly only technical. In fact, it is used in Section 5 to extend the Neumann boundary condition on Γ_N inside Ω .

Theorem 2.1 extends the result of Kotschote [18] to the $L^p - L^q$ framework, $p \neq q$ in general. The blow-up criterion is new in the $L^p - L^q$ setting and the proof is based on a combination of three main ingredients:

- Extension of the conditional regularity estimates obtained for more regular solutions in [5] to a larger class introduced by Cho and Kim [7].
- New *a priori* estimates in the $L^p - L^q$ framework based on conditional regularity estimates.
- Pasting the local solutions (and the corresponding estimates) in Cho–Kim class, with the solutions in the $L^p - L^q$ class.

The rest of the paper is devoted to the proof of Theorem 2.1 based on the above scenario. As a by product, we obtain local existence of solutions in the Cho and Kim class for inhomogeneous boundary data (Theorem 6.2), and new conditional regularity estimates in terms of the amplitude of strong solutions in the $L^p - L^q$ class (Theorem 7.1).

3 New formulation and minimum principle

Using the specific EOS (2.4), we can rewrite (2.1)–(2.3) in the form

$$\partial_t \varrho + \mathbf{u} \cdot \nabla_x \varrho = -\varrho \operatorname{div}_x \mathbf{u}, \quad (3.1)$$

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla_x \mathbf{u} = \frac{1}{\varrho} \operatorname{div}_x \mathbb{S}(\mathbb{D}_x \mathbf{u}) - \vartheta \nabla_x \log(\varrho) - \nabla_x \vartheta + \nabla_x G, \quad (3.2)$$

$$\partial_t \vartheta + \mathbf{u} \cdot \nabla_x \vartheta = \frac{\kappa}{c_v \varrho} \Delta_x \vartheta + \frac{1}{c_v \varrho} \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \mathbb{D}_x \mathbf{u} - \frac{1}{c_v} \vartheta \operatorname{div}_x \mathbf{u}. \quad (3.3)$$

3.1 Minimum principle for the density

The formulation (3.1)–(3.3) anticipates strict positivity of the density ϱ . Formally, this property follows if the same restriction is imposed on the initial data, cf. (2.15). If the velocity field is regular enough, the continuity equation yields

$$\varrho(\tau, \cdot) \geq \min_{\Omega} \varrho_0 \exp \left(- \int_0^\tau \|\operatorname{div}_x \mathbf{u}\|_{L^\infty(\Omega)} dt \right), \quad (3.4)$$

for any $\tau \in [0, T]$. The classes of strong solutions considered in this paper always allow for the Lagrangean formulation of the equation of continuity. Specifically, the velocity field always satisfies

$$\mathbf{u} \in L^1(0, T; W^{1,\infty}(\Omega, \mathbb{R}^3)). \quad (3.5)$$

In particular, the density will be always bounded below away from zero as long as this is true initially. This justifies the lower bound on the density claimed in (2.23).

3.2 Minimum principle for the temperature

Similarly, we multiply equation (3.3) on $\exp\left(\frac{1}{c_v} \int_0^\tau \|\operatorname{div}_x \mathbf{u}\|_{L^\infty(\Omega)}\right)$ obtaining

$$\begin{aligned} \varrho \partial_t \Theta + \varrho \mathbf{u} \cdot \nabla_x \Theta - \frac{\kappa}{c_v} \Delta_x \Theta &\geq 0 \text{ in } (0, T) \times \Omega, \\ \Theta(\tau, \cdot)|_{\Gamma_D} &= \exp\left(\frac{1}{c_v} \int_0^\tau \|\operatorname{div}_x \mathbf{u}\|_{L^\infty(\Omega)}\right) \vartheta_B, \\ \nabla_x \Theta(\tau, \cdot) \cdot \mathbf{n}|_{\Gamma_N} &= \exp\left(\frac{1}{c_v} \int_0^\tau \|\operatorname{div}_x \mathbf{u}\|_{L^\infty(\Omega)}\right) q_B, \end{aligned} \quad (3.6)$$

where we have set

$$\Theta = \vartheta \exp\left(\frac{1}{c_v} \int_0^\tau \|\operatorname{div}_x \mathbf{u}\|_{L^\infty(\Omega)}\right).$$

Next, consider a function F ,

$$F : \mathbb{R} \rightarrow [0, \infty) \text{ convex, } F(z) = 0 \text{ for } z \geq 0, \quad F'(z) < 0 \text{ for } z < 0.$$

Multiplying (3.6) on $F'(\Theta - \underline{\vartheta})$, where

$$\underline{\vartheta} = \min \left\{ \min_{\Omega} \vartheta_0, \min_{\Gamma_D} \vartheta_B \right\},$$

and integrating by parts, we get

$$\frac{d}{dt} \int_{\Omega} \varrho F(\Theta - \underline{\vartheta}) \, dx \leq \frac{\kappa}{c_v} \int_{\Gamma_N} q_B F'(\Theta - \underline{\vartheta}) \, d\sigma$$

Here, we have used the impermeability of the boundary $\mathbf{u} \cdot \mathbf{n}|_{\partial\Omega} = \mathbf{u}_B \cdot \mathbf{n} = 0$. Thus if $q_B \geq 0$ we conclude, similarly to (3.4),

$$\vartheta(\tau, \cdot) \geq \min \left\{ \min_{\Omega} \vartheta_0, \min_{\Gamma_D} \vartheta_B \right\} \exp \left(- \int_0^\tau \frac{1}{c_v} \|\operatorname{div}_x \mathbf{u}\|_{L^\infty(\Omega)} \, dt \right), \quad (3.7)$$

for any $\tau \in [0, T]$. In view of hypotheses (2.16), (2.18), the temperature remains strictly positive as claimed in (2.23).

4 Classes of strong solutions

Unfortunately, the proof of Theorem 2.1 cannot be performed in the function spaces of type $L^p - L^q$ in a direct manner. Instead, we have to introduce intermediate spaces of more regular strong solutions.

4.1 Hilbert space – energy framework

First, we consider the Hilbert space setting introduced by Valli [26], [27] and Valli, Zajaczkowski [28]. Specifically, the velocity and the temperature belong to the class:

$$\vartheta \in L^2(0, T; W^{4,2}(\Omega)), \quad \partial_{tt}^2 \vartheta \in L^2(0, T; L^2(\Omega)), \quad (4.1)$$

$$\mathbf{u} \in L^2(0, T; W^{4,2}(\Omega; R^3)), \quad \partial_{tt}^2 \mathbf{u} \in L^2(0, T; L^2(\Omega; R^3)). \quad (4.2)$$

In particular, given \mathbf{u} satisfying (4.2), the time evolution of the density is uniquely determined by the equation of continuity (3.1),

$$\varrho \in C([0, T]; W^{3,2}(\Omega)), \quad \partial_t \varrho \in C([0, T]; W^{2,2}(\Omega)) \quad (4.3)$$

as long as $\varrho_0 \in W^{3,2}(\Omega)$.

Accordingly, we introduce the *solution* space in the Valli class,

$$\mathcal{S}_V[0, T] = \left\{ (\varrho, \vartheta, \mathbf{u}) \mid (\varrho, \vartheta, \mathbf{u}) \text{ belong to the class (4.1) – (4.3)} \right\}.$$

The class $\mathcal{S}_V[0, T]$ endowed with the norm

$$\begin{aligned} \|(\varrho, \vartheta, \mathbf{u})\|_{\mathcal{S}_V[0, T]} &= \sup_{t \in [0, T]} \|\varrho(t, \cdot)\|_{W^{3,2}(\Omega)} + \sup_{t \in [0, T]} \|\partial_t \varrho(t, \cdot)\|_{W^{2,2}(\Omega)} \\ &\quad + \|(\vartheta, \mathbf{u})\|_{L^2(0, T; W^{4,2}(\Omega; R^4))} + \|\partial_{tt}^2(\vartheta, \mathbf{u})\|_{L^2(0, T; L^2(\Omega; R^4))} \end{aligned} \quad (4.4)$$

is a Banach space. The data belong to the associated trace spaces:

$$\varrho_0 \in W^{3,2}(\Omega), \quad \vartheta_0 \in W^{3,2}(\Omega), \quad \mathbf{u}_0 \in W^{3,2}(\Omega; R^3), \quad (4.5)$$

$$\vartheta_B \in W^{\frac{7}{2}, 2}(\Gamma_D), \quad \mathbf{u}_B \in W^{\frac{7}{2}, 2}(\partial\Omega; R^3), \quad (4.6)$$

$$q_B \in W^{\frac{5}{2}, 2}(\Gamma_N). \quad (4.7)$$

Similarly to the above, we introduce the *data* spaces in the Valli class:

$$\mathcal{D}_{V, I} = \left\{ (\varrho_0, \vartheta_0, \mathbf{u}_0) \mid (\varrho_0, \vartheta_0, \mathbf{u}_0) \text{ belong to the class (4.5)} \right\},$$

with the norm

$$\|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{V, I}} = \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{W^{3,2}(\Omega; R^5)}; \quad (4.8)$$

and

$$\mathcal{D}_{V, B} = \left\{ (\vartheta_B, \mathbf{u}_B, q_B) \mid (\vartheta_B, \mathbf{u}_B, q_B) \text{ belong to the class (4.6), (4.7)} \right\},$$

with the norm

$$\|(\vartheta_B, \mathbf{u}_B, q_B)\|_{\mathcal{D}_{V, B}} = \|\vartheta_B\|_{W^{\frac{7}{2}, 2}(\Gamma_D)} + \|\mathbf{u}_B\|_{W^{\frac{7}{2}, 2}(\partial\Omega; R^3)} + \|q_B\|_{W^{\frac{5}{2}, 2}(\Gamma_N)}. \quad (4.9)$$

Note that

$$\mathcal{S}_V[0, T] \hookrightarrow C([0, T]; \mathcal{D}_{V, I}). \quad (4.10)$$

The following result was proved by Valli [27, Theorem A], see also [27, Remark 3.3].

Proposition 4.1 (Local existence - L^2 -framework). *Let $\Omega \subset R^3$ be a bounded domain of class C^4 , and let $G = G(x)$, $G \in W^{3,2}(\Omega)$ be a given potential. Let the initial data $(\varrho_0, \vartheta_0, \mathbf{u}_0)$ satisfy (4.5), and let the boundary data $(\vartheta_B, \mathbf{u}_B, q_B)$ belong to the class (4.6)–(4.7), where*

$$\min_{\Omega} \varrho_0 \geq \underline{\varrho} > 0, \quad \min_{\Omega} \vartheta_0 \geq \underline{\vartheta} > 0, \quad \min_{\Gamma_D} \vartheta_B \geq \underline{\vartheta} > 0, \quad q_B \geq 0.$$

In addition let the compatibility conditions

$$\vartheta_0|_{\Gamma_D} = \vartheta_B, \quad \mathbf{u}_0|_{\partial\Omega} = \mathbf{u}_B, \quad \nabla_x \vartheta_0 \cdot \mathbf{n}|_{\Gamma_N} = q_B, \quad (4.11)$$

$$\left(-\mathbf{u}_0 \cdot \nabla_x \mathbf{u}_0 - \frac{1}{\varrho_0} \nabla_x p(\varrho_0, \vartheta_0) + \frac{1}{\varrho_0} \operatorname{div}_x \mathbb{S}(\mathbb{D}_x \mathbf{u}_0) + \nabla_x G \right) \Big|_{\partial\Omega} = 0, \quad (4.12)$$

$$\left(-\mathbf{u}_0 \cdot \nabla_x \vartheta_0 - \frac{1}{c_v \varrho_0} \operatorname{div}_x (\kappa \nabla_x \vartheta_0) + \frac{1}{c_v \varrho_0} \mathbb{S}(\mathbb{D}_x \mathbf{u}_0) : \mathbb{D}_x \mathbf{u}_0 - \frac{1}{c_v} \vartheta_0 \operatorname{div}_x \mathbf{u}_0 \right) \Big|_{\Gamma_D} = 0, \quad (4.13)$$

$$\nabla_x \left(-\mathbf{u}_0 \cdot \nabla_x \vartheta_0 - \frac{1}{c_v \varrho_0} \operatorname{div}_x (\kappa \nabla_x \vartheta_0) + \frac{1}{c_v \varrho_0} \mathbb{S}(\mathbb{D}_x \mathbf{u}_0) : \mathbb{D}_x \mathbf{u}_0 - \frac{1}{c_v} \vartheta_0 \operatorname{div}_x \mathbf{u}_0 \right) \cdot \mathbf{n} \Big|_{\Gamma_N} = 0 \quad (4.14)$$

hold. Finally, suppose either $\Gamma_D \neq \emptyset$ or $q_B = 0$.

Then there exists $T > 0$ such that the NSF system (2.1)–(2.6), with the initial conditions (2.7), and the boundary conditions (2.8), (2.10) admits a unique solution $(\varrho, \vartheta, \mathbf{u})$ in the class (4.1)–(4.3). Moreover, the density ϱ and the temperature ϑ are strictly positive, specifically (3.4), (3.7) hold for any $\tau \in [0, T]$.

Given the data $(\varrho_0, \vartheta_0, \mathbf{u}_0; \vartheta_B, \mathbf{u}_B, q_B) \in \mathcal{D}_{V,I} \times \mathcal{D}_{V,B}$, we define

$$\begin{aligned} T_{\max,V} &= T_{\max,V}(\varrho_0, \vartheta_0, \mathbf{u}_0; \vartheta_B, \mathbf{u}_B, q_B) \\ &= \sup \left\{ T \geq 0 \mid \text{the solution of the NSF system exists in } \mathcal{S}_V[0, T] \right\}. \end{aligned}$$

By virtue of Proposition 4.1, $T_{\max,V} > 0$ for any data $(\varrho_0, \vartheta_0, \mathbf{u}_0; \vartheta_B, \mathbf{u}_B, q_B) \in \mathcal{D}_{V,I} \times \mathcal{D}_{V,B}$ satisfying the compatibility conditions (4.11)–(4.14). In general, $T_{\max,V}$ may differ from T_{\max} introduced in Theorem 2.1. As we shall see below, however, they coincide for sufficiently regular data.

One may consider data belonging to higher order Sobolev spaces and satisfying higher order compatibility conditions to obtain higher regularity of the corresponding solutions, cf. Kagei and Kawashima [17].

4.2 Mixed setting

In their quest for a framework that would accommodate vacuum in solutions, Cho and Kim [7] introduced a class of solutions of “mixed” type, specifically,

$$\begin{aligned} \vartheta &\in L^2(0, T; W^{2,q}(\Omega)) \cap C([0, T]; W^{2,2}(\Omega)), \\ \partial_t \vartheta &\in L^2(0, T; W^{1,2}(\Omega)) \cap C([0, T]; L^2(\Omega)), \end{aligned} \quad (4.15)$$

$$\begin{aligned} \mathbf{u} &\in L^2(0, T; W^{2,q}(\Omega; R^3)) \cap C([0, T]; W^{2,2}(\Omega; R^3)), \\ \partial_t \mathbf{u} &\in L^2(0, T; W^{1,2}(\Omega; R^3)) \cap C([0, T]; L^2(\Omega; R^3)), \end{aligned} \quad (4.16)$$

where $3 < q \leq 6$. The density ϱ , again uniquely determined via integration along characteristics, belongs to the class

$$\varrho \in C([0, T]; W^{1,q}(\Omega)), \quad \partial_t \varrho \in C([0, T]; L^q(\Omega)). \quad (4.17)$$

Analogously to the preceding section, we introduce the solution space

$$\mathcal{S}_{ChK} = \left\{ (\varrho, \vartheta, \mathbf{u}) \mid (\varrho, \vartheta, \mathbf{u}) \text{ belongs to the class (4.15)–(4.17)} \right\},$$

with the norm

$$\begin{aligned} \|(\varrho, \vartheta, \mathbf{u})\|_{\mathcal{S}_{ChK}[0,T]} &= \sup_{t \in [0, T]} \|\varrho(t, \cdot)\|_{W^{1,q}(\Omega)} + \sup_{t \in [0, T]} \|\partial_t \varrho(t, \cdot)\|_{L^q(\Omega)} \\ &+ \sup_{t \in [0, T]} \|(\vartheta, \mathbf{u})(t, \cdot)\|_{W^{2,2}(\Omega; R^4)} + \|(\vartheta, \mathbf{u})\|_{L^2(0, T; W^{2,q}(\Omega; R^4))} \\ &+ \sup_{t \in [0, T]} \|(\partial_t \vartheta, \partial_t \mathbf{u})(t, \cdot)\|_{L^2(\Omega; R^4)} + \|(\partial_t \vartheta, \partial_t \mathbf{u})\|_{L^2(0, T; W^{1,2}(\Omega; R^4))}. \end{aligned} \quad (4.18)$$

The associated data belong to the trace space

$$\varrho_0 \in W^{1,q}(\Omega), \quad \vartheta_0 \in W^{2,2}(\Omega), \quad \mathbf{u}_0 \in W^{2,2}(\Omega; R^3), \quad (4.19)$$

$$\mathbf{u}_B \in W^{2-\frac{1}{q}, q}(\partial\Omega), \quad \vartheta_B \in W^{2-\frac{1}{q}, q}(\Gamma_D), \quad (4.20)$$

$$q_B \in W^{1-\frac{1}{q}, q}(\Gamma_N). \quad (4.21)$$

We introduce the data spaces in the Cho-Kim class

$$\mathcal{D}_{ChK,I} = \left\{ (\varrho_0, \vartheta_0, \mathbf{u}_0) \mid (\varrho_0, \vartheta_0, \mathbf{u}_0) \text{ belong to the class (4.19)} \right\},$$

with the norm

$$\|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK,I}} = \|\varrho_0\|_{W^{1,q}(\Omega)} + \|\mathbf{u}_0\|_{W^{2,2}(\Omega; R^3)} + \|\vartheta_0\|_{W^{2,2}(\Omega)}, \quad (4.22)$$

and

$$\mathcal{D}_{ChK,B} = \left\{ (\vartheta_B, \mathbf{u}_B, q_B) \mid (\vartheta_B, \mathbf{u}_B, q_B) \text{ belong to the class (4.20), (4.21)} \right\},$$

with the norm

$$\|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK,B}} = \|\vartheta_B\|_{W^{2-\frac{1}{q}, q}(\Gamma_D)} + \|\mathbf{u}_B\|_{W^{2-\frac{1}{q}, q}(\Omega; R^3)} + \|q_B\|_{W^{1-\frac{1}{q}, q}(\Gamma_N)}. \quad (4.23)$$

As a matter of fact, Cho and Kim [7, Theorem 2.1] proved a local existence of solutions in the above class for the purely Dirichlet problem $\Gamma_N = \emptyset$, $\mathbf{u}_B = 0$, and $\vartheta_B = 0$, where the last condition is rather unphysical. As a by product of our analysis, we extend this result to the inhomogeneous and physically relevant boundary conditions, see Theorem 6.2 below.

4.3 $L^p - L^q$ setting

Finally, we introduce the notation relevant to the $L^p - L^q$ framework used in Theorem 2.1. First observe that the velocity field \mathbf{u} belongs to the space

$$L^p(0, T; W^{2,q}(\Omega; \mathbb{R}^3)), \quad W^{2,q} \hookrightarrow C^{1+\beta} \text{ as } q > 3.$$

In particular, the density may be recovered by the method of characteristics:

$$\varrho \in C([0, T]; W^{1,q}(\Omega)), \quad \partial_t \varrho \in L^p(0, T; L^q(\Omega)) \quad (4.24)$$

as long as $\varrho_0 \in W^{1,q}(\Omega)$, cf. Danchin [9].

Similarly to the preceding sections, we introduce the solution space

$$\mathcal{S}_{p,q}[0, T] = \left\{ (\varrho, \vartheta, \mathbf{u}) \mid (\varrho, \vartheta, \mathbf{u}) \text{ belong to the class (2.11) - (2.13)} \right\}.$$

We define a norm on $\mathcal{S}_{p,q}[0, T]$ as

$$\begin{aligned} \|(\varrho, \vartheta, \mathbf{u})\|_{\mathcal{S}_{p,q}[0, T]} &= \sup_{t \in [0, T]} \|\varrho(t, \cdot)\|_{W^{1,q}(\Omega)} + \|(\vartheta, \mathbf{u})\|_{L^p(0, T; W^{2,q}(\Omega; \mathbb{R}^4))} + \|\partial_t(\varrho, \vartheta, \mathbf{u})\|_{L^p(0, T; L^q(\Omega; \mathbb{R}^5))} \\ &\quad + \|(\vartheta, \mathbf{u})(0, \cdot)\|_{B_{q,p}^{2(1-\frac{1}{p})}(\Omega; \mathbb{R}^4)}. \end{aligned} \quad (4.25)$$

Note carefully that the norm includes the norm of the initial data in the interpolation space. This fact will be exploited later in Section 6.1.

The associated space of initial data reads

$$\mathcal{D}_{p,q,I} = \left\{ (\varrho_0, \vartheta_0, \mathbf{u}_0) \mid (\varrho_0, \vartheta_0, \mathbf{u}_0) \text{ belong to the regularity class (2.15)-(2.17)} \right\},$$

with the norm

$$\|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{p,q,I}} = \|\varrho_0\|_{W^{1,q}(\Omega)} + \|(\vartheta_0, \mathbf{u}_0)\|_{B_{q,p}^{2(1-\frac{1}{p})}(\Omega; \mathbb{R}^4)}. \quad (4.26)$$

The boundary data belong to the space

$$\mathcal{D}_{p,q,B} = \left\{ (\vartheta_B, \mathbf{u}_B, q_B) \mid (\vartheta_B, \mathbf{u}_B, q_B) \text{ belong to the regularity class (2.18)-(2.20)} \right\},$$

with the norm

$$\|(\vartheta_B, \mathbf{u}_B, q_B)\|_{\mathcal{D}_{p,q,B}} = \|\vartheta_B\|_{W^{2-\frac{1}{q},q}(\Gamma_D; \mathbb{R})} + \|\mathbf{u}_B\|_{W^{2-\frac{1}{q},q}(\partial\Omega; \mathbb{R}^3)} + \|q_B\|_{W^{1-\frac{1}{q},q}(\Gamma_N)}.$$

Recall that

$$\mathcal{S}_{p,q}[0, T] \hookrightarrow C([0, T]; \mathcal{D}_{p,q,I}). \quad (4.27)$$

Kotschote [18, Theorem 2.1] showed the local existence for $p = q > 3$, with certain restrictions imposed on the boundary heat flux q_B . Theorem 2.1 extends his result to the general case $p \neq q$ and a larger class of heat fluxes imposed on Γ_N . In particular, we propose an alternative method of proof in the case $p = q$.

5 Conditional regularity estimates

Having collected the necessary preliminary material, we deduce several conditional regularity estimates on the strong solution belonging to the regularity class used by Valli.

First, we need a suitable extension of boundary data. Specifically, we set

$$\operatorname{div}_x \mathbb{S}(\nabla_x \tilde{\mathbf{u}}_B) = 0 \text{ in } \Omega, \quad \tilde{\mathbf{u}}_B|_{\partial\Omega} = \mathbf{u}_B, \quad (5.1)$$

and

$$\begin{aligned} \Delta_x \tilde{\vartheta}_B &= 0 \text{ in } \Omega, \quad \tilde{\vartheta}_B|_{\Gamma_D} = \vartheta_B, \quad \nabla_x \tilde{\vartheta}_B \cdot \mathbf{n}|_{\Gamma_N} = q_B, \text{ if } \Gamma_D \neq \emptyset, \\ \tilde{\vartheta}_B &= \underline{\vartheta} = \inf_{\Omega} \vartheta_0 \text{ if } \Gamma_D = \emptyset. \end{aligned} \quad (5.2)$$

Note that, by virtue of hypothesis (2.18), $\tilde{\vartheta}_B > 0$ in $\bar{\Omega}$. In what follows, we shall identify the boundary data with their extension, $\vartheta_B \approx \tilde{\vartheta}_B$, $\mathbf{u}_B \approx \tilde{\mathbf{u}}_B$.

5.1 Conditional regularity in the mixed and Hilbertian setting

The following result represents a variant of the conditional regularity estimates proved in [5]. Here and hereafter, the symbol Λ denotes a generic positive and non-decreasing function, $\Lambda : [0, \infty) \rightarrow (0, \infty)$.

Proposition 5.1 (Conditional regularity - mixed and Hilbertian framework). *Let the hypotheses of Proposition 4.1 be satisfied. Let $(\varrho, \vartheta, \mathbf{u})$ be the unique strong solution of NSF system defined on some time interval $(0, T)$.*

Then there exists a non-decreasing (positive) function $\Lambda : [0, \infty) \rightarrow (0, \infty)$ such that the following estimates hold:

•

$$\begin{aligned} & \|(\varrho, \vartheta, \mathbf{u})\|_{\mathcal{S}_{ChK}[0, T]} + \sup_{t \in [0, T]} \left\| \left(\frac{1}{\varrho}, \frac{1}{\vartheta} \right) (t, \cdot) \right\|_{C(\bar{\Omega}; R^2)} \\ & \leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK, I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK, B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega}; R^2)} + \|G\|_{W^{2,2}(\Omega)} \right. \\ & \quad \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{C([0, T] \times \bar{\Omega}; R^5)} \right), \quad 3 < q \leq 6; \end{aligned} \quad (5.3)$$

•

$$\begin{aligned} & \|(\varrho, \vartheta, \mathbf{u})\|_{\mathcal{S}_V[0, T]} + \sup_{t \in [0, T]} \left\| \left(\frac{1}{\varrho}, \frac{1}{\vartheta} \right) (t, \cdot) \right\|_{C(\bar{\Omega}; R^2)} \\ & \leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{V, I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{V, B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega}; R^2)} + \|G\|_{W^{3,2}(\Omega)} \right. \\ & \quad \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{C([0, T] \times \bar{\Omega}; R^5)} \right). \end{aligned} \quad (5.4)$$

Remark 5.2. The estimate (5.4) was basically shown in [5], see also [1]. The main novelty is the estimate (5.3) in the class of lower regularity that will be exploited later for proving local existence in the mixed setting.

Remark 5.3. Roughly speaking, the conclusion of Proposition 5.1 can be seen as *a priori* estimates for the NSF system in terms of the data and the L^∞ -norm of the solution.

The rest of the section is devoted to the proof of Proposition 5.1. We proceed in several steps.

5.1.1 Energy estimates

Any bounded solution $(\varrho, \vartheta, \mathbf{u})$ of the NSF system satisfies the energy bounds

$$\begin{aligned} & \int_0^T \left(\|\mathbf{u}\|_{W^{1,2}(\Omega; R^3)}^2 + \|\vartheta\|_{W^{1,2}(\Omega)}^2 \right) dt \\ & \leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK,B}} + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0,T) \times \Omega)} \right). \end{aligned} \quad (5.5)$$

Indeed we multiply the momentum equation (2.2) on $\mathbf{u} - \mathbf{u}_B$ and integrate the resulting expression by parts obtaining

$$\begin{aligned} & \frac{d}{dt} \frac{1}{2} \int_{\Omega} \varrho |\mathbf{u} - \mathbf{u}_B|^2 dx + \int_{\Omega} \mathbb{S}(\mathbb{D}_x \mathbf{u} - \mathbb{D}_x \mathbf{u}_B) : (\mathbb{D}_x \mathbf{u} - \mathbb{D}_x \mathbf{u}_B) dx \\ & = \int_{\Omega} \varrho \mathbf{u} \cdot \nabla_x \mathbf{u}_B \cdot (\mathbf{u} - \mathbf{u}_B) dx + \int_{\Omega} p(\varrho, \vartheta) \operatorname{div}_x (\mathbf{u} - \mathbf{u}_B) dx + \int_{\Omega} \varrho \nabla_x G \cdot (\mathbf{u} - \mathbf{u}_B) dx, \end{aligned} \quad (5.6)$$

where, by virtue of Korn–Poincaré inequality,

$$\int_{\Omega} \mathbb{S}(\mathbb{D}_x \mathbf{u} - \mathbb{D}_x \mathbf{u}_B) : (\mathbb{D}_x \mathbf{u} - \mathbb{D}_x \mathbf{u}_B) dx \gtrsim \|\mathbf{u} - \mathbf{u}_B\|_{W_0^{1,2}(\Omega; R^3)}^2. \quad (5.7)$$

Thus the bound for \mathbf{u} claimed in (5.5) results from integrating (5.6) in time.

Similarly, we multiply the heat equation (2.3) on $(\vartheta - \vartheta_B)$ to deduce

$$\begin{aligned} & c_v \frac{d}{dt} \int_{\Omega} \varrho |\vartheta - \vartheta_B|^2 dx + \kappa \int_{\Omega} |\nabla_x \vartheta - \nabla_x \vartheta_B|^2 dx \\ & = c_v \int_{\Omega} \varrho (\vartheta - \vartheta_B) \mathbf{u} \cdot \nabla_x \vartheta_B dx + \int_{\Omega} (\vartheta - \vartheta_B) \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \mathbb{D}_x \mathbf{u} dx - \int_{\Omega} (\vartheta - \vartheta_B) p(\varrho, \vartheta) \operatorname{div}_x \mathbf{u} dx. \end{aligned} \quad (5.8)$$

Combining (5.8) with the previous estimate on $\nabla_x \mathbf{u}$ we complete the proof of (5.5).

5.1.2 Estimates on the material derivative

Similarly to [5, Sections 4-6], we adapt the arguments of Fang, Zi, and Zhang [13, Section 3] to the present choice of boundary conditions.

We start by introducing the material derivative

$$D_t g = \partial_t g + \mathbf{u} \cdot \nabla_x g.$$

The momentum equation (2.2) takes the form

$$\varrho D_t \mathbf{u} + \nabla_x p = \operatorname{div}_x \mathbb{S}(\mathbb{D}_x \mathbf{u}) + \varrho \nabla_x G. \quad (5.9)$$

Following [5, Section 4.1] we multiply (5.9) on $D_t(\mathbf{u} - \mathbf{u}_B)$ and perform by parts integration. This step is exactly the same as in [5, Section 4.1] yielding finally the inequality

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega} \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \mathbb{D}_x \mathbf{u} \, dx - \frac{d}{dt} \int_{\Omega} p \operatorname{div}_x \mathbf{u} \, dx + \frac{1}{2} \int_{\Omega} \varrho |D_t \mathbf{u}|^2 \, dx \\ & \leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK,B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\overline{\Omega}; R^2)} + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0,T) \times \Omega; R^5)} \right) \times \\ & \times \left(1 + \int_{\Omega} \varrho |D_t \vartheta| |\nabla_x \mathbf{u}| \, dx + \int_{\Omega} |\nabla_x \mathbf{u}|^3 \, dx + \|G\|_{W^{2,2}(\Omega)} \right). \end{aligned} \quad (5.10)$$

Next, following the original idea of Hoff [16] (cf. also [13, Section 3, Lemma 3.3]), we compute the material derivative of the momentum equation (2.2):

$$\begin{aligned} & \varrho D_t^2 \mathbf{u} + \nabla_x \partial_t p + \operatorname{div}_x (\nabla_x p \otimes \mathbf{u}) \\ & = \mu \left(\Delta_x \partial_t \mathbf{u} + \operatorname{div}_x (\Delta_x \mathbf{u} \otimes \mathbf{u}) \right) + \left(\lambda + \frac{\mu}{3} \right) \left(\nabla_x \operatorname{div}_x \partial_t \mathbf{u} + \operatorname{div}_x ((\nabla_x \operatorname{div}_x \mathbf{u}) \otimes \mathbf{u}) \right) + \varrho \mathbf{u} \cdot \nabla_x^2 G. \end{aligned} \quad (5.11)$$

Exactly as in [5, Section 4.2], we may take the scalar product of (5.11) with $D_t(\mathbf{u} - \mathbf{u}_B)$, and, after a lengthy computation, obtaining

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega} \varrho |D_t(\mathbf{u} - \mathbf{u}_B)|^2 \, dx + \mu \int_{\Omega} |\nabla_x D_t(\mathbf{u} - \mathbf{u}_B)|^2 \, dx + \left(\lambda + \frac{\mu}{3} \right) \int_{\Omega} |\operatorname{div}_x D_t(\mathbf{u} - \mathbf{u}_B)|^2 \, dx \\ & \leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK,B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\overline{\Omega})} + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty(0,T) \times \Omega} \right) \times \\ & \times \left(1 + \int_{\Omega} \varrho |D_t \vartheta|^2 \, dx + \int_{\Omega} |\nabla_x \mathbf{u}|^4 \, dx + \int_{\Omega} \varrho |D_t \mathbf{u}|^2 \, dx + \|G\|_{W^{2,2}(\Omega)} \right). \end{aligned} \quad (5.12)$$

We close the estimates following the technique of velocity decomposition proposed in the seminal paper by Sun, Wang, and Zhang [25]. To this end, we first multiply the heat equation (2.3) on $\partial_t \vartheta$ and integrate over Ω :

$$\begin{aligned} & c_v \int_{\Omega} \varrho |D_t \vartheta|^2 \, dx + \frac{\kappa}{2} \frac{d}{dt} \int_{\Omega} |\nabla_x \vartheta|^2 \, dx - \kappa \frac{d}{dt} \int_{\Gamma_N} \vartheta q_B \, d\sigma \\ & = c_v \int_{\Omega} \varrho D_t \vartheta \, \mathbf{u} \cdot \nabla_x \vartheta \, dx - \int_{\Omega} \varrho \vartheta \operatorname{div}_x \mathbf{u} \, D_t \vartheta \, dx + \int_{\Omega} \varrho \vartheta \operatorname{div}_x \mathbf{u} \, \mathbf{u} \cdot \nabla_x \vartheta \, dx \\ & + \frac{d}{dt} \int_{\Omega} \vartheta \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \nabla_x \mathbf{u} \, dx \\ & - \mu \int_{\Omega} \vartheta \left(\nabla_x \mathbf{u} + \nabla_x^t \mathbf{u} - \frac{2}{3} \operatorname{div}_x \mathbf{u} \mathbb{I} \right) : \left(\nabla_x \partial_t \mathbf{u} + \nabla_x^t \partial_t \mathbf{u} - \frac{2}{3} \operatorname{div}_x \partial_t \mathbf{u} \mathbb{I} \right) \, dx \\ & - 2\lambda \int_{\Omega} \vartheta \operatorname{div}_x \mathbf{u} \operatorname{div}_x \partial_t \mathbf{u} \, dx. \end{aligned} \quad (5.13)$$

Here, the boundary integral corresponding to the boundary condition (2.10) reads

$$\begin{aligned} \int_{\Omega} \Delta_x \vartheta \partial_t \vartheta \, dx &= \int_{\partial\Omega} \partial_t \vartheta \nabla_x \vartheta \cdot \mathbf{n} \, d\sigma - \frac{1}{2} \frac{d}{dt} \int_{\Omega} |\nabla_x \vartheta|^2 \, dx \\ &= \frac{d}{dt} \int_{\Gamma_N} \vartheta q_B \, d\sigma - \frac{1}{2} \frac{d}{dt} \int_{\Omega} |\nabla_x \vartheta|^2 \, dx. \end{aligned}$$

It turns out that we may close the estimates exactly as in [5, Sections 4-6].

$$\begin{aligned} &\sup_{t \in [0, T]} \left(\|\mathbf{u}(t, \cdot)\|_{W^{1,2}(\Omega; R^3)} + \|\sqrt{\varrho} D_t \mathbf{u}(t, \cdot)\|_{L^2(\Omega; R^3)} + \|\vartheta(t, \cdot)\|_{W^{1,2}(\Omega)} + \|\nabla_x \mathbf{u}(t, \cdot)\|_{L^4(\Omega; R^{3 \times 3})} \right) \\ &\quad + \int_0^T \int_{\Omega} |\nabla_x D_t \mathbf{u}|^2 \, dx \, dt + \int_0^T \int_{\Omega} \varrho |D_t \vartheta|^2 \, dx \, dt + \int_0^T \|\vartheta\|_{W^{2,2}(\Omega)}^2 \, dt \\ &\leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK, I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK, B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega})} \right. \\ &\quad \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0, T) \times \bar{\Omega}; R^5)} + \|G\|_{W^{2,2}(\Omega)} \right). \end{aligned}$$

Combining the previous estimates as in [5, Sections 5,6], we additionally deduce that

$$\begin{aligned} &\sup_{t \in [0, T]} \left(\|\varrho(t, \cdot)\|_{W^{1,q}(\Omega)} + \|\partial_t \varrho(t, \cdot)\|_{L^q(\Omega)} \right) + \int_0^T \|\mathbf{u}\|_{W^{2,q}(\Omega; R^3)}^2 \, dt \\ &\leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK, I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK, B}} + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0, T) \times \bar{\Omega}; R^5)} + \|G\|_{W^{2,2}(\Omega)} \right). \end{aligned}$$

Moreover, from the fact that

$$\operatorname{div}_x \mathbb{S}(\mathbb{D}_x(\mathbf{u} - \mathbf{u}_B)) = \varrho D_t \mathbf{u} + \vartheta \nabla_x \varrho + \varrho \nabla_x \vartheta - \varrho \nabla_x G$$

and from the classical regularity theory for elliptic equations, we recover that

$$\begin{aligned} &\sup_{t \in [0, T]} \|\mathbf{u}(t, \cdot)\|_{W^{2,2}(\Omega; R^3)} \\ &\lesssim \sup_{t \in [0, T]} \left(\|\sqrt{\varrho} D_t \mathbf{u}(t, \cdot)\|_{L^2(\Omega; R^3)} + \|\nabla_x \varrho(t, \cdot)\|_{L^2(\Omega; R^3)} + \|\nabla_x \vartheta(t, \cdot)\|_{L^2(\Omega; R^3)} + \|\nabla_x G\|_{L^2(\Omega; R^3)} \right) \\ &\leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK, I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK, B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega})} \right. \\ &\quad \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0, T) \times \bar{\Omega}; R^5)} + \|G\|_{W^{2,2}(\Omega)} \right). \end{aligned}$$

We can now use the minimum principle for the density (3.4) to deduce the desired estimates in terms of ϱ and \mathbf{u} appearing in (5.3). To get the remaining estimates for ϑ , we first have to take the time derivative

of equation (2.3), multiply the resulting expression on $\partial_t \vartheta$ and integrate it over Ω , obtaining

$$\begin{aligned}
& c_v \frac{d}{dt} \int_{\Omega} \varrho |\partial_t \vartheta|^2 \, dx + \kappa \int_{\Omega} |\nabla_x \partial_t \vartheta|^2 \, dx \\
&= -c_v \int_{\Omega} \partial_t \varrho |\partial_t \vartheta|^2 \, dx - c_v \int_{\Omega} \partial_t \varrho \partial_t \vartheta \mathbf{u} \cdot \nabla_x \vartheta \, dx - c_v \int_{\Omega} \varrho \partial_t \vartheta \partial_t \mathbf{u} \cdot \nabla_x \vartheta \, dx \\
&\quad + 2 \int_{\Omega} \partial_t \vartheta \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \nabla_x \partial_t \mathbf{u} \, dx \\
&\quad - \int_{\Omega} \vartheta \partial_t \varrho \partial_t \vartheta \operatorname{div}_x \mathbf{u} \, dx - \int_{\Omega} \varrho |\partial_t \vartheta|^2 \operatorname{div}_x \mathbf{u} \, dx - \int_{\Omega} \varrho \vartheta \partial_t \vartheta \operatorname{div}_x \partial_t \mathbf{u} \, dx
\end{aligned} \tag{5.14}$$

By a Grönwall argument, using the previously achieved estimates, it is not difficult to deduce

$$\begin{aligned}
& \sup_{t \in [0, T]} \|\partial_t \vartheta(t, \cdot)\|_{L^2(\Omega)} + \int_0^T \|\partial_t \vartheta\|_{W^{1,2}(\Omega)}^2 \, dt \\
&\leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK,B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega}; \mathbb{R}^2)} \right. \\
&\quad \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0, T) \times \bar{\Omega}; \mathbb{R}^5)} + \|G\|_{W^{2,2}(\Omega)} \right).
\end{aligned}$$

Indeed, denoting with I_k , $k = 1, \dots, 7$, the integrals appearing on the right-hand side of (5.14), we have

$$\begin{aligned}
|I_1| &\lesssim \left\| \frac{1}{\varrho} \right\|_{L^\infty} \|\partial_t \varrho\|_{L^3} \|\sqrt{\varrho} \partial_t \vartheta\|_{L^2} \|\partial_t \vartheta\|_{L^6} \leq \varepsilon \|\nabla_x \partial_t \vartheta\|_{L^2}^2 + C(\varepsilon) \left\| \frac{1}{\varrho} \right\|_{L^\infty}^2 \|\partial_t \varrho\|_{L^q}^2 \|\sqrt{\varrho} \partial_t \vartheta\|_{L^2}^2 \\
|I_2| &\lesssim \|\partial_t \varrho\|_{L^3} \|\partial_t \vartheta\|_{L^6} \|\nabla_x \vartheta\|_{L^2} \leq \varepsilon \|\nabla_x \partial_t \vartheta\|_{L^2}^2 + C(\varepsilon) \|\partial_t \varrho\|_{L^q}^2 \|\vartheta\|_{W^{1,2}}^2 \\
|I_3| &\lesssim \|\partial_t \vartheta\|_{L^6} \|\partial_t \mathbf{u}\|_{L^3} \|\nabla_x \vartheta\|_{L^2} \leq \varepsilon \|\nabla_x \partial_t \vartheta\|_{L^2}^2 + C(\varepsilon) \|\vartheta\|_{W^{1,2}}^2 \|\nabla_x \partial_t \mathbf{u}\|_{L^2}^2 \\
|I_4| &\lesssim \|\partial_t \vartheta\|_{L^6} \|\nabla_x \mathbf{u}\|_{L^3} \|\nabla_x \partial_t \mathbf{u}\|_{L^2} \leq \varepsilon \|\nabla_x \partial_t \vartheta\|_{L^2}^2 + C(\varepsilon) \|\mathbf{u}\|_{W^{2,2}}^2 \|\nabla_x \partial_t \mathbf{u}\|_{L^2}^2 \\
|I_5| &\lesssim \|\partial_t \varrho\|_{L^3} \|\partial_t \vartheta\|_{L^6} \|\nabla_x \mathbf{u}\|_{L^2} \leq \varepsilon \|\nabla_x \partial_t \vartheta\|_{L^2}^2 + C(\varepsilon) \|\partial_t \varrho\|_{L^q}^2 \|\mathbf{u}\|_{W^{1,2}}^2 \\
|I_6| &\lesssim \|\sqrt{\varrho} \partial_t \vartheta\|_{L^2} \|\partial_t \vartheta\|_{L^6} \|\nabla_x \mathbf{u}\|_{L^3} \leq \varepsilon \|\nabla_x \partial_t \vartheta\|_{L^2}^2 + C(\varepsilon) \|\mathbf{u}\|_{W^{2,2}}^2 \|\sqrt{\varrho} \partial_t \vartheta\|_{L^2}^2 \\
|I_7| &\lesssim \|\sqrt{\varrho} \partial_t \vartheta\|_{L^2} \|\nabla_x \partial_t \mathbf{u}\|_{L^2} \lesssim \|\nabla_x \partial_t \mathbf{u}\|_{L^2}^2 + \|\sqrt{\varrho} \partial_t \vartheta\|_{L^2}^2
\end{aligned}$$

Using once again the regularity theory for elliptic equations and the fact that the Sobolev embeddings $W^{1,2}(\Omega) \hookrightarrow L^q(\Omega)$, $W^{1,q}(\Omega) \hookrightarrow L^\infty(\Omega)$ for $3 < q \leq 6$ lead to

$$\begin{aligned}
\int_0^T \|\nabla_x \mathbf{u} : \nabla_x \mathbf{u}\|_{L^q(\Omega)}^2 \, dt &\leq \int_0^T \|\nabla_x \mathbf{u}\|_{L^q(\Omega; \mathbb{R}^{3 \times 3})}^2 \|\nabla_x \mathbf{u}\|_{L^\infty(\Omega; \mathbb{R}^{3 \times 3})}^2 \, dt \\
&\lesssim \int_0^T \|\nabla_x \mathbf{u}\|_{W^{1,2}(\Omega; \mathbb{R}^{3 \times 3})}^2 \|\nabla_x \mathbf{u}\|_{W^{1,q}(\Omega; \mathbb{R}^{3 \times 3})}^2 \, dt \\
&\lesssim \sup_{t \in [0, T]} \|\mathbf{u}(t, \cdot)\|_{W^{2,2}(\Omega; \mathbb{R}^3)}^2 \int_0^T \|\mathbf{u}\|_{W^{2,q}(\Omega; \mathbb{R}^3)}^2 \, dt,
\end{aligned}$$

we get the extra regularity for the temperature ϑ :

$$\begin{aligned} & \sup_{t \in [0, T]} \|\vartheta(t, \cdot)\|_{W^{2,2}(\Omega)} + \int_0^T \|\vartheta\|_{W^{2,q}(\Omega)}^2 dt \\ & \leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK,B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega}; R^2)} \right. \\ & \quad \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0,T) \times \bar{\Omega}; R^5)} + \|G\|_{W^{2,2}(\Omega)} \right). \end{aligned}$$

Indeed,

$$\begin{aligned} \sup_{t \in [0, T]} \|\vartheta(t, \cdot)\|_{W^{2,2}} & \lesssim \sup_{t \in [0, T]} (\|\partial_t \vartheta(t, \cdot)\|_{L^2} + \|\nabla_x \vartheta(t, \cdot)\|_{L^2} + \|\nabla_x \mathbf{u}(t, \cdot)\|_{L^4}^2 + \|\nabla_x \mathbf{u}(t, \cdot)\|_{L^2}) \\ & \lesssim \sup_{t \in [0, T]} (\|\partial_t \vartheta(t, \cdot)\|_{L^2} + \|\vartheta(t, \cdot)\|_{W^{1,2}} + \|\mathbf{u}(t, \cdot)\|_{W^{2,2}}^2 + \|\mathbf{u}(t, \cdot)\|_{W^{1,2}}), \end{aligned}$$

and

$$\begin{aligned} \int_0^T \|\vartheta\|_{W^{2,q}}^2 dt & \lesssim \int_0^T (\|\partial_t \vartheta\|_{L^q}^2 + \|\nabla_x \vartheta\|_{L^q}^2 + \|\nabla_x \mathbf{u}\|_{L^q}^2 + \|\nabla_x \mathbf{u}\|_{L^q}^2) dt \\ & \lesssim \int_0^T (\|\partial_t \vartheta\|_{W^{1,2}}^2 + \|\vartheta\|_{W^{1,q}}^2 + \|\nabla_x \mathbf{u}\|_{L^\infty}^2 \|\nabla_x \mathbf{u}\|_{L^q}^2 + \|\mathbf{u}\|_{W^{1,q}}^2) dt \\ & \lesssim \int_0^T (\|\partial_t \vartheta\|_{W^{1,2}}^2 + \|\vartheta\|_{W^{2,2}}^2 + \|\mathbf{u}\|_{W^{2,q}}^2 \|\mathbf{u}\|_{W^{2,2}}^2 + \|\mathbf{u}\|_{W^{2,2}}^2) dt. \end{aligned}$$

Putting all the estimates together and using the minimum principle for the temperature (3.7), we end up to

$$\begin{aligned} & \sup_{t \in [0, T]} (\|\varrho(t, \cdot)\|_{W^{1,q}(\Omega)} + \|\partial_t \varrho(t, \cdot)\|_{L^q(\Omega)}) \\ & + \sup_{t \in [0, T]} (\|\mathbf{u}(t, \cdot)\|_{W^{2,2}(\Omega; R^3)} + \|\partial_t \mathbf{u}(t, \cdot)\|_{L^2(\Omega; R^3)}) + \sup_{t \in [0, T]} (\|\vartheta(t, \cdot)\|_{W^{2,2}(\Omega)} + \|\partial_t \vartheta(t, \cdot)\|_{L^2(\Omega)}) \\ & + \int_0^T \|\mathbf{u}\|_{W^{2,q}(\Omega; R^3)}^2 dt + \int_0^T \|\partial_t \mathbf{u}\|_{W^{1,2}(\Omega; R^3)}^2 dt + \int_0^T \|\vartheta\|_{W^{2,q}(\Omega)}^2 dt + \int_0^T \|\partial_t \vartheta\|_{W^{1,2}(\Omega)}^2 dt \\ & + \sup_{t \in [0, T]} \left\| \left(\frac{1}{\varrho}, \frac{1}{\vartheta} \right) (t, \cdot) \right\|_{C(\bar{\Omega}; R^2)} \\ & \leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK,B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega}; R^2)} \right. \\ & \quad \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0,T) \times \bar{\Omega}; R^5)} + \|G\|_{W^{2,2}(\Omega)} \right). \end{aligned} \tag{5.15}$$

This is nothing other than (5.3).

The proof of (5.4) can be obtained exactly as in [5] passing to higher order derivatives. We have therefore completed the proof of Proposition 5.1.

5.2 Conditional regularity estimates in the $L^p - L^q$ setting

The reader will have noticed that the conditional regularity estimates in terms of the norm $\|(\varrho, \vartheta, \mathbf{u})\|_{C([0,T] \times \bar{\Omega}; R^5)}$ require differentiating the field equations in time. Unfortunately, this cannot be performed in a direct manner within the $L^p - L^q$ framework. Instead, we report the following auxiliary result.

Proposition 5.4 (Conditional regularity estimates - $L^p - L^q$ framework). *Let the hypotheses of Proposition 4.1 be satisfied. Let $(\varrho, \vartheta, \mathbf{u})$ be the unique solution of NSF system defined on some time interval $(0, T)$. Let*

$$3 < q < \infty, \quad \frac{2q}{2q-3} < p < \infty. \quad (5.16)$$

Then there exists a non-decreasing (positive) function $\Lambda : [0, \infty) \rightarrow (0, \infty)$ such that the following estimate

$$\begin{aligned} & \|(\varrho, \vartheta, \mathbf{u})\|_{\mathcal{S}_{p,q}[0,\tau]} + \sup_{t \in [0,\tau]} \left\| \left(\frac{1}{\varrho}, \frac{1}{\vartheta} \right) (t, \cdot) \right\|_{C(\bar{\Omega}; R^2)} \\ & \leq \Lambda \left(\|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{p,q,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{p,q,B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega}; R^2)} + \|G\|_{W^{1,q}(\Omega)} \right. \\ & \quad \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{C([0,\tau] \times \bar{\Omega}; R^5)} + \int_0^\tau \|(\vartheta, \mathbf{u})\|_{W^{1,\infty}(\Omega; R^4)}^p dt \right) \end{aligned} \quad (5.17)$$

holds for any $0 \leq \tau \leq T_{\text{eff}} \wedge T$, where

$$\begin{aligned} T_{\text{eff}} \geq \Lambda^{-1} & \left(\|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{p,q,I}} + \left\| \frac{1}{\varrho_0} \right\|_{C(\bar{\Omega})} + \|G\|_{W^{1,q}(\Omega)} \right. \\ & \left. + \sup_{t \in [0,T]} \|\vartheta\|_{C(\bar{\Omega})} + \|\mathbf{u}\|_{L^p(0,T; W^{1,\infty}(\Omega; R^3))} \right) > 0. \end{aligned} \quad (5.18)$$

Remark 5.5. Obviously, the conclusion of Proposition 5.4 is “weaker” than that of Proposition 5.1 as the right-hand side of (5.17) depends on the norm $\|(\vartheta, \mathbf{u})\|_{L^p(0,T; W^{1,\infty}(\Omega; R^3))}$, and the result is “local” in view of (5.18). We will remove this drawback in the last part of the paper.

Proof. First, we rewrite the system (3.1)–(3.3) in the form

$$\partial_t b(\varrho) + \mathbf{u} \cdot \nabla_x b(\varrho) = -b'(\varrho) \varrho \operatorname{div}_x \mathbf{u}, \quad (5.19)$$

$$\begin{aligned} \partial_t(\mathbf{u} - \mathbf{u}_B) - \frac{1}{\varrho_0} \operatorname{div}_x \mathbb{S}(\mathbb{D}_x(\mathbf{u} - \mathbf{u}_B)) &= \left(\frac{1}{\varrho} - \frac{1}{\varrho_0} \right) \operatorname{div}_x \mathbb{S}(\mathbb{D}_x(\mathbf{u} - \mathbf{u}_B)) \\ &\quad - \mathbf{u} \cdot \nabla_x \mathbf{u} - \vartheta \nabla_x \log(\varrho) - \nabla_x \vartheta + \nabla_x G, \end{aligned} \quad (5.20)$$

$$\begin{aligned} \partial_t(\vartheta - \vartheta_B) - \frac{\kappa}{c_v \varrho_0} \Delta_x(\vartheta - \vartheta_B) &= \left(\frac{\kappa}{c_v \varrho} - \frac{\kappa}{c_v \varrho_0} \right) \Delta_x(\vartheta - \vartheta_B) - \mathbf{u} \cdot \nabla_x \vartheta \\ &\quad + \frac{1}{c_v \varrho} \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \mathbb{D}_x \mathbf{u} - \frac{1}{c_v} \vartheta \operatorname{div}_x \mathbf{u}, \end{aligned} \quad (5.21)$$

where b in (5.19) is a continuously differentiable function, and $\mathbf{u}_B, \vartheta_B$ are the extensions of the boundary data introduced in (5.1), (5.2). In particular, the functions $(\mathbf{u} - \mathbf{u}_B), (\vartheta - \vartheta_B)$ satisfy the homogeneous boundary conditions.

At this stage, we use the $L^p - L^q$ theory for parabolic equations, see Denk, Hieber, and Prüss [11, Theorem 2.3], to deduce the estimates

$$\begin{aligned}
& \int_0^T \|\partial_t(\mathbf{u} - \mathbf{u}_B)\|_{L^q(\Omega; \mathbb{R}^3)}^p dt + \int_0^T \left\| \frac{1}{\varrho_0} \operatorname{div}_x \mathbb{S}(\mathbb{D}_x(\mathbf{u} - \mathbf{u}_B)) \right\|_{L^q(\Omega; \mathbb{R}^3)}^p dt \\
& \leq \Lambda \left(\left\| \frac{1}{\varrho_0} \right\|_{C(\bar{\Omega})} + \|\varrho_0\|_{W^{1,q}(\Omega)} \right) \left[\int_0^T \left\| \left(\frac{1}{\varrho} - \frac{1}{\varrho_0} \right) \operatorname{div}_x \mathbb{S}(\mathbb{D}_x(\mathbf{u} - \mathbf{u}_B)) \right\|_{L^q(\Omega; \mathbb{R}^3)}^p dt \right. \\
& + \int_0^T \|\mathbf{u} \cdot \nabla_x \mathbf{u}\|_{L^q(\Omega; \mathbb{R}^3)}^p dt + \int_0^T \|\nabla_x \vartheta\|_{L^q(\Omega; \mathbb{R}^3)}^p dt + \int_0^T \|\nabla_x G\|_{L^q(\Omega; \mathbb{R}^3)}^p \\
& \left. + \int_0^T \|\vartheta \nabla_x \log(\varrho)\|_{L^q(\Omega; \mathbb{R}^3)}^p dt + \|\mathbf{u}_0 - \mathbf{u}_B\|_{B_{q,p}^{2(1-\frac{1}{p})}; \mathbb{R}^3}^p \right], \tag{5.22}
\end{aligned}$$

and

$$\begin{aligned}
& \int_0^T \|\partial_t(\vartheta - \vartheta_B)\|_{L^q(\Omega)}^p dt + \int_0^T \left\| \frac{\kappa}{c_v \varrho_0} \Delta_x(\vartheta - \vartheta_B) \right\|_{L^q(\Omega)}^p dt \\
& \leq \Lambda \left(\left\| \frac{1}{\varrho_0} \right\|_{C(\bar{\Omega})} + \|\varrho_0\|_{W^{1,q}(\Omega)} \right) \left[\int_0^T \left\| \left(\frac{1}{\varrho} - \frac{1}{\varrho_0} \right) \Delta_x(\vartheta - \vartheta_B) \right\|_{L^q(\Omega)}^p dt \right. \\
& + \int_0^T \left(\|\mathbf{u} \cdot \nabla_x \vartheta\|_{L^q(\Omega)}^p \right) dt + \int_0^T \left\| \frac{1}{\varrho} \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \mathbb{D}_x \mathbf{u} \right\|_{L^q(\Omega; \mathbb{R}^3)}^p dt + \int_0^T \left\| \frac{1}{c_v} \vartheta \operatorname{div}_x \mathbf{u} \right\|_{L^q(\Omega)}^p dt \\
& \left. + \int_0^T \|\vartheta_0 - \vartheta_B\|_{B_{q,p}^{2(1-\frac{1}{p})}} \right]. \tag{5.23}
\end{aligned}$$

Note that the constant Λ depends, in general, on the modulus of continuity of ϱ_0 and the ellipticity constants of the associated linear operator, see e.g. Krylov [19], Schlag [23], or Solonnikov [24].

Moreover, using (5.1), (5.2) we may rewrite the above inequalities in the form

$$\begin{aligned}
& \int_0^T \left[\|\partial_t \mathbf{u}\|_{L^q(\Omega; \mathbb{R}^3)}^p + \|\mathbf{u}\|_{W^{2,q}(\Omega; \mathbb{R}^3)}^p \right] dt \\
& \lesssim \Lambda \left(\left\| \frac{1}{\varrho_0} \right\|_{C(\bar{\Omega})} + \|\varrho_0\|_{W^{1,q}(\Omega)} \right) \left[\int_0^T \left\| \left(\frac{1}{\varrho} - \frac{1}{\varrho_0} \right) \operatorname{div}_x \mathbb{S}(\mathbb{D}_x \mathbf{u}) \right\|_{L^q(\Omega; \mathbb{R}^3)}^p dt \right. \\
& + \int_0^T \|\mathbf{u} \cdot \nabla_x \mathbf{u}\|_{L^q(\Omega; \mathbb{R}^3)}^p dt + \int_0^T \|\nabla_x \vartheta\|_{L^q(\Omega; \mathbb{R}^3)}^p dt + \int_0^T \|\nabla_x G\|_{L^q(\Omega; \mathbb{R}^3)}^p \\
& \left. + \int_0^T \|\vartheta \nabla_x \log(\varrho)\|_{L^q(\Omega; \mathbb{R}^3)}^p dt + \|\mathbf{u}_0\|_{B_{q,p}^{2(1-\frac{1}{p})}(\Omega; \mathbb{R}^3)}^p + (1+T) \|\mathbf{u}_B\|_{W^{2-\frac{1}{q},q}(\partial\Omega; \mathbb{R}^3)}^p \right], \tag{5.24}
\end{aligned}$$

and

$$\int_0^T \left[\|\partial_t \vartheta\|_{L^q(\Omega)}^p + \|\vartheta\|_{W^{2,q}(\Omega)}^p \right] dt$$

$$\begin{aligned}
&\lesssim \Lambda \left(\left\| \frac{1}{\varrho_0} \right\|_{C(\bar{\Omega})} + \|\varrho_0\|_{W^{1,q}(\Omega)} \right) \left[\int_0^T \left\| \left(\frac{1}{\varrho} - \frac{1}{\varrho_0} \right) \Delta_x \vartheta \right\|_{L^q(\Omega)}^p dt + \int_0^T \left(\|\mathbf{u} \cdot \nabla_x \vartheta\|_{L^q(\Omega)}^p \right) dt \right. \\
&+ \int_0^T \left\| \frac{1}{\varrho} \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \mathbb{D}_x \mathbf{u} \right\|_{L^q(\Omega)}^p dt + \int_0^T \|\vartheta \operatorname{div}_x \mathbf{u}\|_{L^q(\Omega)}^p dt \\
&\left. + \|\vartheta_0\|_{B_{q,p}^{2(1-\frac{1}{p})}(\Omega)} + (1+T)\|\vartheta_B\|_{W^{2-\frac{1}{q},q}(\Gamma_D)}^2 \right]. \tag{5.25}
\end{aligned}$$

Next, we use the estimate on the density gradient proved by Kotschote [18, Lemma 4.2 and comments]:

$$\sup_{\tau \in (0,T)} \|\nabla_x \varrho(\tau, \cdot)\|_{L^q(\Omega; R^3)} \lesssim \exp \left(2 \int_0^T \|\operatorname{div}_x \mathbf{u}\|_{L^\infty(\Omega)} dt \right) \left(\|\nabla_x \varrho_0\|_{L^q(\Omega; R^3)} + \int_0^T \|\mathbf{u}\|_{W^{2,q}(\Omega; R^3)} \right). \tag{5.26}$$

In particular,

$$\begin{aligned}
&\int_0^T \|\vartheta \nabla_x \log(\varrho)\|_{L^q(\Omega)}^p dt \leq T \left[\sup_{t \in (0,T)} \|\vartheta\|_{L^\infty(\Omega)} \sup_{t \in (0,T)} \left\| \frac{1}{\varrho} \right\|_{L^\infty(\Omega)} \sup_{t \in (0,T)} \|\nabla_x \varrho\|_{L^q(\Omega; R^3)}^p \right] \\
&\lesssim T \left[\sup_{t \in (0,T)} \|\vartheta\|_{L^\infty(\Omega)} \sup_{t \in (0,T)} \left\| \frac{1}{\varrho} \right\|_{L^\infty(\Omega)} \exp \left(2p \int_0^T \|\operatorname{div}_x \mathbf{u}\|_{L^\infty(\Omega)} dt \right) \times \right. \\
&\left. \times \left(\|\nabla_x \varrho_0\|_{L^q(\Omega; R^3)}^p + T^{p-1} \int_0^T \|\mathbf{u}\|_{W^{2,q}(\Omega; R^3)}^p \right) \right]. \tag{5.27}
\end{aligned}$$

Finally, computing $\frac{1}{\varrho}$ along characteristics we have

$$\begin{aligned}
\frac{1}{\varrho(t, X(t, x))} &= \frac{1}{\varrho_0(x)} \exp \left(\int_0^t \operatorname{div}_x \mathbf{u}(s, X(s, x)) ds \right), \\
X'(t, x) &= \mathbf{u}(t, X(t, x)), \quad X(0, x) = x.
\end{aligned}$$

Consequently,

$$\frac{1}{\varrho(t, X(t, x))} - \frac{1}{\varrho_0(X(t, x))} = \frac{1}{\varrho_0(x)} \left[\exp \left(\int_0^t \operatorname{div}_x \mathbf{u}(s, X(s, x)) ds \right) - 1 \right] + \frac{1}{\varrho_0(x)} - \frac{1}{\varrho_0(X(t, x))}.$$

As $\varrho_0 \in W^{1,q}(\Omega)$, $q > 3$, ϱ_0 bounded below, the function $\frac{1}{\varrho_0}$ is Hölder continuous. In particular,

$$\left| \frac{1}{\varrho_0(x)} - \frac{1}{\varrho_0(X(t, x))} \right| \lesssim |x - X(t, x)|^\beta = \left| \int_0^t \mathbf{u}(x, X(s, x)) ds \right|^\beta$$

for some $0 < \beta < 1$.

Thus for any $\delta > 0$, we may find $T_{\text{eff}} > 0$ small enough in terms of $\int_0^T \|\mathbf{u}\|_{W^{1,\infty}(\Omega; R^3)}^p$ and the norm of the data so that

$$\sup_{\tau \in (0, T_{\text{eff}})} \left\| \left(\frac{1}{\varrho(\tau, \cdot)} - \frac{1}{\varrho_0} \right) \right\|_{L^\infty(\Omega)} < \delta.$$

In particular, the integrals

$$\int_0^T \left\| \left(\frac{1}{\varrho} - \frac{1}{\varrho_0} \right) \operatorname{div}_x \mathbb{S}(\mathbb{D}_x \mathbf{u}) \right\|_{L^q(\Omega; \mathbb{R}^3)}^p \quad \text{and} \quad \int_0^T \left\| \left(\frac{1}{\varrho} - \frac{1}{\varrho_0} \right) \Delta_x \vartheta \right\|_{L^q(\Omega)}^p dt$$

can be “absorbed” by their counterparts on the left–hand side of the inequalities (5.24) and (5.25) respectively. The same applies to the last integral in (5.27) as long as $T = T_{\text{eff}}$ is small enough. These observations complete the desired bounds in terms of ϱ and \mathbf{u} .

Finally, going back to (5.25) we observe

$$\begin{aligned} & \int_0^T \left\| \frac{1}{\varrho} \mathbb{S}(\mathbb{D}_x \mathbf{u}) : \mathbb{D}_x \mathbf{u} \right\|_{L^q(\Omega)}^p \\ & \lesssim \left(\delta + \sup_{\Omega} \frac{1}{\varrho_0} \right) \int_0^T \|\nabla_x \mathbf{u}\|_{L^q(\Omega)}^{2p} dt = \left(\delta + \sup_{\Omega} \frac{1}{\varrho_0} \right) \int_0^T \|\nabla_x \mathbf{u}\|_{L^{2q}(\Omega; \mathbb{R}^3)}^{2p} dt, \end{aligned}$$

for any $0 < T \leq T_{\text{eff}}$. Using the general version of Gagliardo–Nirenberg inequality (see Brezis, Mironescu [6, Theorem 1]) we obtain

$$\|\mathbf{u}\|_{W^{s,q}(\Omega; \mathbb{R}^3)} \leq \|\mathbf{u}\|_{W^{\alpha,q}(\Omega; \mathbb{R}^3)}^{\frac{1}{2}} \|\mathbf{u}\|_{W^{2,q}(\Omega; \mathbb{R}^3)}^{\frac{1}{2}}, \quad (5.28)$$

where

$$s = 1 + \frac{\alpha}{2}, \quad \text{and} \quad \alpha < 2\left(1 - \frac{1}{p}\right) \Rightarrow B_{q,p}^{2(1-\frac{1}{p})}(\Omega; \mathbb{R}^3) \hookrightarrow B_{q,q}^{\alpha}(\Omega; \mathbb{R}^3) \approx W^{\alpha,q}(\Omega; \mathbb{R}^3).$$

Consequently, we have

$$\sup_{t \in [0, T_{\text{eff}}]} \|\mathbf{u}\|_{W^{\alpha,q}(\Omega)} \leq c(T_{\text{eff}}) \left(\|\mathbf{u}\|_{L^p(0, T_{\text{eff}}; W^{2,q}(\Omega; \mathbb{R}^3))} + \|\partial_t \mathbf{u}\|_{L^p(0, T_{\text{eff}}; L^q(\Omega; \mathbb{R}^3))} \right) \quad (5.29)$$

Note carefully that the constant $c(T_{\text{eff}})$ may blow up for $T_{\text{eff}} \rightarrow 0$. Fortunately, $T_{\text{eff}} > 0$ has already been fixed in the previous step.

Going back to (5.28), we have to show

$$W^{s,q}(\Omega; \mathbb{R}^3) \hookrightarrow W^{1,2q}(\Omega; \mathbb{R}^3) \Leftrightarrow W^{\frac{\alpha}{2},q}(\Omega; \mathbb{R}^3) \hookrightarrow L^{2q}(\Omega; \mathbb{R}^3).$$

This is true if $\frac{\alpha}{2}q > 3$. If $\frac{\alpha}{2}q < 3$, we need

$$\frac{3q}{3 - \frac{\alpha}{2}q} > 2q.$$

As $\frac{\alpha}{2}$ can be chosen arbitrarily close to $(1 - \frac{1}{p})$, the desired conclusion follows from hypothesis (5.16).

The proof is now complete. Note we have used the fact that the constants in the maximal regularity estimates (5.22), (5.23) remain the same replacing T by $T_{\text{eff}} \leq T$. □

6 Local existence of strong solutions

We are in a position to prove the local existence of strong solutions in both the $L^p - L^q$ framework claimed in Theorem 2.1 and the mixed setting for the inhomogeneous Dirichlet boundary conditions. To the best of our knowledge, these results are new. Moreover, the method can be seen as an alternative proof of known results avoiding the explicit construction of approximate schemes.

6.1 Local existence $L^p - L^q$ setting

Our goal is to show the existence of local in time solutions claimed in Theorem 2.1. The solutions are constructed via regular approximations in Valli's class. As already pointed out, this approach requires higher regularity of the underlying spatial domain. We proceed in several steps.

6.1.1 Data approximation

We consider a sequence of initial data

$$(\varrho_0^\varepsilon, \vartheta_0^\varepsilon, \mathbf{u}_0^\varepsilon) \in \mathcal{D}_{V,I} \cap \mathcal{D}_{p,q,I},$$

along with the boundary data

$$(\vartheta_B^\varepsilon, \mathbf{u}_B^\varepsilon, q_B^\varepsilon) \in \mathcal{D}_{V,B} \cap \mathcal{D}_{p,q,B},$$

satisfying the compatibility conditions (4.11)–(4.14), along with a sequence of approximate potentials

$$G^\varepsilon \in W^{3,2}(\Omega).$$

In addition, we suppose

$$\begin{aligned} (\varrho_0^\varepsilon, \vartheta_0^\varepsilon, \mathbf{u}_0^\varepsilon) &\rightarrow (\varrho_0, \vartheta_0, \mathbf{u}_0) \text{ in } \mathcal{D}_{p,q,I}, \\ (\vartheta_B^\varepsilon, \mathbf{u}_B^\varepsilon, q_B^\varepsilon) &\rightarrow (\vartheta_B, \mathbf{u}_B, q_B) \text{ in } \mathcal{D}_{p,q,B}, \\ \varrho_0^\varepsilon &\geq \underline{\varrho} > 0, \vartheta_0^\varepsilon \geq \underline{\vartheta} > 0, \vartheta_B^\varepsilon \geq \underline{\vartheta} > 0, \\ G^\varepsilon &\rightarrow G \text{ in } W^{1,q}(\Omega) \end{aligned} \tag{6.1}$$

as $\varepsilon \rightarrow 0$. Let $(\varrho_\varepsilon, \vartheta_\varepsilon, \mathbf{u}_\varepsilon)_{\varepsilon > 0}$ be the associated sequence of solutions to the NSF system in the space $\mathcal{S}_V[0, T]$, $0 < T < T_{\max, V}^\varepsilon$, the existence of which is guaranteed by Proposition 4.1.

6.1.2 Control functionals and hitting times

We start with the following auxiliary result.

Lemma 6.1. *Suppose*

$$3 < q < \infty, \frac{2q}{2q-3} < p < \infty. \tag{6.2}$$

Then the embeddings

$$\mathcal{S}_{p,q}[0, T] \hookrightarrow C([0, T] \times \bar{\Omega}; R^4), \tag{6.3}$$

and

$$L^p(0, T; W^{2,q}(\Omega)) \cap W^{1,p}(0, T; L^q(\Omega)) \hookrightarrow L^p(0, T; W^{1,\infty}(\Omega)) \tag{6.4}$$

are compact. In both cases, the embedding constant is independent of T .

Proof. First, we show

$$S_{p,q}[0, T] \hookrightarrow C\left([0, T]; W^{1,q}(\Omega) \times B_{q,p}^{2(1-\frac{1}{p})}(\Omega) \times B_{q,p}^{2(1-\frac{1}{p})}(\Omega; R^3)\right), \quad (6.5)$$

where the embedding constant is independent of T . As the embedding

$$W^{1,q}(\Omega) \times B_{q,p}^{2(1-\frac{1}{p})}(\Omega) \times B_{q,p}^{2(1-\frac{1}{p})}(\Omega; R^3) \hookrightarrow C(\overline{\Omega}; R^5)$$

is compact, compactness claimed in (6.3) follows by a direct application of the Banach space version of Arzelà–Ascoli theorem.

First note that (6.5) is obvious for the density ϱ . As for (ϑ, \mathbf{u}) , we first write

$$(\vartheta, \mathbf{u}) = (\vartheta - \tilde{\vartheta}, \mathbf{u} - \tilde{\mathbf{u}}) + (\tilde{\vartheta}, \tilde{\mathbf{u}}),$$

where

$$(\tilde{\mathbf{u}}, \tilde{\vartheta}) \in L^p(0, T; W^{2,q}(\Omega; R^4)) \cap W^{1,p}(0, T; L^q(\Omega; R^4))$$

is an extension of the initial value,

$$(\tilde{\vartheta}, \tilde{\mathbf{u}})(0, \cdot) = (\vartheta, \mathbf{u})(0, \cdot).$$

Accordingly, we get the estimate

$$\sup_{t \in (0, T)} \|(\vartheta - \tilde{\vartheta}, \mathbf{u} - \tilde{\mathbf{u}})(t, \cdot)\|_{B_{q,p}^{2(1-\frac{1}{p})}(\Omega; R^4)}^p \lesssim \int_0^T \left[\|(\vartheta - \tilde{\vartheta}, \mathbf{u} - \tilde{\mathbf{u}})\|_{W^{2,q}(\Omega; R^4)}^p + \|\partial_t(\vartheta - \tilde{\vartheta}, \mathbf{u} - \tilde{\mathbf{u}})\|_{W^{2,q}(\Omega; R^4)}^p \right] \quad (6.6)$$

As $(\vartheta - \tilde{\vartheta}, \mathbf{u} - \tilde{\mathbf{u}})(0, \cdot) = 0$, the embedding constant is independent of T , cf. Denk [10, Lemma 2.4].

Next,

$$\|(\vartheta, \mathbf{u})(0)\|_{B_{q,p}^{2(1-\frac{1}{p})}}^p = \|(\tilde{\vartheta}, \tilde{\mathbf{u}})(0)\|_{B_{q,p}^{2(1-\frac{1}{p})}}^p = \inf \int_0^\infty \left[\|\tilde{\vartheta}, \tilde{\mathbf{u}}\|_{W^{2,q}(\Omega; R^4)}^p + \|\partial_t(\tilde{\vartheta}, \tilde{\mathbf{u}})\|_{W^{2,q}(\Omega; R^4)}^p \right],$$

where the infimum is taken over all possible extensions $(\tilde{\vartheta}, \tilde{\mathbf{u}})$ of the initial data. Consequently, we may choose $(\tilde{\vartheta}, \tilde{\mathbf{u}})$ so that

$$\int_0^\infty \left[\|\tilde{\vartheta}, \tilde{\mathbf{u}}\|_{W^{2,q}(\Omega; R^4)}^p + \|\partial_t(\tilde{\vartheta}, \tilde{\mathbf{u}})\|_{W^{2,q}(\Omega; R^4)}^p \right] \leq 2 \|(\vartheta, \mathbf{u})(0)\|_{B_{q,p}^{2(1-\frac{1}{p})}}^p.$$

Going back to (6.6) we may infer that

$$\begin{aligned} & \sup_{t \in (0, T)} \|(\vartheta - \tilde{\vartheta}, \mathbf{u} - \tilde{\mathbf{u}})(t, \cdot)\|_{B_{q,p}^{2(1-\frac{1}{p})}(\Omega; R^4)}^p \\ & \lesssim \int_0^T \left[\|(\vartheta, \mathbf{u})\|_{W^{2,q}(\Omega; R^4)}^p + \|\partial_t(\vartheta, \mathbf{u})\|_{W^{2,q}(\Omega; R^4)}^p \right] + 2 \|(\vartheta, \mathbf{u})(0)\|_{B_{q,p}^{2(1-\frac{1}{p})}}^p. \end{aligned} \quad (6.7)$$

Finally, we get

$$\sup_{t \in (0, T)} \|(\tilde{\vartheta}, \tilde{\mathbf{u}})(t, \cdot)\|_{B_{q,p}^{2(1-\frac{1}{p})}(\Omega; R^4)}^p \leq \sup_{t \in (0, \infty)} \|(\tilde{\vartheta}, \tilde{\mathbf{u}})(t, \cdot)\|_{B_{q,p}^{2(1-\frac{1}{p})}(\Omega; R^4)}^p$$

$$\lesssim \int_0^\infty \left[\|\tilde{\vartheta}, \tilde{\mathbf{u}}\|_{W^{2,q}(\Omega;R^4)}^p + \|\partial_t(\tilde{\vartheta}, \tilde{\mathbf{u}})\|_{W^{2,q}(\Omega;R^4)}^p \right] \leq 2\|(\vartheta, \mathbf{u})(0)\|_{B_{q,p}^{2(1-\frac{1}{p})}}^p,$$

which completes the proof of (6.5). Note that the norm introduced in (4.25) played a crucial role in the above arguments.

Finally, (6.4) follows from the compact embedding

$$W^{2,q}(\Omega; R^3) \hookrightarrow W^{1,\infty}(\Omega; R^3)$$

combined with the Aubin–Lions compactness argument. □

Given the sequence of solutions $(\varrho_\varepsilon, \vartheta_\varepsilon, \mathbf{u}_\varepsilon)_{\varepsilon>0}$, we define a sequence of *control functionals*

$$\mathcal{F}_\varepsilon(\tau) = \|(\varrho_\varepsilon, \vartheta_\varepsilon, \mathbf{u}_\varepsilon)(\tau, \cdot)\|_{C(\bar{\Omega};R^5)} + \int_0^\tau \|(\vartheta_\varepsilon, \mathbf{u}_\varepsilon)(t, \cdot)\|_{W^{1,\infty}(\Omega;R^4)}^p dt$$

for any $0 \leq \tau < T_{\max, V}^\varepsilon$.

Next, fix $M > 0$ large enough so that

$$\mathcal{F}_\varepsilon(0) = \|(\varrho_0^\varepsilon, \vartheta_0^\varepsilon, \mathbf{u}_0^\varepsilon)\|_{C(\bar{\Omega};R^5)} < \frac{M}{2} \text{ uniformly for } \varepsilon \rightarrow 0. \quad (6.8)$$

In accordance with Lemma 6.1, $\mathcal{F}^\varepsilon \in C[0, T]$ for any $T < T_{\max, V}^\varepsilon$.

Consider the “hitting” time

$$T_M^\varepsilon = \sup_{\tau \in [0, T_{\max}^\varepsilon)} \{\mathcal{F}_\varepsilon(\tau) < M\}.$$

By virtue of (6.8), $T_M^\varepsilon > 0$, and, in accordance with the conditional regularity criterion (5.4), we conclude,

$$0 < T_M^\varepsilon < T_{\max, V}^\varepsilon.$$

Finally, we fix $T_{\text{eff}} > 0$ as in (5.18), specifically,

$$0 < T_{\text{eff}} = \Lambda^{-1} \left(\|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{p,q,I}} + \left\| \frac{1}{\varrho_0} \right\|_{C(\bar{\Omega})} + \|G\|_{W^{1,q}(\Omega)} + M \right).$$

In particular, T_{eff} is positive independent of ε .

Set

$$\bar{T}_\varepsilon = \min\{T_M^\varepsilon; T_{\text{eff}}\} > 0,$$

and

$$\bar{\mathcal{F}}_\varepsilon(\tau) = \mathcal{F}_\varepsilon(\tau \wedge \bar{T}_\varepsilon), \quad \tau \in [0, T_{\text{eff}}].$$

By virtue of the conditional regularity bound (5.17) and Lemma 6.1, the sequence $(\bar{\mathcal{F}}_\varepsilon)_{\varepsilon>0}$ is precompact in $C[0, T_{\text{eff}}]$, as it is precompact in $[0, T_M^\varepsilon]$ independently of ε .

Extracting a suitable subsequence we may assume

$$0 \leq \bar{\mathcal{F}}_\varepsilon \leq M, \quad \bar{\mathcal{F}}_\varepsilon \rightarrow \mathcal{F} \text{ in } C[0, T_{\text{eff}}], \quad \mathcal{F}(0) = \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{C(\bar{\Omega})} \leq \frac{M}{2}.$$

Consequently, there exists $0 < T \leq T_{\text{eff}}$ such that

$$0 \leq \mathcal{F}(\tau) < M \text{ for any } \tau \in [0, T] \Rightarrow 0 \leq \overline{\mathcal{F}}_\varepsilon(\tau) < M \text{ for any } \tau \in [0, T]$$

for all $\varepsilon > 0$ small enough, meaning

$$\overline{\mathcal{F}}_\varepsilon(\tau) = \mathcal{F}_\varepsilon(\tau) = \|(\varrho_\varepsilon, \vartheta_\varepsilon, \mathbf{u}_\varepsilon)(\tau, \cdot)\|_{C(\overline{\Omega}; R^5)} + \int_0^\tau \|(\vartheta_\varepsilon, \mathbf{u}_\varepsilon)(t, \cdot)\|_{W^{1,\infty}(\Omega; R^4)}^p dt < M \text{ for any } \tau \in [0, T]. \quad (6.9)$$

Finally, the estimate (6.9) along with the conditional regularity result (5.17) imply uniform bound,

$$\|(\varrho_\varepsilon, \vartheta_\varepsilon, \mathbf{u}_\varepsilon)\|_{\mathcal{S}_{p,q}[0,T]} + \sup_{t \in [0,T]} \left\| \left(\frac{1}{\varrho_\varepsilon}, \frac{1}{\vartheta_\varepsilon} \right) \right\|_{C(\overline{\Omega})} \leq c(M) \text{ uniformly for } \varepsilon \rightarrow 0.$$

Consequently, passing to a new subsequence as the case may be, we may suppose

$$\begin{aligned} (\vartheta_\varepsilon, \mathbf{u}_\varepsilon) &\rightarrow (\vartheta, \mathbf{u}) \text{ weakly in } L^p(0, T; W^{2,q}(\Omega; R^4)) \cap W^{1,p}(0, T; L^q(\Omega; R^4)), \\ \varrho_\varepsilon &\rightarrow \varrho \text{ weakly-}^* \text{ in } L^\infty(0, T; W^{1,q}(\Omega)). \end{aligned}$$

In view of the strong convergence of the data postulated in (6.1), and the compact embeddings stated in Lemma 6.1, it is a routine matter to conclude that the triple $(\varrho, \vartheta, \mathbf{u})$ is the strong solution of the NSF system in $[0, T]$ claimed in Theorem 2.1.

6.2 Local existence in the mixed setting

Theorem 6.2 (Local existence - mixed framework). *Let*

$$3 < q \leq 6$$

be given. Let $\Omega \subset R^3$ be a bounded domain of class C^4 , and let $G = G(x)$, $G \in W^{2,2}(\Omega)$ be a given potential. Let the initial data $(\varrho_0, \vartheta_0, \mathbf{u}_0)$ satisfy (4.19), and let the boundary data $(\mathbf{u}_B, \vartheta_B)$ belong to the class (4.20), (4.21), respectively, where

$$\min_{\Omega} \varrho_0 \geq \underline{\varrho} > 0, \quad \min_{\Omega} \vartheta_0 \geq \underline{\vartheta} > 0, \quad \min_{\Gamma_D} \vartheta_B \geq \underline{\vartheta} > 0, \quad q_B \geq 0.$$

In addition, let the compatibility conditions

$$\begin{aligned} \mathbf{u}_0|_{\partial\Omega} &= \mathbf{u}_B, \\ \vartheta_0|_{\Gamma_D} &= \vartheta_B, \quad \nabla_x \vartheta_0 \cdot \mathbf{n}|_{\Gamma_N} = q_B \end{aligned}$$

hold. Finally, we suppose either $\Gamma_D \neq \emptyset$ or $q_B = 0$.

Then there exists $T > 0$ such that the NSF system (2.1)–(2.6), with the initial conditions (2.7), and the boundary conditions (2.8) admits a unique solution $(\varrho, \vartheta, \mathbf{u})$ in the class (4.15)–(4.17).

We omit the proof as it can be carried over in the same way as in the preceding section, with Proposition 5.1 in place of Proposition 5.4.

Remark 6.3. The conditional regularity estimates (5.3) established originally for smoother solutions extend by compactness/density arguments to the less regular solutions obtained in Theorem 6.2.

7 Conditional regularity in the $L^p - L^q$ framework

Our ultimate goal is to establish conditional regularity estimates in the $L^p - L^q$ framework and thus complete the proof of Theorem 2.1.

Theorem 7.1 (Conditional regularity - $L^p - L^q$ framework). *Let the hypotheses of Theorem 2.1 be satisfied.*

Then there exists a non-decreasing (positive) function $\Lambda : [0, \infty) \rightarrow (0, \infty)$ such that the solution $(\varrho, \vartheta, \mathbf{u})$ admits the estimate

$$\begin{aligned} & \|(\varrho, \vartheta, \mathbf{u})\|_{S_{p,q}[0,T]} + \sup_{t \in [0,T]} \left\| \left(\frac{1}{\varrho}, \frac{1}{\vartheta} \right) (t, \cdot) \right\|_{C(\bar{\Omega}; R^2)} \\ & \leq \Lambda \left(\|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{p,q,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{p,q,B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega}; R^2)} + \|G\|_{W^{1,q}(\Omega)} \right. \\ & \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{C([0,T] \times \bar{\Omega}; R^5)} \right). \end{aligned} \quad (7.1)$$

for any $0 \leq T < T_{\max,p,q}$.

Similarly to the above, $T_{\max,p,q}$ denotes the life span of the strong solution constructed in Section 6.1. As we shall see below, it is actually independent of the specific value of the exponents p, q .

The rest of this section is devoted to the proof of Theorem 7.1.

7.1 Estimates for more regular data

We start with the case of more regular data, specifically,

$$(\varrho_0, \vartheta_0, \mathbf{u}_0) \in \mathcal{D}_{p,q,I} \cap \mathcal{D}_{ChK,I}, \quad (\mathbf{u}_B, \vartheta_B, q_B) \in \mathcal{D}_{p,q,B} = \mathcal{D}_{ChK,B}, \quad G \in W^{2,q}(\Omega).$$

Accordingly, the solution satisfies the bound (5.3) from Proposition 5.1, cf. also Remark 6.3:

$$\begin{aligned} & \|(\varrho, \vartheta, \mathbf{u})\|_{S_{ChK}[0,T]} + \sup_{t \in [0,T]} \left\| \left(\frac{1}{\varrho}, \frac{1}{\vartheta} \right) (t, \cdot) \right\|_{C(\bar{\Omega}; R^2)} \\ & \leq \Lambda \left(T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{ChK,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{ChK,B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega}; R^2)} \right. \\ & \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0,T) \times \bar{\Omega}; R^5)} + \|G\|_{W^{2,q}(\Omega)} \right), \quad 3 < q \leq 6 \end{aligned} \quad (7.2)$$

for any $0 < T < T_{\max,ChK}$.

Now, revisiting the maximal regularity estimates (5.22), (5.23) we observe that

$$\begin{aligned} \|(\varrho, \vartheta, \mathbf{u})\|_{S_{p,q}[0,T]} & \leq \Lambda \left(\|(\varrho, \vartheta, \mathbf{u})\|_{S_{ChK}[0,T]} + \sup_{t \in [0,T]} \left\| \left(\frac{1}{\varrho}, \frac{1}{\vartheta} \right) (t, \cdot) \right\|_{C(\bar{\Omega}; R^2)} \right. \\ & \left. + T + \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{p,q,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{p,q,B}} + \|G\|_{W^{2,q}(\Omega)} \right). \end{aligned} \quad (7.3)$$

Unlike (7.2) that holds for a restricted range of q amenable to the mixed class, the estimate (7.3) can be extended to arbitrary p, q satisfying (2.14) by an induction argument.

Thus the conclusion of Theorem 7.1 holds for more regular initial data, specifically:

$$\begin{aligned}
& \|(\varrho, \vartheta, \mathbf{u})\|_{\mathcal{S}_{p,q}[0,T]} + \sup_{t \in [0,T]} \left\| \left(\frac{1}{\varrho}, \frac{1}{\vartheta} \right) (t, \cdot) \right\|_{C(\bar{\Omega}; R^2)} \\
& \leq \Lambda \left(\|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{p,q,I} \cap \mathcal{D}_{ChK,I}} + \|(\mathbf{u}_B, \vartheta_B, q_B)\|_{\mathcal{D}_{p,q,B}} + \left\| \left(\frac{1}{\varrho_0}, \frac{1}{\vartheta_0} \right) \right\|_{C(\bar{\Omega}; R^2)} \right. \\
& \left. + \|(\varrho, \vartheta, \mathbf{u})\|_{L^\infty((0,T) \times \bar{\Omega}; R^5)} + \|G\|_{W^{2,q}(\Omega)} \right). \tag{7.4}
\end{aligned}$$

7.2 Initial smoothing

In view of the bound (7.4), the proof of Theorem 7.1 is complete as soon as we observe that solutions emanating from the data in $\mathcal{D}_{p,q,I}$ will immediately become more regular, namely

$$(\varrho(\tau, \cdot), \vartheta(\tau, \cdot), \mathbf{u}(\tau, \cdot)) \in \mathcal{D}_{p,q,I} \cap \mathcal{D}_{ChK,I} \tag{7.5}$$

for any $\tau > 0$. This is obviously the case for the density, where the two classes coincide, while for \mathbf{u}, ϑ , we have

$$(\vartheta, \mathbf{u}) \in L^p(0, T; W^{2,q}(\Omega; R^4)).$$

As $q > 3$, the relation (7.5) holds for a.a. $\tau \in (0, T)$. Moreover, by the mean value theorem,

$$\|(\vartheta, \mathbf{u})(\xi, \cdot)\|_{W^{2,q}(\Omega; R^4)}^p = \frac{1}{T} \|(\vartheta, \mathbf{u})\|_{L^p(0,T; W^{2,q}(\Omega; R^4))}^p \text{ for some } \xi \in [0, T].$$

Consequently, the the proof of Theorem 2.1 is complete as soon as we establish the following result.

Lemma 7.2. *Consider a bounded set of initial/boundary data*

$$\mathcal{B} = \left\{ (\varrho_0, \vartheta_0, \mathbf{u}_0, \vartheta_B, \mathbf{u}_B, q_B) \mid \|(\varrho_0, \vartheta_0, \mathbf{u}_0)\|_{\mathcal{D}_{p,q,I}} + \|(\vartheta_B, \mathbf{u}_B, q_B)\|_{\mathcal{D}_{p,q,B}} \leq K \right\}.$$

Then there exists $T(K) > 0$ such that the local solutions $(\varrho, \vartheta, \mathbf{u})$ corresponding to the data in \mathcal{B} exist on $[0, T]$, and

$$\|(\vartheta, \mathbf{u})\|_{L^p(0,T; W^{2,q}(\Omega; R^4))}^p < \Lambda(K).$$

Proof. The arguments are similar to the existence part of the proof of Theorem 2.1. Given the data

$$(\varrho_0, \vartheta_0, \mathbf{u}_0, \vartheta_B, \mathbf{u}_B, q_B) \in \mathcal{B},$$

we set

$$\mathcal{F}[\varrho_0, \vartheta_0, \mathbf{u}_0, \vartheta_B, \mathbf{u}_B, q_B](\tau) = \|(\varrho, \vartheta, \mathbf{u})(\tau, \cdot)\|_{C(\bar{\Omega}; R^5)} + \int_0^\tau \|(\vartheta, \mathbf{u})(t, \cdot)\|_{W^{1,\infty}(\Omega; R^4)}^p dt,$$

where $(\varrho, \vartheta, \mathbf{u})$ is the corresponding strong solution.

Now, we fix T_{eff} as in (5.18),

$$T_{\text{eff}} = \Lambda^{-1}(K + M) > 0,$$

where K is the “radius” of the data set and $M = M(K) > 0$ is a given number chosen in such a way that

$$\sup \left\{ \mathcal{F}[\varrho_0, \vartheta_0, \mathbf{u}_0; \vartheta_B, \mathbf{u}_B, q_B](0) \mid (\varrho_0, \vartheta_0, \mathbf{u}_0, \vartheta_B, \mathbf{u}_B, q_B) \in \mathcal{B} \right\} \leq \frac{M}{2}. \quad (7.6)$$

Next, we define

$$T_M = T_M[\varrho_0, \vartheta_0, \mathbf{u}_0; \vartheta_B, \mathbf{u}_B, q_B] = \sup \left\{ 0 \leq \tau \leq T_{\text{eff}} \mid \mathcal{F}[\varrho_0, \vartheta_0, \mathbf{u}_0; \vartheta_B, \mathbf{u}_B, q_B](\tau) < M \right\}.$$

Note carefully that $T_M > 0$ because of (7.6). Finally, we set

$$T = T(K) = \inf \left\{ T_M[\varrho_0, \vartheta_0, \mathbf{u}_0; \vartheta_B, \mathbf{u}_B, q_B] \mid (\varrho_0, \vartheta_0, \mathbf{u}_0; \vartheta_B, \mathbf{u}_B, q_B) \in \mathcal{B} \right\}.$$

In view of the conditional regularity estimates (5.17) established in Proposition 5.4, the time T meets the conclusion of Lemma 7.2 as soon as we show $T > 0$.

Arguing by contradiction, we construct a sequence of data

$$(\varrho_0^\varepsilon, \vartheta_0^\varepsilon, \mathbf{u}_0^\varepsilon; \vartheta_B^\varepsilon, \mathbf{u}_B^\varepsilon, q_B^\varepsilon) \in \mathcal{B}$$

such that

$$T_M^\varepsilon = T_M[\varrho_0^\varepsilon, \vartheta_0^\varepsilon, \mathbf{u}_0^\varepsilon; \vartheta_B^\varepsilon, \mathbf{u}_B^\varepsilon, q_B^\varepsilon] \rightarrow 0 \text{ as } \varepsilon \rightarrow 0. \quad (7.7)$$

We consider the associated sequence

$$\mathcal{F}_\varepsilon(\tau) = \mathcal{F}[\varrho_0^\varepsilon, \vartheta_0^\varepsilon, \mathbf{u}_0^\varepsilon; \vartheta_B^\varepsilon, \mathbf{u}_B^\varepsilon, q_B^\varepsilon](\tau),$$

with

$$\overline{\mathcal{F}}_\varepsilon(\tau) = \mathcal{F}_\varepsilon(\tau \wedge T_M^\varepsilon).$$

Now, similarly to the existence part of the proof of Theorem 2.1, we use Lemma 6.1 to conclude

$$\overline{\mathcal{F}}_\varepsilon \rightarrow \mathcal{F} \in C[0, T_{\text{eff}}],$$

where

$$0 \leq \mathcal{F}(\tau) < M \text{ on some interval } [0, \alpha], \quad \alpha > 0.$$

This implies

$$0 \leq \overline{\mathcal{F}}_\varepsilon(\tau) = \mathcal{F}_\varepsilon(\tau) < M \text{ in } [0, \alpha],$$

meaning $T_M^\varepsilon > \alpha > 0$ in contrast with (7.7). □

Acknowledgement

This work was done while the authors were participating in the Oberwolfach Research Fellows Program at Mathematisches Forschungsinstitut Oberwolfach. They warmly thank the Institute for its kind hospitality and the excellent research environment it provided.

References

- [1] A. Abbatiello, D. Basarić, and N. Chaudhuri. On a blow-up criterion for the Navier-Stokes-Fourier system under general equations of state. *Archive Preprint Series*, 2023. **arxiv preprint No. 2310.12230** .
- [2] H. Amann. Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems. In *Function spaces, differential operators and nonlinear analysis (Friedrichroda, 1992)*, volume **133** of *Teubner-Texte Math.*, pages 9–126. Teubner, Stuttgart, 1993.
- [3] H. Amann. *Linear and quasilinear parabolic problems, I*. Birkhäuser Verlag, Basel, 1995.
- [4] H. Amann. Compact embeddings of vector-valued Sobolev and Besov spaces. volume 35(55), pages 161–177. 2000. Dedicated to the memory of Branko Najman.
- [5] D. Basarić, E. Feireisl, and H. Mizerová. Conditional regularity for the Navier-Stokes-Fourier system with Dirichlet boundary conditions. *J. Differential Equations*, **365**:359–378, 2023.
- [6] H. Brezis and P. Mironescu. Gagliardo-Nirenberg inequalities and non-inequalities: the full story. *Ann. Inst. H. Poincaré C Anal. Non Linéaire*, **35**(5):1355–1376, 2018.
- [7] Y. Cho and H. Kim. Existence results for viscous polytropic fluids with vacuum. *J. Differential Equations*, **228**(2):377–411, 2006.
- [8] R. Danchin. Local theory in critical spaces for compressible viscous and heat-conductive gases. *Comm. Partial Differential Equations*, **26**(7-8):1183–1233, 2001.
- [9] R. Danchin. On the solvability of the compressible Navier-Stokes system in bounded domains. *Nonlinearity*, **23**(2):383–407, 2010.
- [10] R. Denk. An introduction to maximal regularity for parabolic evolution equations. In *Nonlinear partial differential equations for future applications*, volume 346 of *Springer Proc. Math. Stat.*, pages 1–70. Springer, Singapore, [2021] ©2021.
- [11] R. Denk, M. Hieber, and J. Prüss. Optimal $L^p - L^q$ -estimates for parabolic boundary value problems with inhomogenous data. *Math. Z.*, **257**:193–224, 2007.
- [12] J. Fan, S. Jiang, and Y. Ou. A blow-up criterion for compressible viscous heat-conductive flows. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **27**(1):337–350, 2010.
- [13] D. Fang, R. Zi, and T. Zhang. A blow-up criterion for two dimensional compressible viscous heat-conductive flows. *Nonlinear Anal.*, **75**(6):3130–3141, 2012.
- [14] E. Feireisl, A. Novotný, and Y. Sun. A regularity criterion for the weak solutions to the Navier-Stokes-Fourier system. *Arch. Ration. Mech. Anal.*, **212**(1):219–239, 2014.
- [15] E. Feireisl, H. Wen, and C. Zhu. On Nash’s conjecture for models of viscous, compressible, and heat conducting fluids. *IM ASCR Prague, preprint No. IM 2022 6*, 2022.

- [16] D. Hoff. Global solutions of the Navier-Stokes equations for multidimensional compressible flow with discontinuous initial data. *J. Differential Equations*, **120**:215–254, 1995.
- [17] Y. Kagei and S. Kawashima. Local solvability of an initial boundary value problem for a quasilinear hyperbolic-parabolic system. *J. Hyperbolic Differ. Equ.*, **3**(2):195–232, 2006.
- [18] M. Kotschote. Strong solutions to the compressible non-isothermal Navier-Stokes equations. *Adv. Math. Sci. Appl.*, **22**(2):319–347, 2012.
- [19] N. V. Krylov. Parabolic equations with VMO coefficients in Sobolev spaces with mixed norms. *J. Funct. Anal.*, **250**(2):521–558, 2007.
- [20] A. Matsumura and T. Nishida. The initial value problem for the equations of motion of compressible and heat conductive fluids. *Comm. Math. Phys.*, **89**:445–464, 1983.
- [21] J. Nash. Continuity of solutions of parabolic and elliptic equations. *Amer. J. Math.*, **80**:931–954, 1958.
- [22] J. Nash. Le problème de cauchy pour les équations différentielles d’un fluide général. *Bull. Soc. Math. France*, **90**:487–497, 1962.
- [23] W. Schlag. Schauder and L^p estimates for parabolic systems via Campanato spaces. *Comm. Partial Differential Equations*, **21**(7-8):1141–1175, 1996.
- [24] V. A. Solonnikov. The solvability of the initial-boundary value problem for the equations of motion of a viscous compressible fluid. *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, **56**:128–142, 197, 1976. Investigations on linear operators and theory of functions, VI.
- [25] Y. Sun, C. Wang, and Z. Zhang. A Beale-Kato-Majda criterion for the 3-D compressible Navier-Stokes equations. *J. Math. Pures Appl.*, **95**(1):36–47, 2011.
- [26] A. Valli. A correction to the paper: “An existence theorem for compressible viscous fluids” [Ann. Mat. Pura Appl. (4) **130** (1982), 197–213; MR 83h:35112]. *Ann. Mat. Pura Appl. (4)*, **132**:399–400 (1983), 1982.
- [27] A. Valli. An existence theorem for compressible viscous fluids. *Ann. Mat. Pura Appl. (4)*, **130**:197–213, 1982.
- [28] A. Valli and M. Zajaczkowski. Navier-Stokes equations for compressible fluids: Global existence and qualitative properties of the solutions in the general case. *Commun. Math. Phys.*, **103**:259–296, 1986.